

Dimethylammonium tetrachlorido-ferrate(III) 18-crown-6 clathrate

Ping Ping Shi and Min Min Zhao*

Ordered Matter Science Research Center, College of Chemistry and Chemical Engineering, Southeast University, Nanjing 211189, People's Republic of China
Correspondence e-mail: zmmzyahfdzg@126.com


Received 20 May 2010; accepted 28 May 2010

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.006\text{ \AA}$; R factor = 0.052; wR factor = 0.128; data-to-parameter ratio = 22.6.

The reaction of dimethylamine hydrochloride, 18-crown-6 and ferric chloride in ethanol yields the title compound, $(\text{C}_2\text{H}_8\text{N})[\text{FeCl}_4]\cdot\text{C}_{12}\text{H}_{24}\text{O}_6$, which exhibits an unusual supramolecular structure. The protonated dimethylamine contains one NH_2^+ group, resulting in a 1:1 supramolecular rotator-stator structure $(\text{CH}_3-\text{NH}_2^+-\text{CH}_3)(18\text{-crown-6})$, through $\text{N}-\text{H}\cdots\text{O}$ hydrogen-bonding interactions between the ammonium group of the cation and the O atoms of the crown ether. In the crystal, all three components lie on a common crystallographic mirror plane normal to [010].

Related literature

For similar 18-crown-6 clathrates, see: Akutagawa *et al.* (2002); Fender *et al.* (2002). For the ferroelectric properties of these materials, see: Zhang *et al.* (2009); Ye *et al.* (2009).



Experimental

Crystal data

$(\text{C}_2\text{H}_8\text{N})[\text{FeCl}_4]\cdot\text{C}_{12}\text{H}_{24}\text{O}_6$	$V = 2448.1 (9)\text{ \AA}^3$
$M_r = 508.06$	$Z = 4$
Orthorhombic, $Pnma$	Mo $K\alpha$ radiation
$a = 9.3035 (19)\text{ \AA}$	$\mu = 1.08\text{ mm}^{-1}$
$b = 11.328 (2)\text{ \AA}$	$T = 293\text{ K}$
$c = 23.230 (5)\text{ \AA}$	$0.40 \times 0.30 \times 0.20\text{ mm}$

Data collection

Rigaku SCXmini diffractometer	23771 measured reflections
Absorption correction: multi-scan (<i>CrystalClear</i> ; Rigaku, 2005)	2940 independent reflections
$T_{\min} = 0.685$, $T_{\max} = 0.806$	1799 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.073$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.052$	130 parameters
$wR(F^2) = 0.128$	H-atom parameters constrained
$S = 0.99$	$\Delta\rho_{\max} = 0.33\text{ e \AA}^{-3}$
2940 reflections	$\Delta\rho_{\min} = -0.23\text{ e \AA}^{-3}$

Table 1

Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H1C···O2 ⁱ	0.90	2.03	2.867 (3)	155
Symmetry code: (i) x , $-y + \frac{1}{2}$, z .				

Data collection: *CrystalClear* (Rigaku, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *PRPKAPPA* (Ferguson, 1999).

The authors are grateful to the starter fund of Southeast University for financial support to buy the X-ray diffractometer.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BH2286).

References

- Akutagawa, T., Hashimoto, A., Nishihara, S., Hasegawa, T. & Nakamura, T. (2002). *J. Supramol. Chem.* **2**, 175–186.
- Fender, N. S., Kahwa, I. A. & Fronczek, F. R. (2002). *J. Solid State Chem.* **163**, 286–293.
- Ferguson, G. (1999). *PRPKAPPA*. University of Guelph, Canada.
- Rigaku (2005). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Ye, H.-Y., Fu, D.-W., Zhang, Y., Zhang, W., Xiong, R.-G. & Huang, S. D. (2009). *J. Am. Chem. Soc.* **131**, 42–43.
- Zhang, W., Cheng, L.-Z., Xiong, R.-G., Nakamura, T. & Huang, S. D. (2009). *J. Am. Chem. Soc.* **131**, 12544–12545.

supporting information

Acta Cryst. (2010). E66, m728 [doi:10.1107/S1600536810020192]

Dimethylammonium tetrachloroferrate(III) 18-crown-6 clathrate

Ping Ping Shi and Min Min Zhao

S1. Comment

There is currently a great deal of interest in crown ethers because of their ability to form non-covalent, H-bonding complexes with ammonium cations, both in solid and in solution (Akutagawa *et al.*, 2002; Fender *et al.*, 2002). Not only the size of the crown ether, but also the nature of the ammonium cation (NH_4^+ , RNH_3^+ , R_2NH_2^+ , etc.) can influence on the stoichiometry and stability of these host–guest complexes. The host molecules combine with the guest species by intermolecular interactions, and, if the host molecule contains some specific sites, it is easy to realise high selectivity in ion or molecular recognition. 18-Crown-6 has the highest affinity for ammonium cations RNH_3^+ . While most studies of 18-crown-6 and its derivatives invariably showed a 1:1 stoichiometry with RNH_3^+ cations, some structurally characterized complexes of crown ethers include R_2NH_2^+ cations.

The present study is a part of systematic investigation of ferroelectric, phase transitions materials (Ye *et al.*, 2009; Zhang *et al.*, 2009) that include metal-organic coordination compounds with organic ligands, or are related to the structures with both organic and inorganic building fragments. In the measured temperature range from 80 to 420 K (m.p. > 430 K), the temperature dependence of the relative permittivity at 1 MHz varied smoothly from 4.6 to 7.2 in the title compound. No dielectric anomaly has been observed. This suggests that this compound is not an actual ferroelectric, or that no distinct phase transition occurred within the probed temperature range.

The title compound is composed of cationic $[\text{C}_2\text{H}_8\text{N} \text{ (18-Crown-6)}]^+$ and one single anionic $[\text{FeCl}_4]^-$ complex (Fig. 1). Supramolecular rotators are assembled between protonated dimethylamine ($\text{CH}_3\text{—NH}_2\text{—CH}_3$) $^+$ and 18-crown-6 by hydrogen-bonding. The ammonium moieties of (NH_2^+) cations interact with two O atoms of the crown ether through two simple N—H \cdots O hydrogen bonds, forming a 1:1 supramolecular rotator-stator structure.

Supramolecular cation structure, $[\text{C}_2\text{H}_8\text{N} \text{ (18-Crown-6)}]^+$, were introduced as counter cations to $[\text{FeCl}_4]^-$ anions. The crown adopts a conformation in which the ring shows some distortion from the mean plane, with the torsion angles: C3—O2—C2—C1 = 178.2 (3) $^\circ$; C2—O2—C3—C4 = 72.6 (4) $^\circ$; C5—O3—C4—C3=178.9 (3) $^\circ$; C4—O3—C5—C6 = 179.5 (3) $^\circ$; O3—C4—C3—O2=61.0 (5) $^\circ$. Fe(III) has a flattened tetrahedral coordination by four Cl $^-$ ions [range of *cis*-bond angles: 108.14 (4)–111.67 (9) $^\circ$; $d_{\text{av}}(\text{Fe—Cl}) = 2.1728 (15)–2.1889 (12) \text{\AA}$].


Fig. 2 shows a view of the crystal structure down the *a* axis. An alternate arrangement of cations and anions layers is elongated along the *b* axis. The title compound is stabilized by intramolecular N—H \cdots O hydrogen bonds, but no significant intermolecular hydrogen bonds are observed.

S2. Experimental


$\text{CH}_3\text{—NH—CH}_3\text{HCl}$ (2 mmol, 0.163 g) and 18-crown-6 (2 mmol, 0.528 g) were dissolved in ethanol. Then, trivalent ferric chloride (2 mmol, 0.54 g) was added to the mixture in concentrated hydrochloric acid medium, the precipitate was filtered and washed with a small amount of ethanol. Five days later, single crystals suitable for X-ray diffraction analysis were obtained from slow evaporation of ethanol and DMF solution at room temperature.

S3. Refinement

All C-bonded H atoms were calculated geometrically with C—H distances fixed to 0.96 Å, and were allowed to ride on the C atoms to which they are bonded, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ or $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$ (methyl groups). The ammonium H atom (H1C) was calculated geometrically and refined using a riding model with N—H = 0.90 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{N})$.

**Figure 1**

The title compound, with the atomic numbering scheme. The displacement ellipsoids are drawn at the 30% probability level.

**Figure 2**

A view of the structure along the a axis. The dashed lines depict the hydrogen bonds.

Dimethylammonium tetrachloroferrate(III)-1,4,7,10,13,16-hexaoxacyclooctadecane (1/1)

Crystal data

$(\text{C}_2\text{H}_8\text{N})[\text{FeCl}_4] \cdot \text{C}_{12}\text{H}_{24}\text{O}_6$
 $M_r = 508.06$
Orthorhombic, $Pnma$
Hall symbol: -P 2ac 2n
 $a = 9.3035$ (19) Å
 $b = 11.328$ (2) Å
 $c = 23.230$ (5) Å
 $V = 2448.1$ (9) Å³
 $Z = 4$

$F(000) = 1060$
 $D_x = 1.378 \text{ Mg m}^{-3}$
Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 17071 reflections
 $\theta = 3.2\text{--}27.8^\circ$
 $\mu = 1.08 \text{ mm}^{-1}$
 $T = 293$ K
Prism, yellow
0.40 × 0.30 × 0.20 mm

Data collection

Rigaku SCXmini
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
Detector resolution: 13.6612 pixels mm⁻¹
CCD_Profile_fitting scans
Absorption correction: multi-scan
(*CrystalClear*; Rigaku, 2005)
 $T_{\min} = 0.685$, $T_{\max} = 0.806$

23771 measured reflections
2940 independent reflections
1799 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.073$
 $\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 3.2^\circ$
 $h = -12 \rightarrow 12$
 $k = -14 \rightarrow 14$
 $l = -30 \rightarrow 30$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.052$
 $wR(F^2) = 0.128$
 $S = 0.99$
2940 reflections

130 parameters
0 restraints
0 constraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.04P)^2 + 1.99P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.33 \text{ e \AA}^{-3}$$

$$\Delta\rho_{\min} = -0.23 \text{ e \AA}^{-3}$$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Fe1	0.47623 (7)	0.2500	0.38202 (3)	0.0633 (2)
Cl2	0.25400 (14)	0.2500	0.35240 (7)	0.0926 (5)
Cl1	0.61871 (16)	0.2500	0.30791 (6)	0.0890 (4)
Cl3	0.51101 (12)	0.09011 (13)	0.43307 (5)	0.1218 (5)
O1	-0.1251 (4)	0.2500	0.57334 (13)	0.0776 (10)
O2	0.0327 (3)	0.0373 (2)	0.58570 (10)	0.0804 (7)
O4	0.0916 (4)	0.2500	0.76754 (13)	0.0731 (9)
O3	0.1171 (3)	0.0390 (2)	0.70458 (11)	0.0814 (7)
C2	-0.1174 (4)	0.0408 (4)	0.57777 (2)	0.0991 (14)
H2A	-0.1646	0.0442	0.6144	0.119*
H2B	-0.1483	-0.0298	0.5584	0.119*
C4	0.0511 (5)	-0.0599 (3)	0.67999 (18)	0.0899 (12)
H4A	-0.0510	-0.0556	0.6856	0.108*
H4B	0.0856	-0.1305	0.6982	0.108*
C6	0.1585 (5)	0.1474 (4)	0.78905 (16)	0.0929 (12)
H6A	0.1503	0.1454	0.8302	0.111*
H6B	0.2587	0.1480	0.7793	0.111*
C3	0.0837 (5)	-0.0633 (3)	0.61760 (18)	0.0901 (12)
H3A	0.1859	-0.0689	0.6128	0.108*
H3B	0.0418	-0.1334	0.6014	0.108*
C1	-0.1581 (5)	0.1448 (4)	0.54302 (18)	0.1033 (15)
H1A	-0.1071	0.1438	0.5071	0.124*
H1B	-0.2592	0.1424	0.5348	0.124*
C5	0.0908 (5)	0.0448 (4)	0.76473 (16)	0.0910 (12)
H5A	0.1280	-0.0249	0.7830	0.109*
H5B	-0.0108	0.0478	0.7717	0.109*
N1	0.1943 (4)	0.2500	0.60266 (16)	0.0631 (10)
H1C	0.1395	0.3142	0.6088	0.076*
C8	0.3114 (6)	0.2500	0.6442 (2)	0.0885 (16)
H8A	0.2746	0.2500	0.6828	0.133*
H8B	0.3692	0.1808	0.6383	0.133*
C7	0.2423 (7)	0.2500	0.5428 (2)	0.107 (2)
H7A	0.1606	0.2500	0.5175	0.161*
H7B	0.2992	0.3192	0.5358	0.161*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Fe1	0.0520 (4)	0.0775 (5)	0.0602 (4)	0.000	-0.0032 (3)	0.000
Cl2	0.0602 (7)	0.0757 (9)	0.1419 (13)	0.000	-0.0276 (8)	0.000

Cl1	0.0893 (10)	0.1058 (11)	0.0719 (8)	0.000	0.0172 (7)	0.000
Cl3	0.0934 (8)	0.1508 (12)	0.1212 (9)	0.0227 (7)	0.0078 (6)	0.0725 (9)
O1	0.076 (2)	0.098 (3)	0.0589 (19)	0.000	-0.0134 (17)	0.000
O2	0.0754 (16)	0.0766 (17)	0.0892 (17)	-0.0121 (13)	0.0188 (13)	-0.0222 (14)
O4	0.080 (2)	0.076 (2)	0.064 (2)	0.000	-0.0125 (17)	0.000
O3	0.0913 (18)	0.0591 (15)	0.0937 (18)	0.0036 (13)	0.0070 (14)	-0.0007 (13)
C2	0.080 (3)	0.098 (3)	0.120 (4)	-0.019 (2)	0.008 (2)	-0.045 (3)
C4	0.097 (3)	0.058 (2)	0.115 (3)	0.007 (2)	0.017 (2)	-0.001 (2)
C6	0.092 (3)	0.113 (3)	0.074 (2)	0.018 (3)	-0.015 (2)	0.019 (2)
C3	0.103 (3)	0.050 (2)	0.117 (3)	0.003 (2)	0.031 (3)	-0.013 (2)
C1	0.082 (3)	0.145 (4)	0.083 (3)	-0.005 (3)	-0.020 (2)	-0.034 (3)
C5	0.106 (3)	0.082 (3)	0.085 (3)	0.012 (2)	-0.007 (2)	0.022 (2)
N1	0.050 (2)	0.060 (2)	0.080 (3)	0.000	0.0050 (19)	0.000
C8	0.066 (3)	0.114 (5)	0.086 (4)	0.000	-0.009 (3)	0.000
C7	0.113 (5)	0.141 (6)	0.069 (4)	0.000	0.010 (3)	0.000

Geometric parameters (\AA , °)

Fe1—Cl1	2.1728 (15)	C4—H4B	0.9599
Fe1—Cl2	2.1791 (14)	C6—C5	1.437 (5)
Fe1—Cl3 ⁱ	2.1889 (12)	C6—H6A	0.9601
Fe1—Cl3	2.1889 (12)	C6—H6B	0.9600
O1—C1	1.418 (4)	C3—H3A	0.9600
O1—C1 ⁱ	1.418 (4)	C3—H3B	0.9601
O2—C2	1.410 (4)	C1—H1A	0.9599
O2—C3	1.440 (4)	C1—H1B	0.9600
O4—C6 ⁱ	1.410 (4)	C5—H5A	0.9600
O4—C6	1.410 (4)	C5—H5B	0.9600
O3—C4	1.399 (4)	N1—C8	1.455 (6)
O3—C5	1.420 (4)	N1—C7	1.461 (6)
C2—C1	1.476 (6)	N1—H1C	0.9000
C2—H2A	0.9601	C8—H8A	0.9599
C2—H2B	0.9600	C8—H8B	0.9601
C4—C3	1.481 (5)	C7—H7A	0.9600
C4—H4A	0.9600	C7—H7B	0.9600
Cl1—Fe1—Cl2	109.19 (7)	O2—C3—C4	114.5 (3)
Cl1—Fe1—Cl3 ⁱ	109.82 (4)	O2—C3—H3A	108.6
Cl2—Fe1—Cl3 ⁱ	108.14 (4)	C4—C3—H3A	108.6
Cl1—Fe1—Cl3	109.82 (4)	O2—C3—H3B	108.6
Cl2—Fe1—Cl3	108.14 (4)	C4—C3—H3B	108.7
Cl3 ⁱ —Fe1—Cl3	111.67 (9)	H3A—C3—H3B	107.6
C1—O1—C1 ⁱ	114.3 (4)	O1—C1—C2	110.1 (3)
C2—O2—C3	114.6 (3)	O1—C1—H1A	109.6
C6 ⁱ —O4—C6	111.1 (4)	C2—C1—H1A	109.7
C4—O3—C5	111.3 (3)	O1—C1—H1B	109.6
O2—C2—C1	110.4 (4)	C2—C1—H1B	109.6
O2—C2—H2A	109.7	H1A—C1—H1B	108.2

C1—C2—H2A	109.5	O3—C5—C6	110.4 (3)
O2—C2—H2B	109.6	O3—C5—H5A	109.6
C1—C2—H2B	109.5	C6—C5—H5A	109.4
H2A—C2—H2B	108.1	O3—C5—H5B	109.6
O3—C4—C3	109.3 (3)	C6—C5—H5B	109.6
O3—C4—H4A	109.8	H5A—C5—H5B	108.1
C3—C4—H4A	109.7	C8—N1—C7	113.7 (4)
O3—C4—H4B	109.9	C8—N1—H1C	108.6
C3—C4—H4B	109.9	C7—N1—H1C	109.0
H4A—C4—H4B	108.3	N1—C8—H8A	110.6
O4—C6—C5	109.5 (3)	N1—C8—H8B	108.9
O4—C6—H6A	109.7	H8A—C8—H8B	109.5
C5—C6—H6A	109.8	N1—C7—H7A	109.9
O4—C6—H6B	109.8	N1—C7—H7B	109.3
C5—C6—H6B	109.8	H7A—C7—H7B	109.5
H6A—C6—H6B	108.2		

Symmetry code: (i) $x, -y+1/2, z$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D\cdots H$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
N1—H1C \cdots O2 ⁱ	0.90	2.03	2.867 (3)	155

Symmetry code: (i) $x, -y+1/2, z$.