

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## 4,4'-Diphenyl-2,2'-bi-1,3-thiazole

Seik Weng Ng

Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia  
Correspondence e-mail: seikweng@um.edu.my


Received 8 July 2010; accepted 9 July 2010

Key indicators: single-crystal X-ray study;  $T = 100$  K; mean  $\sigma(\text{C}-\text{C}) = 0.002$  Å;  
 $R$  factor = 0.028;  $wR$  factor = 0.079; data-to-parameter ratio = 17.3.

In the centrosymmetric title compound,  $\text{C}_{18}\text{H}_{12}\text{N}_2\text{S}_2$ , the five- (r.m.s. deviation = 0.002 Å) and six-membered (r.m.s. deviation = 0.002 Å) rings are essentially coplanar [dihedral angle between rings = 1.9 (1)°].

## Related literature

For the crystal structures of other 4,4'-disubstituted compounds, see: Bolognesi *et al.* (1987); Craig *et al.* (1988); Curtis *et al.* (2004).



## Experimental

## Crystal data

 $\text{C}_{18}\text{H}_{12}\text{N}_2\text{S}_2$  $M_r = 320.42$ 

Monoclinic,  $P2_1/c$   
 $a = 5.7769$  (4) Å  
 $b = 7.6573$  (5) Å  
 $c = 17.1960$  (12) Å  
 $\beta = 99.614$  (1)°  
 $V = 749.99$  (9) Å<sup>3</sup>

$Z = 2$   
Mo  $K\alpha$  radiation  
 $\mu = 0.35$  mm<sup>-1</sup>  
 $T = 100$  K  
 $0.30 \times 0.10 \times 0.10$  mm

## Data collection

Bruker SMART APEX  
diffractometer  
Absorption correction: multi-scan  
(*SADABS*; Sheldrick, 1996)  
 $T_{\min} = 0.902$ ,  $T_{\max} = 0.966$

6993 measured reflections  
1730 independent reflections  
1575 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.026$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.028$   
 $wR(F^2) = 0.079$   
 $S = 1.04$   
1730 reflections

100 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.42$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.24$  e Å<sup>-3</sup>

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINTE* (Bruker, 2009); data reduction: *SAINTE*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *pubCIF* (Westrip, 2010).

I thank the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NK2046).

## References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.  
Bolognesi, A., Catellani, M., Destri, S. & Porzio, W. (1987). *Acta Cryst.* **C43**, 1171–1173.  
Bruker (2009). *APEX2* and *SAINTE*. Bruker AXS Inc., Madison, Wisconsin, USA.  
Craig, D. C., Goodwin, H. A., Onggo, D. & Rae, A. D. (1988). *Aust. J. Chem.* **41**, 1625–1644.  
Curtis, D., Cao, J. & Kampf, J. F. (2004). *J. Am. Chem. Soc.* **126**, 4318–4328.  
Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.  
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

## supporting information

*Acta Cryst.* (2010). E66, o2030 [https://doi.org/10.1107/S1600536810027261]

## 4,4'-Diphenyl-2,2'-bi-1,3-thiazole

Seik Weng Ng

### S1. Comment

2,2'-Bithiazole and other 4,4'-disubstituted derivatives possess a pair of nitrogen-donor sites that renders such molecules capable of chelating metal atoms. The crystal structure of the parent compound as well as those of the methyl and ethyl substituted derivatives have been reported (Bolognesi *et al.*, 1987; Craig *et al.*, 1988, Curtis *et al.*, 2004). These molecules are centrosymmetric compounds having an inversion center midway along the C<sub>azoly1</sub>–C<sub>azoly1</sub> bond. In the parent compound, this bond is 1.468 (6) Å (Bolognesi *et al.*, 1987). The bond is somewhat shortened to 1.455 (2) Å in the phenyl analog (Scheme I, Fig. 1).

### S2. Experimental

The organic compound was returned unchanged in an attempted reaction of lead(II) nitrate (0.13 mmol, 0.04 g) with 4,4'-diphenyl-2,2'-bithiazole (0.25 mmol, 0.08 g) in the presence of potassium thiocyanate (0.25 mmol, 0.03 g) in a methanol/THF mixture. Crystals were obtained after one week of setting the mixture aside.

### S3. Refinement

Hydrogen atoms were placed in calculated positions (C–H 0.95 Å) and included in the refinement in the riding model approximation, with  $U(\text{H})$  set to  $1.2U_{\text{eq}}(\text{C})$ .




Figure 1

Displacement ellipsoid plot (Barbour, 2001) of C<sub>18</sub>H<sub>12</sub>N<sub>2</sub>S<sub>2</sub> at the 70% probability level. Hydrogen atoms are drawn as spheres of arbitrary radius. The molecule lies on an inversion center.

## 4,4'-Diphenyl-2,2'-bi-1,3-thiazole

## Crystal data

C<sub>18</sub>H<sub>12</sub>N<sub>2</sub>S<sub>2</sub> $M_r = 320.42$ Monoclinic,  $P2_1/c$ 

Hall symbol: -P 2ybc

 $a = 5.7769$  (4) Å $b = 7.6573$  (5) Å $c = 17.1960$  (12) Å $\beta = 99.614$  (1)° $V = 749.99$  (9) Å<sup>3</sup> $Z = 2$  $F(000) = 332$  $D_x = 1.419$  Mg m<sup>-3</sup>Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 4091 reflections

 $\theta = 2.4$ – $28.3$ ° $\mu = 0.35$  mm<sup>-1</sup> $T = 100$  K

Prism, colorless

 $0.30 \times 0.10 \times 0.10$  mm

## Data collection

Bruker SMART APEX

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 $\omega$  scans

Absorption correction: multi-scan

(SADABS; Sheldrick, 1996)

 $T_{\min} = 0.902$ ,  $T_{\max} = 0.966$ 

6993 measured reflections

1730 independent reflections

1575 reflections with  $I > 2\sigma(I)$  $R_{\text{int}} = 0.026$  $\theta_{\text{max}} = 27.5$ °,  $\theta_{\text{min}} = 2.4$ ° $h = -7 \rightarrow 7$  $k = -9 \rightarrow 9$  $l = -21 \rightarrow 22$ 

## Refinement

Refinement on  $F^2$ 

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.028$  $wR(F^2) = 0.079$  $S = 1.04$ 

1730 reflections

100 parameters

0 restraints

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0421P)^2 + 0.350P]$ where  $P = (F_o^2 + 2F_c^2)/3$  $(\Delta/\sigma)_{\text{max}} = 0.001$  $\Delta\rho_{\text{max}} = 0.42$  e Å<sup>-3</sup> $\Delta\rho_{\text{min}} = -0.24$  e Å<sup>-3</sup>Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å<sup>2</sup>)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.76174 (5)	0.35670 (4)	0.446214 (17)	0.01797 (12)
N1	0.35546 (18)	0.49301 (13)	0.39798 (6)	0.0151 (2)
C1	0.2750 (2)	0.43006 (15)	0.25545 (7)	0.0139 (2)
C2	0.0586 (2)	0.51551 (16)	0.24862 (7)	0.0156 (2)
H2	0.0132	0.5690	0.2936	0.019*
C3	-0.0906 (2)	0.52260 (16)	0.17611 (7)	0.0179 (3)
H3	-0.2377	0.5804	0.1719	0.022*
C4	-0.0255 (2)	0.44574 (17)	0.10999 (7)	0.0198 (3)
H4	-0.1278	0.4505	0.0607	0.024*
C5	0.1903 (2)	0.36157 (16)	0.11614 (7)	0.0193 (3)
H5	0.2355	0.3094	0.0708	0.023*
C6	0.3399 (2)	0.35345 (15)	0.18827 (7)	0.0165 (3)
H6	0.4868	0.2957	0.1921	0.020*

C7	0.4293 (2)	0.41960 (15)	0.33292 (7)	0.0140 (2)
C8	0.6444 (2)	0.34026 (16)	0.34867 (7)	0.0167 (3)
H8	0.7186	0.2835	0.3103	0.020*
C9	0.5133 (2)	0.46895 (15)	0.46100 (7)	0.0150 (2)

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
S1	0.01731 (17)	0.02092 (18)	0.01515 (17)	0.00472 (11)	0.00112 (12)	-0.00100 (11)
N1	0.0165 (5)	0.0146 (5)	0.0145 (5)	-0.0003 (4)	0.0035 (4)	0.0003 (4)
C1	0.0159 (6)	0.0113 (5)	0.0149 (6)	-0.0023 (4)	0.0038 (4)	0.0007 (4)
C2	0.0166 (6)	0.0142 (6)	0.0167 (6)	-0.0010 (4)	0.0046 (5)	-0.0004 (4)
C3	0.0156 (6)	0.0155 (6)	0.0221 (6)	-0.0006 (4)	0.0014 (5)	0.0004 (5)
C4	0.0229 (7)	0.0179 (6)	0.0167 (6)	-0.0023 (5)	-0.0020 (5)	-0.0003 (5)
C5	0.0256 (7)	0.0179 (6)	0.0147 (6)	-0.0011 (5)	0.0044 (5)	-0.0030 (4)
C6	0.0180 (6)	0.0149 (6)	0.0172 (6)	0.0006 (4)	0.0040 (5)	-0.0007 (4)
C7	0.0168 (6)	0.0115 (5)	0.0145 (5)	-0.0017 (4)	0.0043 (4)	-0.0004 (4)
C8	0.0187 (6)	0.0176 (6)	0.0141 (5)	0.0006 (4)	0.0034 (4)	-0.0013 (4)
C9	0.0164 (6)	0.0134 (5)	0.0157 (6)	0.0001 (4)	0.0041 (4)	0.0001 (4)

*Geometric parameters (Å, °)*

S1—C8	1.7056 (12)	C3—H3	0.9500
S1—C9	1.7278 (12)	C4—C5	1.3914 (18)
N1—C9	1.3076 (15)	C4—H4	0.9500
N1—C7	1.3817 (15)	C5—C6	1.3897 (17)
C1—C2	1.3981 (16)	C5—H5	0.9500
C1—C6	1.4013 (16)	C6—H6	0.9500
C1—C7	1.4762 (16)	C7—C8	1.3690 (17)
C2—C3	1.3934 (17)	C8—H8	0.9500
C2—H2	0.9500	C9—C9 <sup>i</sup>	1.455 (2)
C3—C4	1.3870 (18)		
C8—S1—C9	88.74 (6)	C6—C5—H5	119.9
C9—N1—C7	110.30 (10)	C4—C5—H5	119.9
C2—C1—C6	119.02 (11)	C5—C6—C1	120.33 (11)
C2—C1—C7	119.86 (11)	C5—C6—H6	119.8
C6—C1—C7	121.11 (11)	C1—C6—H6	119.8
C3—C2—C1	120.28 (11)	C8—C7—N1	114.40 (11)
C3—C2—H2	119.9	C8—C7—C1	126.45 (11)
C1—C2—H2	119.9	N1—C7—C1	119.14 (10)
C4—C3—C2	120.33 (11)	C7—C8—S1	111.12 (9)
C4—C3—H3	119.8	C7—C8—H8	124.4
C2—C3—H3	119.8	S1—C8—H8	124.4
C3—C4—C5	119.76 (11)	N1—C9—C9 <sup>i</sup>	123.47 (14)
C3—C4—H4	120.1	N1—C9—S1	115.44 (9)
C5—C4—H4	120.1	C9 <sup>i</sup> —C9—S1	121.09 (12)
C6—C5—C4	120.27 (11)		

---

C6—C1—C2—C3	0.60 (17)	C6—C1—C7—C8	0.67 (18)
C7—C1—C2—C3	-178.53 (11)	C2—C1—C7—N1	0.83 (16)
C1—C2—C3—C4	-0.32 (18)	C6—C1—C7—N1	-178.29 (11)
C2—C3—C4—C5	-0.17 (18)	N1—C7—C8—S1	-0.35 (13)
C3—C4—C5—C6	0.36 (19)	C1—C7—C8—S1	-179.35 (9)
C4—C5—C6—C1	-0.06 (18)	C9—S1—C8—C7	0.34 (10)
C2—C1—C6—C5	-0.41 (17)	C7—N1—C9—C9 <sup>i</sup>	-179.94 (14)
C7—C1—C6—C5	178.71 (11)	C7—N1—C9—S1	0.14 (13)
C9—N1—C7—C8	0.13 (15)	C8—S1—C9—N1	-0.28 (10)
C9—N1—C7—C1	179.22 (10)	C8—S1—C9—C9 <sup>i</sup>	179.80 (14)
C2—C1—C7—C8	179.79 (12)		

---

Symmetry code: (i)  $-x+1, -y+1, -z+1$ .