

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Dibromidobis(*N,N,N',N'*-tetramethylthiourea- κ S)cadmium(II)Sidra Nawaz,^a Sana Sadaf,^a Mohammed Fettouhi,^b Atif Fazal^b and Saeed Ahmad^{a*}^aDepartment of Chemistry, University of Engineering and Technology, Lahore 54890, Pakistan, and ^bDepartment of Chemistry, King Fahd University of Petroleum and Minerals, Dhahran 31261, Saudi Arabia

Correspondence e-mail: saeed_a786@hotmail.com


Received 3 July 2010; accepted 14 July 2010

Key indicators: single-crystal X-ray study; $T = 292$ K; mean $\sigma(\text{N}-\text{C}) = 0.005$ Å; R factor = 0.021; wR factor = 0.052; data-to-parameter ratio = 25.9.

In the title compound, $[\text{CdBr}_2(\text{C}_5\text{H}_{12}\text{N}_2\text{S})_2]$, the Cd^{II} atom lies on a twofold rotation axis. It exhibits a distorted tetrahedral coordination environment defined by two S atoms of two tetramethylthiourea (tmtu) ligands and two bromide ions. The crystal structure is consolidated by $\text{C}-\text{H}\cdots\text{N}$ and $\text{C}-\text{H}\cdots\text{S}$ hydrogen bonds.

Related literature

For crystallographic and spectroscopic studies of thiourea complexes, see: Al-Arfaj *et al.* (1998); Ali *et al.* (2009); Isab *et al.* (2009); Lobana *et al.* (2008); Marcos *et al.* (1998); Moloto *et al.* (2003). The structure of the title compound is isotopic with $[\text{Cd}(\text{tmtu})_2\text{I}_2]$ (Nawaz *et al.*, 2010a) and $[\text{Hg}(\text{tmtu})_2\text{Cl}_2]$ (Nawaz *et al.*, 2010b).



Experimental

Crystal data

$[\text{CdBr}_2(\text{C}_5\text{H}_{12}\text{N}_2\text{S})_2]$
 $M_r = 536.67$
 Monoclinic, $C2/c$
 $a = 18.6133$ (17) Å

$b = 10.0690$ (9) Å
 $c = 13.4600$ (12) Å
 $\beta = 130.834$ (1)°
 $V = 1908.6$ (3) Å³

$Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 5.54$ mm⁻¹

$T = 292$ K
 $0.24 \times 0.23 \times 0.20$ mm

Data collection

Bruker SMART APEX area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{\text{min}} = 0.350$, $T_{\text{max}} = 0.404$

12678 measured reflections
 2379 independent reflections
 2114 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.028$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.021$
 $wR(F^2) = 0.052$
 $S = 1.05$
 2379 reflections

92 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.44$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.45$ e Å⁻³

Table 1

Selected bond lengths (Å).

Cd1—S1	2.5580 (6)	Cd1—Br1	2.5735 (3)
--------	------------	---------	------------

Table 2

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C2—H2A \cdots N2	0.96	2.53	2.855 (4)	100
C5—H5A \cdots S1	0.96	2.65	3.026 (3)	104

Data collection: SMART (Bruker, 2008); cell refinement: SAINT (Bruker, 2008); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

We gratefully acknowledge King Fahd University of Petroleum and Minerals, Dhahran, Saudi Arabia, for providing the X-ray facility.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: WM2371).

References

- Al-Arfaj, A. R., Reibenspies, J. H., Isab, A. A. & Hussain, M. S. (1998). *Acta Cryst.* **C54**, 51–53.
 Ali, S., Malik, M. R., Isab, A. A. & Ahmad, S. (2009). *J. Coord. Chem.* **62**, 475–480.
 Bruker (2008). SMART and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Isab, A. A., Wazeer, M. I. M. & Ashraf, W. (2009). *Spectrochim. Acta Part A*, **72**, 218–221.
 Lobana, T. S., Sharma, R., Sharma, R., Sultana, R. & Butcher, R. J. (2008). *Z. Anorg. Allg. Chem.* **634**, 718–723.
 Marcos, C., Alía, J. M., Adovasio, V., Prieto, M. & García-Granda, S. (1998). *Acta Cryst.* **C54**, 1225–1229.
 Moloto, M. J., Malik, M. A., O'Brien, P., Motevalli, M. & Kolawole, G. A. (2003). *Polyhedron*, **22**, 595–603.
 Nawaz, S., Sadaf, S., Fettouhi, M., Fazal, A. & Ahmad, S. (2010a). *Acta Cryst.* **E66**, m951.
 Nawaz, S., Sadaf, H., Fettouhi, M., Fazal, A. & Ahmad, S. (2010b). *Acta Cryst.* **E66**, m952.
 Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supporting information

Acta Cryst. (2010). E66, m950 [https://doi.org/10.1107/S1600536810028102]

Dibromidobis(*N,N,N',N'*-tetramethylthiourea- κ S)cadmium(II)**Sidra Nawaz, Sana Sadaf, Mohammed Fettouhi, Atif Fazal and Saeed Ahmad****S1. Comment**

The coordination chemistry of thioureas with metal ions has been the subject of several recent investigations because of their variable binding modes and because of the relevance of their binding sites to those in living systems.

Crystallographic reports about d^{10} metal complexes of thioureas established that these ligands are coordinated *via* the sulfur atom (Al-Arfaj *et al.*, 1998; Moloto *et al.*, 2003). Spectroscopic data is also consistent with this finding (Isab *et al.*, 2009; Ali *et al.*, 2009). Herein, we report the crystal structure of a cadmium bromide complex with tetramethylthiourea (tmtu), [Cd(C₅H₁₂N₂S₂)₂Br₂], (I).

The crystal structure of (I) consists of discrete molecular species in which the cadmium atom is located on a twofold rotation axis (Fig. 1). It exhibits a distorted tetrahedral coordination environment defined by two tetramethylthiourea (tmtu) ligands and two bromide ions. The tmtu ligand is terminally bound to the Cd^{II} atom *via* coordination of the S1 atom. The Cd—S and Cd—Br bond lengths are 2.5580 (6) and 2.5735 (3) Å, respectively. These values are in agreement with those reported for related compounds, e.g. (Al-Arfaj *et al.*, 1998; Lobana *et al.* 2008; Marcos *et al.*, 1998; Moloto *et al.*, 2003). The bond angles around Cd are indicative of a slight tetrahedral distortion, with the S—Cd—S angle showing the largest deviation (117.70 (3)°) from the ideal value. The SCN₂— moiety of the tmtu ligand is essentially planar, the maximum deviation from the mean plane being 0.007 (2) Å for the carbon atom. The fragments N1—C1—C2—C3 and N2—C1—C4—C5 are also close to planarity. The maximum deviations from the mean planes are 0.072 (2) Å and 0.065 (2) Å for N1 and N2, respectively. These values are consistent with a significant C—N double bond character and electron delocalization in the SCN₂— moiety. The steric effect of the two adjacent 1,3-methyl groups imposes a dihedral angle of 47.4 (2) ° for the two mean planes and therefore a tilted conformation. The latter is likely stabilized by non-classical intramolecular hydrogen bonding interactions involving methyl H atoms with sulfur and nitrogen atoms (C2—H2A \cdots N2 and C5—H5A \cdots S1). The molecules pack to form columns approximately parallel to [110] direction (Fig. 2).

The structure of (I) is isotopic with [Cd(tmtu)₂I₂] (Nawaz *et al.*, 2010*a*), with an equivalent degree of distortion from the ideal tetrahedral configuration and similar Cd—S and C—N bond lengths. The tmtu bond lengths are also consistent with those found for the likewise isotopic compound [Hg(tmtu)₂Cl₂] (Nawaz *et al.*, 2010*b*), in which a significantly higher distortion of the metal ion coordination sphere is observed.

S2. Experimental

0.35 g (1.0 mmol) cadmium(II) bromide tetrahydrate dissolved in 10 ml water were added to two equivalents of tetramethylthiourea in methanol. A white precipitate formed and was filtered off. The filtrate was kept for crystallization. As a result, an off-white crystalline product suitable for single crystal X-ray diffraction was obtained.

S3. Refinement

H atoms were placed in calculated positions with a C—H distance of 0.96 Å and $U_{\text{iso}}(\text{H}) = 1.5 U_{\text{eq}}(\text{C})$.




Figure 1

The molecular structure of title compound with the atomic numbering scheme. Displacement ellipsoids are drawn at the 30% probability level. H-atoms are omitted for clarity.




Figure 2

Packing diagram of the title complex

Dibromidobis(*N,N,N',N'*-tetramethylthiourea- κ S)cadmium(II)

Crystal data

[CdBr₂(C₅H₁₂N₂S)₂]
 $M_r = 536.67$

Monoclinic, $C2/c$
 Hall symbol: $-C 2yc$

$a = 18.6133 (17) \text{ \AA}$
 $b = 10.0690 (9) \text{ \AA}$
 $c = 13.4600 (12) \text{ \AA}$
 $\beta = 130.834 (1)^\circ$
 $V = 1908.6 (3) \text{ \AA}^3$
 $Z = 4$
 $F(000) = 1048$
 $D_x = 1.868 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
 Cell parameters from 12678 reflections
 $\theta = 2.5\text{--}28.3^\circ$
 $\mu = 5.54 \text{ mm}^{-1}$
 $T = 292 \text{ K}$
 Block, colorless
 $0.24 \times 0.23 \times 0.20 \text{ mm}$

Data collection

Bruker SMART APEX area-detector
 diffractometer
 Radiation source: normal-focus sealed tube
 Graphite monochromator
 ω scans
 Absorption correction: multi-scan
 (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.350, T_{\max} = 0.404$

12678 measured reflections
 2379 independent reflections
 2114 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.028$
 $\theta_{\max} = 28.3^\circ, \theta_{\min} = 2.5^\circ$
 $h = -24 \rightarrow 24$
 $k = -13 \rightarrow 13$
 $l = -17 \rightarrow 17$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.021$
 $wR(F^2) = 0.052$
 $S = 1.05$
 2379 reflections
 92 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods
 Secondary atom site location: difference Fourier
 map

Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.025P)^2 + 1.3001P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.44 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.45 \text{ e \AA}^{-3}$
 Extinction correction: SHELXTL (Sheldrick,
 2008), $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
 Extinction coefficient: 0.0068 (2)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cd1	1.0000	0.70441 (2)	0.2500	0.03969 (8)
Br1	1.149352 (17)	0.56647 (3)	0.34643 (3)	0.05549 (9)
S1	1.03619 (4)	0.83583 (7)	0.44089 (5)	0.04794 (14)
N1	0.91758 (14)	0.76405 (19)	0.4771 (2)	0.0474 (4)
N2	0.85096 (13)	0.8900 (2)	0.29223 (18)	0.0471 (4)
C1	0.92634 (14)	0.82923 (19)	0.39910 (19)	0.0366 (4)
C2	0.8527 (2)	0.8077 (3)	0.4968 (3)	0.0672 (7)

H2A	0.8267	0.8928	0.4562	0.101*
H2B	0.8867	0.8145	0.5891	0.101*
H2C	0.8022	0.7444	0.4580	0.101*
C3	0.9861 (2)	0.6637 (3)	0.5706 (3)	0.0724 (8)
H3A	1.0111	0.6189	0.5359	0.109*
H3B	0.9552	0.6004	0.5851	0.109*
H3C	1.0369	0.7056	0.6522	0.109*
C4	0.75337 (18)	0.8418 (3)	0.2186 (3)	0.0694 (8)
H4A	0.7550	0.7542	0.2480	0.104*
H4B	0.7211	0.8394	0.1266	0.104*
H4C	0.7203	0.9005	0.2330	0.104*
C5	0.8609 (2)	0.9921 (3)	0.2254 (3)	0.0697 (8)
H5A	0.9218	1.0340	0.2870	0.105*
H5B	0.8118	1.0573	0.1892	0.105*
H5C	0.8557	0.9524	0.1561	0.105*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cd1	0.04286 (13)	0.04283 (13)	0.04311 (13)	0.000	0.03237 (11)	0.000
Br1	0.05068 (15)	0.05784 (16)	0.05843 (16)	0.01407 (10)	0.03588 (13)	0.00638 (11)
S1	0.0391 (3)	0.0665 (4)	0.0425 (3)	-0.0082 (2)	0.0286 (2)	-0.0132 (2)
N1	0.0572 (11)	0.0471 (10)	0.0559 (11)	0.0041 (9)	0.0448 (10)	0.0036 (8)
N2	0.0448 (10)	0.0540 (11)	0.0427 (10)	0.0038 (8)	0.0287 (9)	-0.0020 (8)
C1	0.0406 (10)	0.0363 (10)	0.0371 (10)	-0.0014 (8)	0.0272 (9)	-0.0061 (8)
C2	0.0799 (19)	0.0800 (19)	0.0815 (19)	-0.0012 (15)	0.0702 (18)	-0.0047 (15)
C3	0.090 (2)	0.0600 (16)	0.082 (2)	0.0174 (15)	0.0627 (19)	0.0238 (15)
C4	0.0393 (13)	0.101 (2)	0.0573 (16)	0.0026 (14)	0.0268 (12)	-0.0123 (15)
C5	0.0804 (19)	0.0721 (18)	0.0571 (15)	0.0165 (15)	0.0451 (15)	0.0201 (14)

Geometric parameters (Å, °)

Cd1—S1	2.5580 (6)	C2—H2B	0.9600
Cd1—S1 ⁱ	2.5580 (6)	C2—H2C	0.9600
Cd1—Br1 ⁱ	2.5735 (3)	C3—H3A	0.9600
Cd1—Br1	2.5735 (3)	C3—H3B	0.9600
S1—C1	1.731 (2)	C3—H3C	0.9600
N1—C1	1.335 (3)	C4—H4A	0.9600
N1—C3	1.460 (3)	C4—H4B	0.9600
N1—C2	1.463 (3)	C4—H4C	0.9600
N2—C1	1.331 (3)	C5—H5A	0.9600
N2—C5	1.455 (3)	C5—H5B	0.9600
N2—C4	1.471 (3)	C5—H5C	0.9600
C2—H2A	0.9600		
S1—Cd1—S1 ⁱ	117.70 (3)	H2A—C2—H2C	109.5
S1—Cd1—Br1 ⁱ	105.899 (14)	H2B—C2—H2C	109.5
S1 ⁱ —Cd1—Br1 ⁱ	106.524 (15)	N1—C3—H3A	109.5

S1—Cd1—Br1	106.524 (15)	N1—C3—H3B	109.5
S1 ⁱ —Cd1—Br1	105.899 (14)	H3A—C3—H3B	109.5
Br1 ⁱ —Cd1—Br1	114.676 (17)	N1—C3—H3C	109.5
C1—S1—Cd1	100.04 (7)	H3A—C3—H3C	109.5
C1—N1—C3	122.1 (2)	H3B—C3—H3C	109.5
C1—N1—C2	122.3 (2)	N2—C4—H4A	109.5
C3—N1—C2	114.2 (2)	N2—C4—H4B	109.5
C1—N2—C5	121.5 (2)	H4A—C4—H4B	109.5
C1—N2—C4	122.8 (2)	N2—C4—H4C	109.5
C5—N2—C4	114.6 (2)	H4A—C4—H4C	109.5
N2—C1—N1	119.41 (19)	H4B—C4—H4C	109.5
N2—C1—S1	121.32 (16)	N2—C5—H5A	109.5
N1—C1—S1	119.26 (16)	N2—C5—H5B	109.5
N1—C2—H2A	109.5	H5A—C5—H5B	109.5
N1—C2—H2B	109.5	N2—C5—H5C	109.5
H2A—C2—H2B	109.5	H5A—C5—H5C	109.5
N1—C2—H2C	109.5	H5B—C5—H5C	109.5

Symmetry code: (i) $-x+2, y, -z+1/2$.

Hydrogen-bond geometry (Å, °)

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
C2—H2A...N2	0.96	2.53	2.855 (4)	100
C5—H5A...S1	0.96	2.65	3.026 (3)	104