

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

# 1-sec-Butyl-3-[hydroxy(1-methyl-1H-indol-3-yl)methylidene]pyrrolidine-2,4-dione

Hai-zhen Xu<sup>a\*</sup> and You-Quan Zhu<sup>b</sup>

<sup>a</sup>College of Chemistry, Tianjin Normal University, 393 Binshuixi Road, Xiqing District, Tianjin 300387, People's Republic of China, and <sup>b</sup>State Key Laboratory of Elemento-Organic Chemistry, Nankai University, Tianjin 300071, People's Republic of China

Correspondence e-mail: hxyxhz@126.com


Received 24 July 2010; accepted 1 August 2010

Key indicators: single-crystal X-ray study;  $T = 113$  K; mean  $\sigma(\text{C}-\text{C}) = 0.003$  Å; disorder in main residue;  $R$  factor = 0.055;  $wR$  factor = 0.168; data-to-parameter ratio = 17.0.

In the title compound,  $\text{C}_{18}\text{H}_{20}\text{N}_2\text{O}_3$ , the dihedral angle between the indole ring system (r.m.s. deviation = 0.018 Å) and the hydroxymethylenepyrrrolidine-2,4-dione plane (r.m.s. deviation = 0.036 Å) is  $9.87(7)^\circ$ . The keto and enol groups are involved in an intramolecular  $\text{O}-\text{H}\cdots\text{O}$  hydrogen bond. An intramolecular  $\text{C}-\text{H}\cdots\text{O}$  interaction also occurs. The *sec*-butyl group is disordered over two orientations corresponding to an approximate  $180^\circ$  rotation about the  $\text{N}-\text{C}$  bond, with occupancies of 0.670 (6) and 0.330 (6). In the crystal, molecules are linked into chains along the  $c$  axis by  $\text{C}-\text{H}\cdots\text{O}$  hydrogen bonds.

## Related literature

For the antibiotic activity of 3-acylpyrrolidine-2,4-dione compounds, see: Baan *et al.* (1978); Holzapfel *et al.* (1970); Mackellar *et al.* (1971); Matsuo *et al.* (1980); Rinehart *et al.* (1963); Sticking (1959); Wu *et al.* (2002). For a related structure, see: Ellis & Spek (2001). For bond-length data, see: Allen *et al.* (1987).



## Experimental

### Crystal data

 $\text{C}_{18}\text{H}_{20}\text{N}_2\text{O}_3$  $M_r = 312.36$ 

Monoclinic,  $P2_1/c$   
 $a = 11.781(2)$  Å  
 $b = 10.529(2)$  Å  
 $c = 12.644(3)$  Å  
 $\beta = 97.18(3)^\circ$   
 $V = 1556.1(5)$  Å<sup>3</sup>

$Z = 4$   
 Mo  $K\alpha$  radiation  
 $\mu = 0.09$  mm<sup>-1</sup>  
 $T = 113$  K  
 $0.18 \times 0.16 \times 0.10$  mm

### Data collection

Rigaku Saturn diffractometer  
 Absorption correction: multi-scan  
 (*CrystalClear*; Rigaku, 2005)  
 $T_{\min} = 0.984$ ,  $T_{\max} = 0.991$

12618 measured reflections  
 3698 independent reflections  
 2687 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.034$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.055$   
 $wR(F^2) = 0.168$   
 $S = 1.13$   
 3698 reflections  
 218 parameters

10 restraints  
 H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.66$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.57$  e Å<sup>-3</sup>

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O1}-\text{H1}\cdots\text{O3}$	0.84	1.72	2.5003 (19)	154
$\text{C8}-\text{H8}\cdots\text{O2}$	0.95	2.12	2.916 (2)	140
$\text{C9}-\text{H9C}\cdots\text{O2}^i$	0.98	2.51	3.441 (3)	159

Symmetry code: (i)  $x, -y + \frac{1}{2}, z - \frac{1}{2}$ .

Data collection: *CrystalClear* (Rigaku, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors gratefully acknowledge financial support from the National Natural Science Foundation of China (grant No. 20772066).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI5139).

## References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Baan, J. L. van der, Barnick, J. W. F. K. & Bickelhaupt, F. (1978). *Tetrahedron*, **34**, 223–231.
- Ellis, D. D. & Spek, A. L. (2001). *Acta Cryst.* **C57**, 433–434.
- Holzapfel, C. W., Hutchison, R. D. & Wilkins, D. C. (1970). *Tetrahedron*, **26**, 5239–5246.
- Mackellar, F. A., Grostic, M. F., Olson, E. C., Wnuk, R. J., Branfman, A. R. & Rinehart, K. L. Jr (1971). *J. Am. Chem. Soc.* **93**, 4943–4945.
- Matsuo, K., Kitaguchi, I., Takata, Y. & Tanaka, K. (1980). *Chem. Pharm. Bull.* **28**, 2494–2502.
- Rigaku (2005). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.
- Rinehart, K. L., Beck, J. R., Borders, D. B., Kinstle, T. H. & Krauss, D. (1963). *J. Am. Chem. Soc.* **85**, 4038–4039.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Sticking, C. E. (1959). *Biochem. J.* **72**, 332–334.
- Wu, C.-S., Huang, J.-L., Sun, Y.-S. & Yang, D.-Y. (2002). *J. Med. Chem.* **45**, 2222–2228.

## supporting information

*Acta Cryst.* (2010). E66, o2248 [https://doi.org/10.1107/S1600536810030679]

## 1-sec-Butyl-3-[hydroxy(1-methyl-1*H*-indol-3-yl)methylidene]pyrrolidine-2,4-dione

Hai-zhen Xu and You-Quan Zhu

### S1. Comment

Many compounds containing the 3-acylpyrrolidine-2,4-dione moiety are novel heterocyclic compounds with antibiotic activity. Some of them are tenuazonic (Sticking, 1959), streptolydigin (Rinehart *et al.*, 1963), tirandamycin (Mackellar *et al.*, 1971), malonomycin (Baan *et al.*, 1978), alpha-cyclopiazonic acid (Sticking, 1959) and bata-cyclopiazonic acid (Holzapfel *et al.*, 1970). All these compounds possess a 3-acyltetramic acid moiety as a tricarbonylmethane structure and their hydrogen chemical shift of the enol hydroxy is about 11 p.p.m. (Wu *et al.*, 2002). On the other hand, most of the excellent inhibitors of *p*-hydroxyphenylpyruvate dioxygenase also possess similar characteristics, which are crucial for their bioactivity. Up to now, we have synthesized a series of 3-(un)substituted aroyl-1-benzylpyrrolidine-2,4-dione compounds and some of them have high herbicidal activity. The structure of the title compound, (I), helps us to investigate the relationship between structure and herbicidal activity.

The molecular structure of (I) is shown in Fig. 1. Atom H1, involved in intramolecular hydrogen bonding between O1 and O3, was assigned to O1 rather than to O3, based on bond lengths. The C14—O3 distance is 1.254 (2) Å, which is longer than the C12—O2 distance of 1.231 (2) Å. In contrast, the C10—O1 distance [1.322 (2) Å] is intermediate between the normal carbonyl bond and the C—O single bond length (Allen *et al.* 1987). A similar situation has been found in 3-(1-hydroxyethylidene)-1-phenylpyrrolidine-2,4-dione, which contains the same pyrrolidine skeleton (Ellis & Spek, 2001). The dihedral angle formed by the indole ring system (r.m.s. deviation 0.018 Å) and the hydroxymethylene-pyrrolidine-2,4-dione plane (r.m.s. deviation 0.036 Å) is 9.87 (7)°.

### S2. Experimental

The title compound was obtained according to the reported procedure of Matsuo *et al.* (1980). Colourless single crystals were obtained by recrystallization of the title compound from petroleum ether and ethyl acetate.

### S3. Refinement

The sec-butyl group is disordered over two orientations corresponding to an approximately 180° rotation about the N2—C15 bond, with refined occupancies of 0.670 (6) and 0.330 (6). All C—C distances in this group were restrained to 1.540 (5) Å. H atoms were placed in calculated positions, with C—H = 0.95–0.98 Å and O—H = 0.84 Å, and included in the final cycles of refinement using a riding model, with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$  and  $1.5U_{\text{eq}}(\text{O})$ .




Figure 1

The molecular structure of the title compound, showing 30% probability displacement ellipsoids and the atom-numbering scheme. Both disorder components are shown.

### 1-sec-Butyl-3-[hydroxy(1-methyl-1*H*-indol-3-yl)methylidene]pyrrolidine-2,4-dione

#### Crystal data

$C_{18}H_{20}N_2O_3$   
 $M_r = 312.36$   
 Monoclinic,  $P2_1/c$   
 Hall symbol: -P 2ybc  
 $a = 11.781$  (2) Å  
 $b = 10.529$  (2) Å  
 $c = 12.644$  (3) Å  
 $\beta = 97.18$  (3)°  
 $V = 1556.1$  (5) Å<sup>3</sup>  
 $Z = 4$

$F(000) = 664$   
 $D_x = 1.333$  Mg m<sup>-3</sup>  
 Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å  
 Cell parameters from 4289 reflections  
 $\theta = 2.5$ – $27.9$ °  
 $\mu = 0.09$  mm<sup>-1</sup>  
 $T = 113$  K  
 Prism, yellow  
 $0.18 \times 0.16 \times 0.10$  mm

#### Data collection

Rigaku Saturn  
 diffractometer  
 Radiation source: fine-focus sealed tube  
 Graphite monochromator  
 $\omega$  scans  
 Absorption correction: multi-scan  
 (*CrystalClear*; Rigaku, 2005)  
 $T_{\min} = 0.984$ ,  $T_{\max} = 0.991$

12618 measured reflections  
 3698 independent reflections  
 2687 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.034$   
 $\theta_{\max} = 27.9$ °,  $\theta_{\min} = 2.6$ °  
 $h = -9 \rightarrow 15$   
 $k = -13 \rightarrow 13$   
 $l = -16 \rightarrow 16$

#### Refinement

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.055$   
 $wR(F^2) = 0.168$   
 $S = 1.13$   
 3698 reflections  
 218 parameters  
 10 restraints  
 Primary atom site location: structure-invariant  
 direct methods

Secondary atom site location: difference Fourier  
 map  
 Hydrogen site location: inferred from  
 neighbouring sites  
 H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0947P)^2 + 0.1812P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} = 0.001$   
 $\Delta\rho_{\max} = 0.66$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.57$  e Å<sup>-3</sup>

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
O1	0.66886 (10)	1.07272 (11)	0.42060 (10)	0.0232 (3)	
H1	0.6268	1.0643	0.4691	0.035*	
O2	0.93397 (11)	0.78344 (12)	0.54741 (11)	0.0297 (3)	
O3	0.59380 (11)	1.00771 (11)	0.58894 (10)	0.0258 (3)	
N1	0.98265 (12)	0.96858 (14)	0.26733 (12)	0.0224 (3)	
N2	0.69798 (14)	0.85187 (15)	0.68474 (13)	0.0282 (4)	
C1	0.83455 (14)	1.00283 (15)	0.35989 (13)	0.0194 (4)	
C2	0.81236 (14)	1.06960 (15)	0.25917 (13)	0.0202 (4)	
C3	0.72166 (16)	1.14252 (16)	0.20819 (15)	0.0253 (4)	
H3	0.6560	1.1595	0.2425	0.030*	
C4	0.72960 (18)	1.18928 (18)	0.10697 (15)	0.0306 (4)	
H4	0.6677	1.2368	0.0714	0.037*	
C5	0.82620 (19)	1.16822 (18)	0.05608 (16)	0.0319 (5)	
H5	0.8299	1.2041	-0.0123	0.038*	
C6	0.91688 (17)	1.09601 (17)	0.10329 (15)	0.0278 (4)	
H6	0.9827	1.0810	0.0687	0.033*	
C7	0.90694 (15)	1.04631 (16)	0.20424 (14)	0.0216 (4)	
C8	0.93954 (14)	0.94263 (16)	0.35943 (14)	0.0211 (4)	
H8	0.9757	0.8910	0.4154	0.025*	
C9	1.08481 (15)	0.91093 (18)	0.23351 (16)	0.0285 (4)	
H9A	1.1285	0.8681	0.2943	0.043*	
H9B	1.1322	0.9770	0.2065	0.043*	
H9C	1.0623	0.8489	0.1770	0.043*	
C10	0.75981 (14)	0.99897 (15)	0.44100 (13)	0.0186 (4)	
C11	0.76929 (14)	0.92599 (15)	0.53493 (14)	0.0201 (4)	
C12	0.84678 (15)	0.82736 (17)	0.57785 (14)	0.0236 (4)	
C13	0.80043 (16)	0.77699 (18)	0.67725 (16)	0.0298 (4)	
H13A	0.7816	0.6855	0.6696	0.036*	
H13B	0.8569	0.7890	0.7413	0.036*	
C14	0.67860 (15)	0.93502 (16)	0.60328 (14)	0.0220 (4)	
C15	0.61807 (18)	0.8307 (2)	0.76310 (17)	0.0412 (6)	
H15	0.5520	0.8898	0.7455	0.049*	
C16	0.6765 (2)	0.8655 (4)	0.87609 (19)	0.0717 (10)	
H16A	0.7046	0.9511	0.8762	0.108*	0.670 (6)
H16B	0.6218	0.8582	0.9260	0.108*	0.670 (6)

H16C	0.7391	0.8085	0.8962	0.108*	0.670 (6)
H16D	0.7043	0.9511	0.8756	0.086*	0.330 (6)
H16E	0.7411	0.8106	0.8945	0.086*	0.330 (6)
C17	0.5719 (2)	0.6974 (2)	0.7563 (2)	0.0582 (8)	
H17A	0.5187	0.6874	0.8073	0.070*	0.670 (6)
H17B	0.6336	0.6388	0.7749	0.070*	0.670 (6)
H17C	0.5373	0.6820	0.6846	0.087*	0.330 (6)
H17D	0.6328	0.6378	0.7745	0.087*	0.330 (6)
H17E	0.5155	0.6875	0.8043	0.087*	0.330 (6)
C18	0.5146 (3)	0.6638 (3)	0.6507 (3)	0.0524 (13)	0.670 (6)
H18A	0.4453	0.7153	0.6346	0.079*	0.670 (6)
H18B	0.4939	0.5736	0.6495	0.079*	0.670 (6)
H18C	0.5662	0.6800	0.5971	0.079*	0.670 (6)
C18'	0.5954 (8)	0.8519 (12)	0.9604 (7)	0.087 (4)	0.330 (6)
H18D	0.5255	0.9007	0.9387	0.130*	0.330 (6)
H18E	0.6326	0.8840	1.0287	0.130*	0.330 (6)
H18F	0.5758	0.7621	0.9678	0.130*	0.330 (6)

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O1	0.0214 (6)	0.0279 (7)	0.0218 (7)	0.0049 (5)	0.0084 (5)	0.0028 (5)
O2	0.0252 (7)	0.0340 (7)	0.0313 (8)	0.0094 (6)	0.0096 (6)	0.0056 (6)
O3	0.0230 (6)	0.0314 (7)	0.0240 (7)	0.0073 (5)	0.0073 (5)	0.0040 (5)
N1	0.0202 (7)	0.0254 (7)	0.0229 (8)	-0.0016 (6)	0.0077 (6)	-0.0021 (6)
N2	0.0258 (8)	0.0336 (8)	0.0275 (8)	0.0063 (7)	0.0119 (7)	0.0098 (6)
C1	0.0201 (8)	0.0194 (8)	0.0193 (8)	-0.0042 (6)	0.0051 (7)	-0.0015 (6)
C2	0.0228 (8)	0.0190 (8)	0.0198 (8)	-0.0036 (6)	0.0063 (7)	-0.0012 (6)
C3	0.0251 (9)	0.0254 (9)	0.0268 (10)	-0.0003 (7)	0.0085 (8)	0.0016 (7)
C4	0.0356 (10)	0.0301 (9)	0.0265 (10)	0.0024 (8)	0.0062 (8)	0.0053 (8)
C5	0.0441 (12)	0.0315 (10)	0.0222 (9)	0.0004 (9)	0.0123 (9)	0.0059 (8)
C6	0.0333 (10)	0.0275 (9)	0.0248 (9)	-0.0033 (8)	0.0124 (8)	-0.0010 (7)
C7	0.0235 (8)	0.0198 (8)	0.0227 (9)	-0.0035 (7)	0.0072 (7)	-0.0025 (6)
C8	0.0203 (8)	0.0242 (8)	0.0194 (9)	-0.0030 (7)	0.0049 (7)	-0.0021 (6)
C9	0.0219 (9)	0.0339 (10)	0.0318 (10)	0.0016 (8)	0.0120 (8)	-0.0040 (8)
C10	0.0181 (8)	0.0182 (8)	0.0200 (9)	-0.0017 (6)	0.0039 (6)	-0.0027 (6)
C11	0.0196 (8)	0.0207 (8)	0.0205 (9)	-0.0002 (6)	0.0051 (7)	0.0003 (6)
C12	0.0233 (9)	0.0260 (9)	0.0222 (9)	-0.0014 (7)	0.0058 (7)	0.0010 (7)
C13	0.0282 (9)	0.0324 (10)	0.0304 (10)	0.0071 (8)	0.0103 (8)	0.0106 (8)
C14	0.0219 (8)	0.0247 (8)	0.0198 (9)	-0.0015 (7)	0.0046 (7)	0.0006 (7)
C15	0.0355 (11)	0.0539 (13)	0.0387 (12)	0.0133 (10)	0.0218 (10)	0.0209 (10)
C16	0.0500 (16)	0.129 (3)	0.0391 (15)	0.0170 (17)	0.0182 (13)	0.0283 (16)
C17	0.0434 (13)	0.0516 (14)	0.086 (2)	0.0139 (11)	0.0349 (15)	0.0354 (14)
C18	0.042 (2)	0.0387 (19)	0.080 (3)	-0.0026 (16)	0.020 (2)	0.0083 (18)
C18'	0.079 (6)	0.126 (8)	0.058 (6)	0.012 (6)	0.021 (5)	0.010 (5)

*Geometric parameters (Å, °)*

O1—C10	1.322 (2)	C10—C11	1.407 (2)
O1—H1	0.84	C11—C12	1.443 (2)
O2—C12	1.231 (2)	C11—C14	1.459 (2)
O3—C14	1.254 (2)	C12—C13	1.527 (2)
N1—C8	1.355 (2)	C13—H13A	0.99
N1—C7	1.387 (2)	C13—H13B	0.99
N1—C9	1.459 (2)	C15—C17	1.504 (3)
N2—C14	1.349 (2)	C15—C16	1.550 (3)
N2—C13	1.455 (2)	C15—H15	1.00
N2—C15	1.467 (2)	C16—C18'	1.525 (5)
C1—C8	1.390 (2)	C16—H16A	0.96
C1—C10	1.433 (2)	C16—H16B	0.96
C1—C2	1.450 (2)	C16—H16C	0.96
C2—C3	1.405 (2)	C16—H16D	0.96
C2—C7	1.406 (2)	C16—H16E	0.96
C3—C4	1.385 (3)	C17—C18	1.462 (4)
C3—H3	0.95	C17—H17A	0.96
C4—C5	1.393 (3)	C17—H17B	0.96
C4—H4	0.95	C17—H17C	0.96
C5—C6	1.384 (3)	C17—H17D	0.96
C5—H5	0.95	C17—H17E	0.96
C6—C7	1.398 (3)	C18—H18A	0.98
C6—H6	0.95	C18—H18B	0.98
C8—H8	0.95	C18—H18C	0.98
C9—H9A	0.98	C18'—H18D	0.98
C9—H9B	0.98	C18'—H18E	0.98
C9—H9C	0.98	C18'—H18F	0.98
C10—O1—H1	109.5	N2—C15—H15	107.6
C8—N1—C7	109.27 (14)	C17—C15—H15	107.6
C8—N1—C9	125.41 (16)	C16—C15—H15	107.6
C7—N1—C9	124.87 (15)	C18'—C16—C15	112.1 (5)
C14—N2—C13	111.38 (14)	C18'—C16—H16A	109.5
C14—N2—C15	123.56 (16)	C15—C16—H16A	109.6
C13—N2—C15	124.71 (15)	C15—C16—H16B	109.2
C8—C1—C10	128.05 (16)	H16A—C16—H16B	109.5
C8—C1—C2	106.28 (14)	C18'—C16—H16C	106.5
C10—C1—C2	125.66 (15)	C15—C16—H16C	109.6
C3—C2—C7	118.22 (15)	H16A—C16—H16C	109.5
C3—C2—C1	135.34 (15)	H16B—C16—H16C	109.5
C7—C2—C1	106.40 (15)	C18'—C16—H16D	109.7
C4—C3—C2	118.86 (17)	C15—C16—H16D	109.2
C4—C3—H3	120.6	H16B—C16—H16D	109.7
C2—C3—H3	120.6	H16C—C16—H16D	109.7
C3—C4—C5	121.56 (19)	C18'—C16—H16E	108.7
C3—C4—H4	119.2	C15—C16—H16E	109.2

C5—C4—H4	119.2	H16A—C16—H16E	107.6
C6—C5—C4	121.27 (17)	H16B—C16—H16E	111.7
C6—C5—H5	119.4	H16D—C16—H16E	107.8
C4—C5—H5	119.4	C18—C17—C15	113.6 (2)
C5—C6—C7	116.82 (17)	C18—C17—H17A	108.8
C5—C6—H6	121.6	C15—C17—H17A	109.0
C7—C6—H6	121.6	C18—C17—H17B	108.5
N1—C7—C6	128.67 (16)	C15—C17—H17B	109.1
N1—C7—C2	108.13 (14)	H17A—C17—H17B	107.6
C6—C7—C2	123.20 (17)	C15—C17—H17C	108.7
N1—C8—C1	109.93 (16)	H17A—C17—H17C	112.4
N1—C8—H8	125.0	H17B—C17—H17C	110.0
C1—C8—H8	125.0	C18—C17—H17D	107.9
N1—C9—H9A	109.5	C15—C17—H17D	110.0
N1—C9—H9B	109.5	H17A—C17—H17D	107.3
H9A—C9—H9B	109.5	H17C—C17—H17D	109.5
N1—C9—H9C	109.5	C18—C17—H17E	105.9
H9A—C9—H9C	109.5	C15—C17—H17E	109.8
H9B—C9—H9C	109.5	H17B—C17—H17E	109.7
O1—C10—C11	117.53 (14)	H17C—C17—H17E	109.5
O1—C10—C1	113.52 (15)	H17D—C17—H17E	109.5
C11—C10—C1	128.93 (15)	C17—C18—H18A	109.5
C10—C11—C12	133.64 (15)	H17C—C18—H18A	107.5
C10—C11—C14	118.53 (15)	C17—C18—H18B	109.5
C12—C11—C14	107.45 (14)	H17C—C18—H18B	118.4
O2—C12—C11	131.76 (16)	H18A—C18—H18B	109.5
O2—C12—C13	121.67 (16)	C17—C18—H18C	109.5
C11—C12—C13	106.56 (14)	H17C—C18—H18C	102.1
N2—C13—C12	104.50 (14)	H18A—C18—H18C	109.5
N2—C13—H13A	110.9	H18B—C18—H18C	109.5
C12—C13—H13A	110.9	C16—C18'—H18D	109.5
N2—C13—H13B	110.9	H16B—C18'—H18D	104.2
C12—C13—H13B	110.9	C16—C18'—H18E	109.5
H13A—C13—H13B	108.9	H16B—C18'—H18E	113.9
O3—C14—N2	124.21 (16)	H18D—C18'—H18E	109.5
O3—C14—C11	125.70 (16)	C16—C18'—H18F	109.5
N2—C14—C11	110.08 (15)	H16B—C18'—H18F	110.2
N2—C15—C17	111.19 (18)	H18D—C18'—H18F	109.5
N2—C15—C16	109.75 (18)	H18E—C18'—H18F	109.5
C17—C15—C16	112.9 (2)		
C8—C1—C2—C3	176.72 (18)	C1—C10—C11—C12	-5.6 (3)
C10—C1—C2—C3	-1.9 (3)	O1—C10—C11—C14	0.5 (2)
C8—C1—C2—C7	-0.62 (18)	C1—C10—C11—C14	-177.42 (16)
C10—C1—C2—C7	-179.19 (15)	C10—C11—C12—O2	5.9 (3)
C7—C2—C3—C4	-0.7 (3)	C14—C11—C12—O2	178.41 (19)
C1—C2—C3—C4	-177.76 (18)	C10—C11—C12—C13	-173.30 (18)
C2—C3—C4—C5	-1.6 (3)	C14—C11—C12—C13	-0.80 (19)

C3—C4—C5—C6	2.1 (3)	C14—N2—C13—C12	1.2 (2)
C4—C5—C6—C7	-0.4 (3)	C15—N2—C13—C12	174.59 (18)
C8—N1—C7—C6	180.00 (18)	O2—C12—C13—N2	-179.49 (17)
C9—N1—C7—C6	-7.3 (3)	C11—C12—C13—N2	-0.2 (2)
C8—N1—C7—C2	-0.17 (19)	C13—N2—C14—O3	177.45 (17)
C9—N1—C7—C2	172.49 (15)	C15—N2—C14—O3	4.0 (3)
C5—C6—C7—N1	177.90 (17)	C13—N2—C14—C11	-1.8 (2)
C5—C6—C7—C2	-1.9 (3)	C15—N2—C14—C11	-175.24 (17)
C3—C2—C7—N1	-177.39 (15)	C10—C11—C14—O3	-3.8 (3)
C1—C2—C7—N1	0.49 (18)	C12—C11—C14—O3	-177.61 (17)
C3—C2—C7—C6	2.5 (3)	C10—C11—C14—N2	175.43 (15)
C1—C2—C7—C6	-179.67 (16)	C12—C11—C14—N2	1.6 (2)
C7—N1—C8—C1	-0.24 (19)	C14—N2—C15—C17	115.6 (2)
C9—N1—C8—C1	-172.85 (15)	C13—N2—C15—C17	-57.0 (3)
C10—C1—C8—N1	179.06 (16)	C14—N2—C15—C16	-118.7 (2)
C2—C1—C8—N1	0.53 (18)	C13—N2—C15—C16	68.7 (3)
C8—C1—C10—O1	176.19 (15)	N2—C15—C16—C18'	177.0 (5)
C2—C1—C10—O1	-5.5 (2)	C17—C15—C16—C18'	-58.3 (6)
C8—C1—C10—C11	-5.8 (3)	N2—C15—C17—C18	-57.2 (3)
C2—C1—C10—C11	172.44 (16)	C16—C15—C17—C18	178.9 (2)
O1—C10—C11—C12	172.36 (17)		

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
O1—H1...O3	0.84	1.72	2.5003 (19)	154
C8—H8...O2	0.95	2.12	2.916 (2)	140
C9—H9C...O2 <sup>i</sup>	0.98	2.51	3.441 (3)	159

Symmetry code: (i) *x*, -*y*+3/2, *z*-1/2.