

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

[Amino(iminio)methyl]phosphonate

Ting-Hai Yang,^{a*} Wei Zhuang,^a Wei Wei,^b Yong-Bing Yang^a and Qiang Chen^a^aHigh-Tech Institute of Nanjing University, Changzhou 213164, People's Republic of China, and ^bSchool of Chemistry & Chemical Engineering, Changzhou University, Changzhou 213164, People's Republic of China

Correspondence e-mail: Tinghaiyang@gmail.com, Chem100@nju.edu.cn


Received 30 July 2010; accepted 10 August 2010

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{N}-\text{C}) = 0.005$ Å; R factor = 0.034; wR factor = 0.161; data-to-parameter ratio = 13.4.

The title compound, $\text{CH}_5\text{N}_2\text{O}_3\text{P}$, exists as a zwitterion. The N atom of the imino group is protonated and the phosphonic acid group is deprotonated. The molecular geometry about the central C atom of this zwitterionic species was found to be strictly planar with the sum of the three angles about C being precisely 360° . In the crystal, the molecules are interlinked by $\text{O}-\text{H}\cdots\text{O}$ and $\text{N}-\text{H}\cdots\text{O}$ hydrogen-bonding interactions, forming a three-dimensional supramolecular network structure.

Related literature

For background to phosphonic acid and metal phosphonate compounds, see: Ayyappan *et al.* (2001); Clearfield (1998); Haga *et al.* (2007); Vivani *et al.* (2008); Bao *et al.* (2007); Cave *et al.* (2006); Cao *et al.* (1992); Ma *et al.* (2006, 2008). For a related structure, see Makarov *et al.* (1999).



Experimental

Crystal data

$\text{CH}_5\text{N}_2\text{O}_3\text{P}$
 $M_r = 124.04$
 Triclinic, $P\bar{1}$
 $a = 4.8559$ (17) Å
 $b = 5.910$ (2) Å
 $c = 8.101$ (3) Å
 $\alpha = 99.570$ (6)°
 $\beta = 90.784$ (6)°

$\gamma = 101.546$ (6)°
 $V = 224.36$ (14) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 0.50$ mm⁻¹
 $T = 296$ K
 $0.20 \times 0.18 \times 0.16$ mm

Data collection

Bruker SMART APEX CCD diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2007)
 $T_{\min} = 0.907$, $T_{\max} = 0.924$

1324 measured reflections
 855 independent reflections
 840 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.014$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.034$
 $wR(F^2) = 0.161$
 $S = 1.01$
 855 reflections

64 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.62$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.77$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O3}-\text{H3B}\cdots\text{O1}^{\text{i}}$	0.96	1.75	2.611 (4)	147
$\text{N1}-\text{H1A}\cdots\text{O2}^{\text{ii}}$	0.86	2.06	2.903 (4)	167
$\text{N2}-\text{H2A}\cdots\text{O1}^{\text{ii}}$	0.86	2.09	2.924 (4)	164
$\text{N1}-\text{H1B}\cdots\text{O2}^{\text{iii}}$	0.86	2.02	2.812 (4)	153
$\text{N2}-\text{H2B}\cdots\text{O3}^{\text{iv}}$	0.86	2.21	3.008 (4)	154

Symmetry codes: (i) $x+1, y, z$; (ii) $x, y-1, z$; (iii) $-x+2, -y+1, -z+1$; (iv) $-x+1, -y+1, -z+2$.

Data collection: SMART (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: DIAMOND (Brandenburg, 1999); software used to prepare material for publication: SHELXTL.

The authors acknowledge the Science and Technology Bureau of Changzhou City (project No. CQ20090004) for supporting this work.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: JJ2049).

References

- Ayyappan, P., Evans, O. R., Foxman, B. M., Wheeler, K. A., Warren, T. H. & Lin, W. B. (2001). *Inorg. Chem.* **40**, 5954–5961.
 Bao, S. S., Ma, L. F., Wang, Y., Fang, L., Zhu, C. J., Li, Y. Z. & Zheng, L. M. (2007). *Chem. Eur. J.* **13**, 2333–2343.
 Brandenburg, K. (1999). DIAMOND. Crystal Impact GbR, Bonn, Germany.
 Bruker (2007). SMART, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
 Cao, G., Hong, H. & Mallouk, T. E. (1992). *Acc. Chem. Res.* **25**, 420–427.
 Cave, D., Coomer, F. C., Molinos, E., Klauss, H. H. & Wood, P. T. (2006). *Angew. Chem. Int. Ed.* **45**, 803–806.
 Clearfield, A. (1998). *Prog. Inorg. Chem.* **47**, 371–510.
 Haga, M. A., Kobayashi, K. & Terada, K. (2007). *Coord. Chem. Rev.*, **251**, 2688–2701.
 Ma, Y. S., Li, Y. Z., Song, Y. & Zheng, L. M. (2008). *Inorg. Chem.* **47**, 4536–4544.
 Ma, Y. S., Song, Y., Du, W. X., Li, Y. Z. & Zheng, L. M. (2006). *Dalton Trans.* pp. 3228–3235.
 Makarov, S. V., Mundoma, C., Penn, J. H., Petersen, J. L., Svarovsky, S. A. & Simoyi, R. H. (1999). *Inorg. Chim. Acta*, **286**, 149–154.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Vivani, R., Alberti, G., Costantino, F. & Nocchetti, M. (2008). *Microporous Mesoporous Mater.* **107**, 58–70.

supporting information

Acta Cryst. (2010). E66, o2326 [https://doi.org/10.1107/S1600536810032083]

[Amino(iminio)methyl]phosphonate**Ting-Hai Yang, Wei Zhuang, Wei Wei, Yong-Bing Yang and Qiang Chen****S1. Comment**

In the last decade considerable attention has been afforded to the synthesis of metal phosphonates due to their potential applications in ion-exchange and sorption, catalysis, magnetism and sensors (Ayyappan *et al.*, 2001; Clearfield, 1998; Haga *et al.*, 2007; Vivani *et al.*, 2008; Bao *et al.*, 2007; Cave *et al.*, 2006; Cao *et al.*, 1992; Ma *et al.*, 2006, 2008). In order to synthesize metal phosphonates with novel structures and properties, many kinds of phosphonic acid ligands have been used. In order to study the crystal structure of phosphonic acid, we synthesized and determined the structure of the title compound (Fig. 1). As shown in Scheme 1, the molecular exists as a zwitterion, the imino group being protonated and the phosphonic acid group being deprotonated. The molecular geometry about the central C atom is strictly planar with the sum of the three angles about C being precisely 360°. The three bonds about the central carbon atom consist of two nearly equivalent C–N1 and C–N2 distances of 1.299 (5) Å and 1.314 (5) Å, respectively, and a C–P bond distance of 1.845 (3) Å. These two C–N bonds are considerably shorter than a typical C–N single bond distance of 1.47 Å. Similar zwitterions have been formed by other aminoiminomethanesulfonic acids (Makarov *et al.*, 1999). The P–O distances in these compounds range from 1.4872 (2) Å to 1.5872 (2) Å. By comparison of individual P–O distances, the H atom can be located on O3. In our crystal structure, three intermolecular hydrogen-bond interactions exist, *viz.* between the N atom and the phosphonate O atom [N1—H1A···O2, N2—H2A···O1, N1—H1B···O2, N2—H2B···O3], and between two phosphonate O atoms [O3—H3B···O1] (Table 1). Thus the molecules are interlinked by these intermolecular hydrogen bonds, forming a three-dimensional supramolecular network structure (Fig.2).

S2. Experimental

All solvents and chemicals were of analytical grade and were used without further purification. The title compound was prepared by the following reaction: A sample of 2,4,6-tri-(phosphonate ethyl)-1,3,5-triazine (9.8 g, 20 mmol) was dissolved in 6 mol/ ml HCl (20 ml), The mixture was heated (100 °C, 10 h) and then evaporated to dryness leaving a white solid. Crystallization was carried out by dissolution of 0.62 g of the title compound (about 0.5 mmol) in 10 ml water, followed by evaporation at room temperature. After two weeks, colorless block crystals obtained.

S3. Refinement

All non-hydrogen atoms were refined anisotropically, whereas the positions of all H atoms bonded to nitrogen were fixed geometrically (N—H = 0.86 Å), and included in the refinement in the riding mode, with $U_{iso}(H) = 1.2U_{eq}(N)$. The H atom in P—O—H was located in a difference Fourier map and refined with a distance restraint of O—H = 0.96 Å, and with $U_{iso}(H) = 1.5U_{eq}(O)$.




Figure 1

Molecular structure of the title compound, $(\text{NH}_2)_2\text{CPO}_3\text{H}$, showing 50% probability displacement ellipsoids




Figure 2

The cell packing diagram for the title compound viewed down the a axis.

[Amino(iminio)methyl]phosphonate

Crystal data

$\text{CH}_5\text{N}_2\text{O}_3\text{P}$
 $M_r = 124.04$
 Triclinic, $P\bar{1}$

Hall symbol: $-P\ 1$
 $a = 4.8559(17)\ \text{\AA}$
 $b = 5.910(2)\ \text{\AA}$

$c = 8.101 (3) \text{ \AA}$
 $\alpha = 99.570 (6)^\circ$
 $\beta = 90.784 (6)^\circ$
 $\gamma = 101.546 (6)^\circ$
 $V = 224.36 (14) \text{ \AA}^3$
 $Z = 2$
 $F(000) = 128$
 $D_x = 1.836 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
 Cell parameters from 1436 reflections
 $\theta = 2.6\text{--}30.3^\circ$
 $\mu = 0.50 \text{ mm}^{-1}$
 $T = 296 \text{ K}$
 Block, colorless
 $0.20 \times 0.18 \times 0.16 \text{ mm}$

Data collection

Bruker SMART APEX CCD
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 phi and ω scans
 Absorption correction: multi-scan
 (SADABS; Bruker, 2007)
 $T_{\min} = 0.907$, $T_{\max} = 0.924$

1324 measured reflections
 855 independent reflections
 840 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.014$
 $\theta_{\max} = 26.0^\circ$, $\theta_{\min} = 2.6^\circ$
 $h = -5 \rightarrow 5$
 $k = -6 \rightarrow 7$
 $l = -9 \rightarrow 9$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.034$
 $wR(F^2) = 0.161$
 $S = 1.01$
 855 reflections
 64 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.1046P)^2 + 0.7669P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.62 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.77 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
P1	0.66113 (16)	0.60637 (13)	0.74674 (10)	0.0149 (4)
O1	0.3963 (5)	0.6733 (4)	0.8092 (3)	0.0225 (6)
O2	0.7837 (5)	0.6917 (4)	0.5956 (3)	0.0224 (6)
O3	0.8828 (5)	0.6695 (5)	0.9016 (3)	0.0224 (6)
H3B	1.0481	0.6105	0.8696	0.034*
N1	0.7663 (7)	0.1822 (5)	0.6080 (4)	0.0224 (7)
H1A	0.7485	0.0324	0.5929	0.027*
H1B	0.8918	0.2651	0.5566	0.027*
N2	0.4088 (7)	0.1649 (5)	0.7899 (4)	0.0229 (7)

H2A	0.3854	0.0148	0.7776	0.027*
H2B	0.3047	0.2374	0.8554	0.027*
C1	0.6036 (7)	0.2834 (6)	0.7084 (4)	0.0168 (7)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
P1	0.0130 (6)	0.0125 (5)	0.0216 (6)	0.0045 (3)	0.0061 (4)	0.0070 (3)
O1	0.0156 (13)	0.0197 (13)	0.0350 (14)	0.0076 (10)	0.0086 (10)	0.0073 (10)
O2	0.0245 (13)	0.0199 (13)	0.0271 (14)	0.0067 (10)	0.0113 (10)	0.0124 (10)
O3	0.0150 (13)	0.0271 (14)	0.0247 (13)	0.0047 (10)	0.0029 (10)	0.0032 (10)
N1	0.0243 (16)	0.0139 (14)	0.0318 (17)	0.0057 (12)	0.0124 (13)	0.0088 (12)
N2	0.0263 (16)	0.0145 (14)	0.0299 (17)	0.0052 (12)	0.0138 (13)	0.0074 (12)
C1	0.0170 (16)	0.0147 (15)	0.0207 (16)	0.0044 (12)	0.0026 (12)	0.0073 (12)

Geometric parameters (Å, °)

P1—O2	1.487 (2)	N1—H1A	0.8600
P1—O1	1.490 (3)	N1—H1B	0.8600
P1—O3	1.587 (3)	N2—C1	1.314 (5)
P1—C1	1.845 (3)	N2—H2A	0.8600
O3—H3B	0.9600	N2—H2B	0.8600
N1—C1	1.299 (5)		
O2—P1—O1	119.46 (15)	C1—N1—H1B	120.0
O2—P1—O3	111.94 (15)	H1A—N1—H1B	120.0
O1—P1—O3	106.97 (15)	C1—N2—H2A	120.0
O2—P1—C1	108.20 (15)	C1—N2—H2B	120.0
O1—P1—C1	107.89 (15)	H2A—N2—H2B	120.0
O3—P1—C1	100.70 (15)	N1—C1—N2	122.4 (3)
P1—O3—H3B	109.3	N1—C1—P1	118.7 (3)
C1—N1—H1A	120.0	N2—C1—P1	118.8 (3)
O2—P1—C1—N1	28.6 (3)	O2—P1—C1—N2	-154.6 (3)
O1—P1—C1—N1	159.2 (3)	O1—P1—C1—N2	-24.1 (3)
O3—P1—C1—N1	-88.9 (3)	O3—P1—C1—N2	87.8 (3)

Hydrogen-bond geometry (Å, °)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O3—H3B \cdots O1 ⁱ	0.96	1.75	2.611 (4)	147
N1—H1A \cdots O2 ⁱⁱ	0.86	2.06	2.903 (4)	167
N2—H2A \cdots O1 ⁱⁱ	0.86	2.09	2.924 (4)	164
N1—H1B \cdots O2 ⁱⁱⁱ	0.86	2.02	2.812 (4)	153
N2—H2B \cdots O3 ^{iv}	0.86	2.21	3.008 (4)	154

Symmetry codes: (i) $x+1, y, z$; (ii) $x, y-1, z$; (iii) $-x+2, -y+1, -z+1$; (iv) $-x+1, -y+1, -z+2$.