

Ethyl {4-[{(1,5-dimethyl-2,4-dioxo-2,3,4,5-tetrahydro-1H-1,5-benzodiazepin-3-yl)methyl]-1,2,3-triazol-1-yl}acetate

Rachida Dardouri,^a Youssef Kandri Rodi,^a Sonia Ladeira,^b El Mokhtar Essassi^a and Seik Weng Ng^{*a}

^aLaboratoire de Chimie Organique Hétérocyclique, Pôle de Compétences Pharmaco-chimie, Université Mohammed V-Agdal, BP 1014 Avenue Ibn Batout, Rabat, Morocco, ^bService Commun Rayons-X, Laboratoire de Chimie de Coordination, 205 Route de Narbonne, Toulouse, France, and ^cDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: seikweng@um.edu.my


Received 27 October 2010; accepted 28 October 2010

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.045; wR factor = 0.124; data-to-parameter ratio = 16.7.

In the title compound, $\text{C}_{18}\text{H}_{21}\text{N}_5\text{O}_4$, the diazepine ring adopts a boat conformation with the triazolylmethyl-bearing C atom as the prow and the C atoms at the ring junction as the stern.

Related literature

For the structure of 1,5-dimethyl-3-propargyl-1,5-benzodiazepine-2,4-dione, see: Dardouri *et al.* (2010).



Experimental

Crystal data

$\text{C}_{18}\text{H}_{21}\text{N}_5\text{O}_4$	$V = 1821.56(9)\text{ \AA}^3$
$M_r = 371.40$	$Z = 4$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 8.5452(2)\text{ \AA}$	$\mu = 0.10\text{ mm}^{-1}$
$b = 15.9993(5)\text{ \AA}$	$T = 293\text{ K}$
$c = 13.9215(4)\text{ \AA}$	$0.40 \times 0.10 \times 0.05\text{ mm}$
$\beta = 106.853(1)^\circ$	

Data collection

Bruker X8 APEXII diffractometer	2909 reflections with $I > 2\sigma(I)$
15511 measured reflections	$R_{\text{int}} = 0.037$
4129 independent reflections	Standard reflections: 0

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.045$	247 parameters
$wR(F^2) = 0.124$	H-atom parameters constrained
$S = 1.03$	$\Delta\rho_{\text{max}} = 0.30\text{ e \AA}^{-3}$
4129 reflections	$\Delta\rho_{\text{min}} = -0.31\text{ e \AA}^{-3}$

Data collection: *APEX2* (Bruker, 2008); cell refinement: *SAINT* (Bruker, 2008); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2010).

We thank Université Mohammed V-Agdal and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5398).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Bruker (2008). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Dardouri, R., Ouazzani Chahdi, F., Saffon, N., Essassi, E. M. & Ng, S. W. (2010). *Acta Cryst. E66*, o1797.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2010). E66, o3039 [https://doi.org/10.1107/S1600536810044120]

Ethyl {4-[(1,5-dimethyl-2,4-dioxo-2,3,4,5-tetrahydro-1*H*-1,5-benzodiazepin-3-yl)methyl]-1,2,3-triazol-1-yl}acetate

Rachida Dardouri, Youssef Kandri Rodi, Sonia Ladeira, El Mokhtar Essassi and Seik Weng Ng

S1. Comment


1,5-Dimethyl-3-propargyl-1,5-benzodiazepine-2,4-dione, whose synthesis was reported recently (Dardouri *et al.*, 2010), possess an acetylenic linkage that can be exploited for the synthesis of other 1,5-benzodiazepine-2,4-dione derivatives. In this study, the compound is reacted with ethyl 2-azidoacetate to yield the title compound (Scheme I, Fig. 1). The ester provides three nitrogen atoms necessary for the formation of the triazolyl ring.

S2. Experimental

To a solution of 3-propargyl-1,5-dimethyl-1,5-benzodiazepine-2,4-dione (0.23 g, 1 mmol) in a *t*-butyl alcohol/water mixture (1:2, 8 ml) was added copper sulfate pentahydrate (0.25 g, 1 mmol), sodium ascorbate (0.29 g, 2 mmol) and ethyl 2-azidoacetate (0.64 g, 5 mmol). The mixture was stirred for two hours. Water (20 ml) was added and the organic compound was extracted with ethyl acetate (2 x 20 ml). The extracts were washed with brine and then dried over sodium sulfate. The compound was recrystallized from an *n*-hexane/ethyl acetate mixture to give colorless crystals.

S3. Refinement

H-atoms were placed in calculated positions (C—H 0.93–0.98 Å) and were included in the refinement in the riding model approximation, with $U(H)$ set to 1.2–1.5 $U_{eq}(C)$.

**Figure 1**

Anisotropic displacement ellipsoid plot (Barbour, 2001) of $C_{18}H_{21}N_5O_4$ at the 50% probability level; hydrogen atoms are drawn as arbitrary radius.

Ethyl {4-[{(1,5-dimethyl-2,4-dioxo-2,3,4,5-tetrahydro-1*H*-1,5-benzodiazepin-3-yl)methyl]-1,2,3-triazol-1-yl}acetate

Crystal data

$C_{18}H_{21}N_5O_4$
 $M_r = 371.40$
Monoclinic, $P2_1/c$
Hall symbol: -P 2ybc
 $a = 8.5452 (2)$ Å
 $b = 15.9993 (5)$ Å
 $c = 13.9215 (4)$ Å
 $\beta = 106.853 (1)^\circ$
 $V = 1821.56 (9)$ Å³
 $Z = 4$

$F(000) = 784$
 $D_x = 1.354 \text{ Mg m}^{-3}$
Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 3798 reflections
 $\theta = 2.5\text{--}26.4^\circ$
 $\mu = 0.10 \text{ mm}^{-1}$
 $T = 293$ K
Prism, colorless
 $0.40 \times 0.10 \times 0.05$ mm

Data collection

Bruker X8 APEXII
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 φ and ω scans
15511 measured reflections
4129 independent reflections

2909 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.037$
 $\theta_{\text{max}} = 27.5^\circ$, $\theta_{\text{min}} = 2.8^\circ$
 $h = -10 \rightarrow 10$
 $k = -20 \rightarrow 20$
 $l = -18 \rightarrow 18$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.045$
 $wR(F^2) = 0.124$
 $S = 1.03$

4129 reflections
247 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.052P)^2 + 0.6008P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 0.30 \text{ e \AA}^{-3}$$

$$\Delta\rho_{\min} = -0.31 \text{ e \AA}^{-3}$$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.56199 (17)	0.18555 (8)	0.67466 (9)	0.0438 (3)
O2	0.81954 (18)	0.21921 (9)	0.49678 (10)	0.0505 (4)
O3	0.41269 (16)	0.12018 (9)	-0.04809 (9)	0.0440 (3)
O4	0.25337 (19)	0.10260 (14)	0.05096 (12)	0.0829 (7)
N1	0.78013 (17)	0.10102 (8)	0.73772 (10)	0.0285 (3)
N2	0.96549 (17)	0.11692 (10)	0.59493 (10)	0.0344 (4)
N3	0.52214 (17)	0.02996 (9)	0.35097 (10)	0.0314 (3)
N4	0.52862 (17)	0.02234 (9)	0.25819 (10)	0.0328 (3)
N5	0.52498 (17)	0.10028 (9)	0.22117 (10)	0.0291 (3)
C1	0.88147 (19)	0.03536 (10)	0.72159 (12)	0.0271 (4)
C2	0.8961 (2)	-0.03761 (11)	0.77820 (13)	0.0347 (4)
H2	0.8390	-0.0426	0.8255	0.042*
C3	0.9939 (2)	-0.10250 (12)	0.76516 (15)	0.0421 (5)
H3	1.0037	-0.1505	0.8042	0.050*
C4	1.0772 (2)	-0.09629 (13)	0.69423 (15)	0.0449 (5)
H4	1.1407	-0.1407	0.6840	0.054*
C5	1.0662 (2)	-0.02442 (13)	0.63862 (13)	0.0395 (5)
H5	1.1237	-0.0205	0.5914	0.047*
C6	0.97042 (19)	0.04275 (11)	0.65161 (12)	0.0300 (4)
C7	0.7764 (2)	0.11896 (12)	0.84068 (13)	0.0385 (4)
H7A	0.7659	0.1781	0.8486	0.058*
H7B	0.8758	0.0996	0.8875	0.058*
H7C	0.6849	0.0909	0.8531	0.058*
C8	0.6620 (2)	0.13566 (10)	0.66069 (12)	0.0301 (4)
C9	0.6685 (2)	0.11352 (10)	0.55590 (11)	0.0263 (3)
H9	0.6798	0.0528	0.5514	0.032*
C10	0.8228 (2)	0.15520 (11)	0.54480 (12)	0.0332 (4)
C11	1.1183 (2)	0.15061 (15)	0.58176 (15)	0.0499 (5)
H11A	1.1209	0.2101	0.5909	0.075*
H11B	1.1237	0.1378	0.5154	0.075*
H11C	1.2100	0.1258	0.6304	0.075*
C12	0.5152 (2)	0.14153 (11)	0.47517 (12)	0.0314 (4)
H12A	0.4195	0.1194	0.4904	0.038*
H12B	0.5083	0.2020	0.4755	0.038*
C13	0.51605 (19)	0.11257 (10)	0.37318 (12)	0.0274 (3)
C14	0.5173 (2)	0.15758 (11)	0.29023 (12)	0.0295 (4)
H14	0.5136	0.2154	0.2830	0.035*
C15	0.5444 (2)	0.11311 (12)	0.12218 (12)	0.0339 (4)
H15A	0.6163	0.0703	0.1096	0.041*

H15B	0.5963	0.1668	0.1207	0.041*
C16	0.3849 (2)	0.11069 (11)	0.03967 (13)	0.0366 (4)
C17	0.2699 (3)	0.11868 (14)	-0.13697 (14)	0.0487 (5)
H17A	0.1960	0.0748	-0.1294	0.058*
H17B	0.3051	0.1058	-0.1956	0.058*
C18	0.1815 (2)	0.19999 (12)	-0.15270 (14)	0.0435 (5)
H18A	0.0900	0.1969	-0.2121	0.065*
H18B	0.2542	0.2436	-0.1603	0.065*
H18C	0.1431	0.2120	-0.0958	0.065*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0564 (9)	0.0414 (7)	0.0342 (7)	0.0164 (6)	0.0141 (6)	-0.0056 (6)
O2	0.0584 (9)	0.0477 (8)	0.0410 (8)	-0.0153 (7)	0.0074 (7)	0.0153 (6)
O3	0.0389 (7)	0.0690 (9)	0.0205 (6)	0.0060 (6)	0.0030 (5)	-0.0024 (6)
O4	0.0371 (9)	0.1597 (19)	0.0459 (10)	-0.0232 (10)	0.0022 (7)	0.0413 (11)
N1	0.0337 (8)	0.0327 (7)	0.0201 (7)	-0.0024 (6)	0.0091 (6)	-0.0031 (5)
N2	0.0301 (8)	0.0491 (9)	0.0242 (7)	-0.0106 (7)	0.0080 (6)	0.0014 (6)
N3	0.0320 (8)	0.0344 (8)	0.0264 (7)	-0.0013 (6)	0.0063 (6)	0.0015 (6)
N4	0.0352 (8)	0.0332 (8)	0.0274 (7)	-0.0031 (6)	0.0050 (6)	0.0005 (6)
N5	0.0287 (7)	0.0341 (8)	0.0223 (7)	-0.0019 (6)	0.0040 (5)	0.0016 (6)
C1	0.0246 (8)	0.0334 (8)	0.0217 (8)	-0.0037 (7)	0.0043 (6)	-0.0038 (6)
C2	0.0307 (9)	0.0398 (10)	0.0312 (9)	-0.0051 (7)	0.0053 (7)	0.0024 (7)
C3	0.0365 (10)	0.0369 (10)	0.0446 (11)	0.0011 (8)	-0.0013 (8)	0.0032 (8)
C4	0.0312 (10)	0.0519 (12)	0.0437 (11)	0.0111 (8)	-0.0017 (8)	-0.0094 (9)
C5	0.0256 (9)	0.0631 (13)	0.0273 (9)	0.0038 (8)	0.0035 (7)	-0.0092 (9)
C6	0.0239 (8)	0.0429 (10)	0.0203 (8)	-0.0053 (7)	0.0018 (6)	-0.0034 (7)
C7	0.0472 (11)	0.0469 (11)	0.0227 (9)	-0.0019 (9)	0.0119 (8)	-0.0072 (8)
C8	0.0376 (10)	0.0269 (8)	0.0262 (8)	-0.0007 (7)	0.0098 (7)	-0.0017 (6)
C9	0.0310 (9)	0.0263 (8)	0.0214 (8)	0.0005 (7)	0.0069 (6)	-0.0008 (6)
C10	0.0395 (10)	0.0387 (9)	0.0204 (8)	-0.0080 (8)	0.0069 (7)	-0.0015 (7)
C11	0.0375 (11)	0.0799 (15)	0.0324 (10)	-0.0239 (10)	0.0104 (8)	0.0009 (10)
C12	0.0314 (9)	0.0361 (9)	0.0259 (8)	0.0072 (7)	0.0072 (7)	0.0018 (7)
C13	0.0229 (8)	0.0319 (8)	0.0255 (8)	0.0013 (6)	0.0042 (6)	0.0024 (7)
C14	0.0299 (9)	0.0296 (8)	0.0272 (8)	0.0010 (7)	0.0051 (7)	0.0016 (7)
C15	0.0329 (9)	0.0448 (10)	0.0225 (8)	-0.0031 (8)	0.0055 (7)	0.0001 (7)
C16	0.0348 (10)	0.0418 (10)	0.0293 (9)	-0.0072 (8)	0.0033 (7)	0.0055 (7)
C17	0.0509 (12)	0.0617 (13)	0.0231 (9)	0.0071 (10)	-0.0056 (8)	-0.0067 (9)
C18	0.0438 (11)	0.0441 (11)	0.0352 (10)	-0.0068 (9)	-0.0002 (8)	0.0007 (8)

Geometric parameters (\AA , $^\circ$)

O1—C8	1.225 (2)	C5—H5	0.9300
O2—C10	1.219 (2)	C7—H7A	0.9600
O3—C16	1.319 (2)	C7—H7B	0.9600
O3—C17	1.464 (2)	C7—H7C	0.9600
O4—C16	1.186 (2)	C8—C9	1.518 (2)

N1—C8	1.360 (2)	C9—C12	1.525 (2)
N1—C1	1.420 (2)	C9—C10	1.524 (2)
N1—C7	1.471 (2)	C9—H9	0.9800
N2—C10	1.362 (2)	C11—H11A	0.9600
N2—C6	1.419 (2)	C11—H11B	0.9600
N2—C11	1.473 (2)	C11—H11C	0.9600
N3—N4	1.3145 (19)	C12—C13	1.495 (2)
N3—C13	1.362 (2)	C12—H12A	0.9700
N4—N5	1.3462 (19)	C12—H12B	0.9700
N5—C14	1.344 (2)	C13—C14	1.364 (2)
N5—C15	1.450 (2)	C14—H14	0.9300
C1—C2	1.394 (2)	C15—C16	1.507 (2)
C1—C6	1.404 (2)	C15—H15A	0.9700
C2—C3	1.377 (3)	C15—H15B	0.9700
C2—H2	0.9300	C17—C18	1.489 (3)
C3—C4	1.378 (3)	C17—H17A	0.9700
C3—H3	0.9300	C17—H17B	0.9700
C4—C5	1.374 (3)	C18—H18A	0.9600
C4—H4	0.9300	C18—H18B	0.9600
C5—C6	1.394 (3)	C18—H18C	0.9600
C16—O3—C17	116.81 (15)	C10—C9—H9	108.9
C8—N1—C1	121.62 (13)	O2—C10—N2	122.21 (16)
C8—N1—C7	117.89 (14)	O2—C10—C9	122.71 (16)
C1—N1—C7	119.01 (14)	N2—C10—C9	115.05 (15)
C10—N2—C6	122.59 (14)	N2—C11—H11A	109.5
C10—N2—C11	117.80 (16)	N2—C11—H11B	109.5
C6—N2—C11	119.40 (15)	H11A—C11—H11B	109.5
N4—N3—C13	109.17 (13)	N2—C11—H11C	109.5
N3—N4—N5	106.71 (13)	H11A—C11—H11C	109.5
C14—N5—N4	111.01 (13)	H11B—C11—H11C	109.5
C14—N5—C15	128.78 (14)	C13—C12—C9	111.66 (13)
N4—N5—C15	119.93 (14)	C13—C12—H12A	109.3
C2—C1—C6	119.01 (15)	C9—C12—H12A	109.3
C2—C1—N1	119.29 (15)	C13—C12—H12B	109.3
C6—C1—N1	121.68 (15)	C9—C12—H12B	109.3
C3—C2—C1	121.03 (17)	H12A—C12—H12B	107.9
C3—C2—H2	119.5	C14—C13—N3	108.06 (15)
C1—C2—H2	119.5	C14—C13—C12	130.07 (15)
C2—C3—C4	119.94 (18)	N3—C13—C12	121.82 (14)
C2—C3—H3	120.0	N5—C14—C13	105.05 (15)
C4—C3—H3	120.0	N5—C14—H14	127.5
C5—C4—C3	119.93 (18)	C13—C14—H14	127.5
C5—C4—H4	120.0	N5—C15—C16	113.16 (14)
C3—C4—H4	120.0	N5—C15—H15A	108.9
C4—C5—C6	121.26 (17)	C16—C15—H15A	108.9
C4—C5—H5	119.4	N5—C15—H15B	108.9
C6—C5—H5	119.4	C16—C15—H15B	108.9

C5—C6—C1	118.78 (16)	H15A—C15—H15B	107.8
C5—C6—N2	119.78 (15)	O4—C16—O3	124.56 (17)
C1—C6—N2	121.44 (15)	O4—C16—C15	125.75 (17)
N1—C7—H7A	109.5	O3—C16—C15	109.69 (15)
N1—C7—H7B	109.5	O3—C17—C18	111.90 (16)
H7A—C7—H7B	109.5	O3—C17—H17A	109.2
N1—C7—H7C	109.5	C18—C17—H17A	109.2
H7A—C7—H7C	109.5	O3—C17—H17B	109.2
H7B—C7—H7C	109.5	C18—C17—H17B	109.2
O1—C8—N1	122.20 (15)	H17A—C17—H17B	107.9
O1—C8—C9	121.75 (15)	C17—C18—H18A	109.5
N1—C8—C9	115.91 (14)	C17—C18—H18B	109.5
C8—C9—C12	112.12 (13)	H18A—C18—H18B	109.5
C8—C9—C10	105.62 (13)	C17—C18—H18C	109.5
C12—C9—C10	112.32 (14)	H18A—C18—H18C	109.5
C8—C9—H9	108.9	H18B—C18—H18C	109.5
C12—C9—H9	108.9		
C13—N3—N4—N5	-0.60 (17)	O1—C8—C9—C10	107.96 (18)
N3—N4—N5—C14	0.41 (17)	N1—C8—C9—C10	-67.92 (18)
N3—N4—N5—C15	174.79 (13)	C6—N2—C10—O2	179.53 (16)
C8—N1—C1—C2	-127.81 (17)	C11—N2—C10—O2	-5.9 (3)
C7—N1—C1—C2	38.0 (2)	C6—N2—C10—C9	1.3 (2)
C8—N1—C1—C6	53.7 (2)	C11—N2—C10—C9	175.88 (15)
C7—N1—C1—C6	-140.50 (16)	C8—C9—C10—O2	-103.15 (18)
C6—C1—C2—C3	-1.1 (2)	C12—C9—C10—O2	19.4 (2)
N1—C1—C2—C3	-179.65 (15)	C8—C9—C10—N2	75.09 (17)
C1—C2—C3—C4	-0.9 (3)	C12—C9—C10—N2	-162.40 (14)
C2—C3—C4—C5	1.8 (3)	C8—C9—C12—C13	-174.76 (14)
C3—C4—C5—C6	-0.7 (3)	C10—C9—C12—C13	66.48 (18)
C4—C5—C6—C1	-1.3 (2)	N4—N3—C13—C14	0.58 (18)
C4—C5—C6—N2	178.22 (15)	N4—N3—C13—C12	-177.28 (14)
C2—C1—C6—C5	2.2 (2)	C9—C12—C13—C14	-117.25 (19)
N1—C1—C6—C5	-179.29 (15)	C9—C12—C13—N3	60.1 (2)
C2—C1—C6—N2	-177.35 (14)	N4—N5—C14—C13	-0.06 (18)
N1—C1—C6—N2	1.2 (2)	C15—N5—C14—C13	-173.80 (15)
C10—N2—C6—C5	131.51 (17)	N3—C13—C14—N5	-0.31 (18)
C11—N2—C6—C5	-43.0 (2)	C12—C13—C14—N5	177.32 (16)
C10—N2—C6—C1	-49.0 (2)	C14—N5—C15—C16	-97.7 (2)
C11—N2—C6—C1	136.53 (17)	N4—N5—C15—C16	89.08 (19)
C1—N1—C8—O1	171.63 (16)	C17—O3—C16—O4	-1.3 (3)
C7—N1—C8—O1	5.6 (2)	C17—O3—C16—C15	179.53 (16)
C1—N1—C8—C9	-12.5 (2)	N5—C15—C16—O4	2.8 (3)
C7—N1—C8—C9	-178.49 (14)	N5—C15—C16—O3	-178.05 (15)
O1—C8—C9—C12	-14.7 (2)	C16—O3—C17—C18	79.5 (2)
N1—C8—C9—C12	169.44 (14)		