

catena-Poly[[1,10-phenanthroline- $\kappa^2 N,N'$ cadmium(II)]- μ -oxalato- $\kappa^4 O^1,O^2;O^{1'},O^{2'}$]

Yao-Kang Lv, Li-Hua Gan,* Yong-Jie Cao, Biao-Feng Gao and Liu-Hua Chen

Chemistry Department, Tongji University, Shanghai 200092, People's Republic of China

Correspondence e-mail: ganlh@tongji.edu.cn


Received 25 September 2010; accepted 8 October 2010

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(C-C) = 0.008$ Å; R factor = 0.027; wR factor = 0.066; data-to-parameter ratio = 15.1.

In the title complex, $[Cd(C_2O_4)(C_{12}H_8N_2)]_n$, the Cd^{II} atom has a distorted octahedral coordination, defined by four O atoms from two symmetry-related oxalate ligands and by two N atoms from a bidentate 1,10-phenanthroline ligand. Each oxalate ligand bridges two Cd^{II} atoms, generating a zigzag chain structure propagating along [100]. The packing of the structure is consolidated by non-classical C—H···O hydrogen-bonding interactions.

Related literature

For general background to the rational design and synthesis of metal-organic polymers, see: Kondrashev *et al.* (1985); Orioli *et al.* (2002); Athar *et al.* (2008); Lv *et al.* (2010). Wu *et al.* (2003). For related structures, see: Cao *et al.* (2009); Jeanneau *et al.* (2001).



Experimental

Crystal data

$[Cd(C_2O_4)(C_{12}H_8N_2)]$

$M_r = 380.62$

Orthorhombic, $Pna2_1$

$a = 9.7199$ (2) Å

$b = 10.3338$ (2) Å

$c = 13.1638$ (2) Å

$V = 1322.22$ (4) Å³

$Z = 4$

Mo $K\alpha$ radiation

$\mu = 1.67$ mm⁻¹

$T = 296$ K

0.29 × 0.14 × 0.10 mm

Data collection

Bruker APEXII area-detector

diffractometer

Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.76$, $T_{\max} = 0.85$

11056 measured reflections

2892 independent reflections

2386 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.034$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.027$

$wR(F^2) = 0.066$

$S = 1.00$

2892 reflections

191 parameters

1 restraint

H-atom parameters constrained

$\Delta\rho_{\max} = 0.37$ e Å⁻³

$\Delta\rho_{\min} = -0.27$ e Å⁻³

Absolute structure: Flack (1983), 1310 Friedel pairs

Flack parameter: 0.33 (4)

Table 1
Selected bond lengths (Å).

Cd1—O1	2.258 (3)	Cd1—O3 ⁱ	2.294 (3)
Cd1—O4 ⁱ	2.269 (3)	Cd1—N2	2.307 (3)
Cd1—O2	2.271 (3)	Cd1—N1	2.338 (3)

Symmetry code: (i) $x - \frac{1}{2}, -y - \frac{1}{2}, z$.

Table 2

Hydrogen-bond geometry (Å, °).

$D—H\cdots A$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
C4—H4A···O2 ⁱⁱ	0.93	2.38	3.106 (6)	134
C7—H7A···O1 ⁱⁱⁱ	0.93	2.57	3.289 (6)	134
C11—H11A···O3 ^{iv}	0.93	2.42	3.302 (7)	159

Symmetry codes: (ii) $-x + \frac{1}{2}, y + \frac{1}{2}, z - \frac{1}{2}$; (iii) $x - \frac{1}{2}, -y + \frac{1}{2}, z$; (iv) $-x + \frac{3}{2}, y + \frac{1}{2}, z + \frac{1}{2}$.

Data collection: *APEX2* (Bruker, 2006); cell refinement: *SAINT* (Bruker, 2006); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *DIAMOND* (Brandenburg & Putz, 2004); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: WM2410).

References

- Athar, M., Li, G. H., Shi, Z., Chen, Y. & Feng, S. H. (2008). *Solid State Sci.* **10**, 1853–1859.
- Brandenburg, K. & Putz, H. (2004). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Bruker (2006). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cao, X. Y., Yao, Y. G., Qin, Y. Y., Lin, Q. P., Li, Z. J., Cheng, J. K. & Hur, N. H. (2009). *CrystEngComm*, **11**, 1815–1818.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Jeanneau, E., Audebrand, N. & Louër, D. (2001). *Acta Cryst.* **C57**, 1012–1013.
- Kondrashev, Y. D., Bogdanov, V. S., Golubev, S. N. & Pron, G. F. (1985). *Zh. Strukt. Khim.* **26**, 90–93.
- Lv, Y. K., Zhan, C. H., Jiang, Z. G. & Feng, Y. L. (2010). *Inorg. Chem. Commun.* **13**, 440–444.
- Orioli, P., Bruni, B., Vaira, M. D., Messori, L. & Piccioli, F. (2002). *Inorg. Chem.* **41**, 4312–4314.
- Sheldrick, G. M. (1996). *SADABS*. University of Göettingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Wu, C. D., Lu, C. Z., Lu, S. F., Zhuang, H. H. & Huang, J. S. (2003). *Dalton Trans.* pp. 3192–3198.

supporting information

Acta Cryst. (2010). E66, m1440 [https://doi.org/10.1107/S1600536810040341]

catena-Poly[[(1,10-phenanthroline- κ^2N,N')cadmium(II)]- μ -oxalato- $\kappa^4O^1,O^2;O'^1,O'^2]$

Yao-Kang Lv, Li-Hua Gan, Yong-Jie Cao, Biao-Feng Gao and Liu-Hua Chen

S1. Comment

Rational design and synthesis of metal-organic polymers is of current interest in the field of supramolecular chemistry and crystal engineering (Athar *et al.*, 2008; Lv *et al.*, 2010; Wu *et al.*, 2003). Among the anions involved in the formation of such solids, the oxalate anion, which possesses four donor O atoms, plays an important role. Indeed, it can act either as a monodentate or a bidentate chelating ligand and can thus bridge two or more metal atoms in a variety of arrangements, as recently shown with a number of compounds (Kondrashev *et al.*, 1985; Jeanneau *et al.*, 2001; Cao *et al.*, 2009; Orioli *et al.*, 2002). We report here on the synthesis and structure of the title compound, $[Cd(C_2O_4)(C_{12}H_8N_2)]$.


As shown in Fig. 1, the central Cd(II) atom is six-coordinated by four O atoms from two symmetry-related oxalate ligands and two N atoms from a bidendate 1,10-phenanthroline ligand (Table 1), forming a distorted octahedral geometry. Each oxalate ligand bridges two cadmium(II) atoms generating a zigzag chain structure propagating along [100]. Furthermore, there are non-classical C—H \cdots O hydrogen bonds present (Table 2). Together with van der Waals forces they interconnect the zigzag chains and construct a supramolecular network (Fig. 2).

S2. Experimental


The synthesis of the title complex (I) was carried out by hydrothermal reaction. A mixture of $Cd(NO_3)_2 \cdot 4H_2O$ (1.0 mmol), $K_2C_2O_4$ (1.0 mmol), 1,10-phenanthroline (1.0 mmol) in 20 ml water was placed in a Teflon-lined stainless steel autoclave and heated at 433 K for 72 h, and then cooled to room temperature over 3 days. The resulting colorless crystals suitable for X-ray analysis were obtained in about 36% yield.

S3. Refinement

The H atoms bonded to C atoms were positioned geometrically [$C—H$ 0.93 Å $U_{iso}(H) = 1.2U_{eq}(C)$]. The crystal measured was an inversion twin with a 2:1 ratio for the twin domains.

**Figure 1**

The asymmetric unit and some symmetry-related atoms of the title complex. Displacement ellipsoids are plotted at the 30% probability level and H atoms are not labeled. [Symmetry codes: (i) $x-0.5, -y-0.5, z$; (ii) $x+0.5, -y-0.5, z$.]

**Figure 2**

Packing diagram of the title compound viewed down [010]; all H atoms were omitted for clarity.

catena-Poly[[1,10-phenanthroline- κ^2N,N')cadmium(II)]- μ -oxalato- $\kappa^4O^1,O^2;O^{1'},O^{2'}$]

Crystal data

[Cd(C₂O₄)(C₁₂H₈N₂)]
 $M_r = 380.62$

Orthorhombic, $Pna2_1$
Hall symbol: P 2c -2n

$a = 9.7199 (2)$ Å
 $b = 10.3338 (2)$ Å
 $c = 13.1638 (2)$ Å
 $V = 1322.22 (4)$ Å³
 $Z = 4$
 $F(000) = 744$
 $D_x = 1.912$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 2979 reflections
 $\theta = 2.5\text{--}27.4^\circ$
 $\mu = 1.67$ mm⁻¹
 $T = 296$ K
Block, colourless
 $0.29 \times 0.14 \times 0.10$ mm

Data collection

Bruker APEXII area-detector
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 ω scans
Absorption correction: multi-scan
(SADABS; Sheldrick, 1996)
 $T_{\min} = 0.76$, $T_{\max} = 0.85$

11056 measured reflections
2892 independent reflections
2386 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.034$
 $\theta_{\max} = 27.4^\circ$, $\theta_{\min} = 2.5^\circ$
 $h = -12 \rightarrow 12$
 $k = -13 \rightarrow 13$
 $l = -17 \rightarrow 17$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.027$
 $wR(F^2) = 0.066$
 $S = 1.00$
2892 reflections
191 parameters
1 restraint
Primary atom site location: structure-invariant
direct methods
Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0359P)^2]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.37$ e Å⁻³
 $\Delta\rho_{\min} = -0.27$ e Å⁻³
Absolute structure: Flack (1983), 1310 Friedel
pairs
Absolute structure parameter: 0.33 (4)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cd1	0.38159 (2)	-0.09300 (2)	0.33607 (7)	0.04611 (10)
O1	0.5669 (3)	-0.1452 (3)	0.2410 (2)	0.0499 (7)
O2	0.5193 (3)	-0.2166 (3)	0.4361 (2)	0.0579 (8)
O3	0.7580 (3)	-0.2606 (3)	0.2421 (2)	0.0558 (8)
O4	0.6953 (3)	-0.3510 (3)	0.4301 (2)	0.0475 (7)
N1	0.4604 (4)	0.1011 (3)	0.4063 (3)	0.0488 (9)
N2	0.2834 (4)	0.0751 (3)	0.2463 (3)	0.0480 (8)
C1	0.6195 (3)	-0.2664 (4)	0.3923 (4)	0.0395 (11)

C2	0.6499 (5)	-0.2203 (4)	0.2815 (4)	0.0413 (11)
C3	0.1925 (5)	0.0621 (5)	0.1726 (3)	0.0591 (12)
H3A	0.1661	-0.0210	0.1539	0.071*
C4	0.1343 (5)	0.1662 (7)	0.1214 (4)	0.0689 (15)
H4A	0.0695	0.1532	0.0704	0.083*
C5	0.1737 (6)	0.2856 (6)	0.1474 (4)	0.0677 (15)
H5A	0.1375	0.3563	0.1128	0.081*
C6	0.2686 (4)	0.3061 (4)	0.2260 (4)	0.0537 (11)
C7	0.3150 (6)	0.4299 (5)	0.2568 (5)	0.0691 (14)
H7A	0.2827	0.5028	0.2231	0.083*
C8	0.4037 (5)	0.4444 (4)	0.3329 (9)	0.0735 (13)
H8A	0.4315	0.5273	0.3512	0.088*
C9	0.4585 (5)	0.3335 (4)	0.3883 (4)	0.0566 (11)
C10	0.5520 (6)	0.3424 (6)	0.4680 (4)	0.0738 (15)
H10A	0.5841	0.4231	0.4886	0.089*
C11	0.5964 (6)	0.2351 (7)	0.5155 (5)	0.0736 (18)
H11A	0.6584	0.2408	0.5691	0.088*
C12	0.5478 (6)	0.1150 (5)	0.4831 (4)	0.0666 (14)
H12A	0.5780	0.0412	0.5169	0.080*
C13	0.4163 (4)	0.2091 (4)	0.3576 (3)	0.0436 (11)
C14	0.3214 (4)	0.1959 (4)	0.2760 (3)	0.0424 (9)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cd1	0.04105 (14)	0.04202 (14)	0.05528 (16)	0.00004 (11)	-0.0025 (2)	0.0066 (2)
O1	0.0474 (16)	0.0561 (17)	0.0462 (17)	0.0046 (15)	0.0020 (14)	0.0163 (16)
O2	0.0529 (18)	0.070 (2)	0.0505 (17)	0.0190 (16)	0.0150 (16)	0.0229 (17)
O3	0.0573 (19)	0.0592 (17)	0.0508 (17)	0.0164 (15)	0.0141 (15)	0.0105 (15)
O4	0.0469 (15)	0.0471 (15)	0.0484 (16)	0.0047 (13)	0.0033 (13)	0.0097 (14)
N1	0.0444 (19)	0.056 (2)	0.0457 (19)	-0.0024 (16)	-0.0074 (15)	0.0050 (18)
N2	0.0466 (19)	0.051 (2)	0.0460 (19)	0.0014 (15)	-0.0064 (16)	0.0079 (17)
C1	0.042 (3)	0.038 (2)	0.039 (2)	-0.0019 (18)	-0.0002 (18)	0.001 (2)
C2	0.044 (2)	0.032 (2)	0.049 (3)	-0.0045 (19)	0.006 (2)	0.006 (2)
C3	0.057 (3)	0.069 (3)	0.051 (3)	-0.002 (2)	-0.011 (2)	0.006 (2)
C4	0.063 (3)	0.089 (4)	0.054 (3)	0.001 (3)	-0.013 (2)	0.018 (3)
C5	0.061 (3)	0.074 (4)	0.068 (4)	0.020 (3)	0.002 (3)	0.031 (3)
C6	0.047 (2)	0.050 (2)	0.064 (3)	0.0105 (19)	0.013 (2)	0.015 (2)
C7	0.070 (3)	0.049 (3)	0.088 (4)	0.013 (2)	0.009 (3)	0.010 (3)
C8	0.082 (3)	0.0364 (19)	0.102 (4)	-0.0012 (19)	0.031 (5)	-0.002 (5)
C9	0.051 (3)	0.049 (3)	0.070 (3)	-0.013 (2)	0.013 (2)	-0.010 (2)
C10	0.074 (4)	0.074 (4)	0.074 (4)	-0.020 (3)	0.012 (3)	-0.015 (3)
C11	0.072 (4)	0.100 (5)	0.049 (3)	-0.026 (3)	-0.011 (3)	-0.008 (3)
C12	0.066 (3)	0.074 (3)	0.060 (3)	-0.014 (3)	-0.018 (3)	0.012 (3)
C13	0.0400 (18)	0.045 (2)	0.046 (3)	-0.0005 (15)	0.0091 (18)	0.004 (2)
C14	0.043 (2)	0.041 (2)	0.043 (2)	0.0026 (17)	0.0074 (18)	0.0054 (18)

Geometric parameters (\AA , $\text{^{\circ}}$)

Cd1—O1	2.258 (3)	C4—C5	1.336 (9)
Cd1—O4 ⁱ	2.269 (3)	C4—H4A	0.9300
Cd1—O2	2.271 (3)	C5—C6	1.402 (7)
Cd1—O3 ⁱ	2.294 (3)	C5—H5A	0.9300
Cd1—N2	2.307 (3)	C6—C14	1.411 (5)
Cd1—N1	2.338 (3)	C6—C7	1.416 (7)
O1—C2	1.240 (5)	C7—C8	1.330 (11)
O2—C1	1.242 (5)	C7—H7A	0.9300
O3—C2	1.244 (5)	C8—C9	1.459 (9)
O3—Cd1 ⁱⁱ	2.294 (3)	C8—H8A	0.9300
O4—C1	1.247 (5)	C9—C10	1.392 (7)
O4—Cd1 ⁱⁱ	2.269 (3)	C9—C13	1.409 (6)
N1—C12	1.329 (6)	C10—C11	1.344 (9)
N1—C13	1.356 (5)	C10—H10A	0.9300
N2—C3	1.319 (6)	C11—C12	1.394 (8)
N2—C14	1.359 (5)	C11—H11A	0.9300
C1—C2	1.563 (5)	C12—H12A	0.9300
C3—C4	1.391 (7)	C13—C14	1.422 (6)
C3—H3A	0.9300		
O1—Cd1—O4 ⁱ	151.38 (10)	C5—C4—C3	118.3 (5)
O1—Cd1—O2	73.54 (9)	C5—C4—H4A	120.9
O4 ⁱ —Cd1—O2	90.60 (10)	C3—C4—H4A	120.9
O1—Cd1—O3 ⁱ	87.77 (11)	C4—C5—C6	121.1 (5)
O4 ⁱ —Cd1—O3 ⁱ	73.03 (10)	C4—C5—H5A	119.5
O2—Cd1—O3 ⁱ	104.50 (12)	C6—C5—H5A	119.5
O1—Cd1—N2	103.06 (12)	C5—C6—C14	117.5 (4)
O4 ⁱ —Cd1—N2	98.14 (12)	C5—C6—C7	123.9 (5)
O2—Cd1—N2	164.61 (13)	C14—C6—C7	118.6 (5)
O3 ⁱ —Cd1—N2	90.22 (13)	C8—C7—C6	121.6 (5)
O1—Cd1—N1	99.37 (13)	C8—C7—H7A	119.2
O4 ⁱ —Cd1—N1	105.35 (12)	C6—C7—H7A	119.2
O2—Cd1—N1	93.45 (13)	C7—C8—C9	121.7 (5)
O3 ⁱ —Cd1—N1	161.94 (12)	C7—C8—H8A	119.2
N2—Cd1—N1	72.07 (12)	C9—C8—H8A	119.2
C2—O1—Cd1	115.5 (3)	C10—C9—C13	117.9 (5)
C1—O2—Cd1	115.2 (3)	C10—C9—C8	124.3 (5)
C2—O3—Cd1 ⁱⁱ	116.0 (3)	C13—C9—C8	117.8 (5)
C1—O4—Cd1 ⁱⁱ	115.6 (3)	C11—C10—C9	120.4 (5)
C12—N1—C13	118.3 (4)	C11—C10—H10A	119.8
C12—N1—Cd1	127.1 (3)	C9—C10—H10A	119.8
C13—N1—Cd1	114.5 (3)	C10—C11—C12	119.0 (5)
C3—N2—C14	119.1 (4)	C10—C11—H11A	120.5
C3—N2—Cd1	125.3 (3)	C12—C11—H11A	120.5
C14—N2—Cd1	115.6 (3)	N1—C12—C11	123.0 (5)
O2—C1—O4	124.6 (5)	N1—C12—H12A	118.5

O2—C1—C2	117.1 (4)	C11—C12—H12A	118.5
O4—C1—C2	118.3 (4)	N1—C13—C9	121.5 (4)
O1—C2—O3	125.5 (5)	N1—C13—C14	118.9 (4)
O1—C2—C1	117.9 (4)	C9—C13—C14	119.5 (4)
O3—C2—C1	116.6 (4)	N2—C14—C6	120.5 (4)
N2—C3—C4	123.4 (5)	N2—C14—C13	118.7 (3)
N2—C3—H3A	118.3	C6—C14—C13	120.7 (4)
C4—C3—H3A	118.3		
O4 ⁱ —Cd1—O1—C2	−54.2 (5)	O2—C1—C2—O3	172.9 (5)
O2—Cd1—O1—C2	4.5 (3)	O4—C1—C2—O3	−8.0 (5)
O3 ⁱ —Cd1—O1—C2	−101.3 (4)	C14—N2—C3—C4	1.2 (7)
N2—Cd1—O1—C2	169.0 (3)	Cd1—N2—C3—C4	179.0 (4)
N1—Cd1—O1—C2	95.4 (3)	N2—C3—C4—C5	0.9 (8)
O1—Cd1—O2—C1	−7.7 (3)	C3—C4—C5—C6	−1.6 (8)
O4 ⁱ —Cd1—O2—C1	148.1 (3)	C4—C5—C6—C14	0.4 (7)
O3 ⁱ —Cd1—O2—C1	75.5 (3)	C4—C5—C6—C7	179.7 (5)
N2—Cd1—O2—C1	−87.0 (6)	C5—C6—C7—C8	179.0 (6)
N1—Cd1—O2—C1	−106.5 (3)	C14—C6—C7—C8	−1.8 (8)
O1—Cd1—N1—C12	−78.2 (4)	C6—C7—C8—C9	0.3 (10)
O4 ⁱ —Cd1—N1—C12	87.2 (4)	C7—C8—C9—C10	179.6 (6)
O2—Cd1—N1—C12	−4.4 (4)	C7—C8—C9—C13	1.1 (9)
O3 ⁱ —Cd1—N1—C12	169.5 (4)	C13—C9—C10—C11	−2.0 (7)
N2—Cd1—N1—C12	−179.0 (4)	C8—C9—C10—C11	179.4 (6)
O1—Cd1—N1—C13	97.6 (3)	C9—C10—C11—C12	0.4 (9)
O4 ⁱ —Cd1—N1—C13	−96.9 (3)	C13—N1—C12—C11	−0.3 (8)
O2—Cd1—N1—C13	171.5 (3)	Cd1—N1—C12—C11	175.5 (4)
O3 ⁱ —Cd1—N1—C13	−14.6 (6)	C10—C11—C12—N1	0.8 (9)
N2—Cd1—N1—C13	−3.2 (3)	C12—N1—C13—C9	−1.5 (6)
O1—Cd1—N2—C3	88.0 (4)	Cd1—N1—C13—C9	−177.7 (3)
O4 ⁱ —Cd1—N2—C3	−72.7 (4)	C12—N1—C13—C14	−179.2 (4)
O2—Cd1—N2—C3	163.3 (4)	Cd1—N1—C13—C14	4.5 (5)
O3 ⁱ —Cd1—N2—C3	0.2 (4)	C10—C9—C13—N1	2.6 (6)
N1—Cd1—N2—C3	−176.3 (4)	C8—C9—C13—N1	−178.7 (5)
O1—Cd1—N2—C14	−94.2 (3)	C10—C9—C13—C14	−179.6 (4)
O4 ⁱ —Cd1—N2—C14	105.1 (3)	C8—C9—C13—C14	−1.0 (6)
O2—Cd1—N2—C14	−18.9 (6)	C3—N2—C14—C6	−2.5 (6)
O3 ⁱ —Cd1—N2—C14	178.0 (3)	Cd1—N2—C14—C6	179.5 (3)
N1—Cd1—N2—C14	1.6 (3)	C3—N2—C14—C13	178.1 (4)
Cd1—O2—C1—O4	−169.5 (3)	Cd1—N2—C14—C13	0.1 (5)
Cd1—O2—C1—C2	9.5 (4)	C5—C6—C14—N2	1.7 (6)
Cd1 ⁱⁱ —O4—C1—O2	−174.2 (3)	C7—C6—C14—N2	−177.6 (4)
Cd1 ⁱⁱ —O4—C1—C2	6.8 (4)	C5—C6—C14—C13	−178.9 (4)
Cd1—O1—C2—O3	−179.9 (4)	C7—C6—C14—C13	1.8 (6)
Cd1—O1—C2—C1	−1.5 (5)	N1—C13—C14—N2	−3.2 (6)
Cd1 ⁱⁱ —O3—C2—O1	−176.9 (4)	C9—C13—C14—N2	179.0 (4)
Cd1 ⁱⁱ —O3—C2—C1	4.6 (4)	N1—C13—C14—C6	177.4 (4)

O2—C1—C2—O1	−5.6 (5)	C9—C13—C14—C6	−0.4 (6)
O4—C1—C2—O1	173.4 (5)		

Symmetry codes: (i) $x-1/2, -y-1/2, z$; (ii) $x+1/2, -y-1/2, z$.

Hydrogen-bond geometry (\AA , °)

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C4—H4A···O2 ⁱⁱⁱ	0.93	2.38	3.106 (6)	134
C7—H7A···O1 ^{iv}	0.93	2.57	3.289 (6)	134
C11—H11A···O3 ^v	0.93	2.42	3.302 (7)	159

Symmetry codes: (iii) $-x+1/2, y+1/2, z-1/2$; (iv) $x-1/2, -y+1/2, z$; (v) $-x+3/2, y+1/2, z+1/2$.