

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Bis(2,3,5,6-tetra-2-pyridylpyrazine- κ^3N^2,N^1,N^6)nickel(II) dithiocyanate dihydrate

 Noelia De la Pinta,^a M. Luz Fidalgo,^b José M. Ezpeleta,^c Roberto Cortés^a and Gotzon Madariaga^{d*}

^aDepartamento de Química Inorgánica, Facultad de Ciencia y Tecnología, Universidad del País Vasco, Apdo. 644, E-48080 Bilbao, Spain, ^bDepartamento de Química Inorgánica, Facultad de Farmacia, Universidad del País Vasco, Apdo. 450, E-01080 Vitoria, Spain, ^cDepartamento de Física Aplicada II, Facultad de Farmacia, Universidad del País Vasco, Apdo. 450, E-01080 Vitoria, Spain, and ^dDepartamento de Física de la Materia Condensada, Facultad de Ciencia y Tecnología, Universidad del País Vasco, Apdo. 644, E-48080 Bilbao, Spain
Correspondence e-mail: gotzon.madariaga@ehu.es


Received 14 December 2010; accepted 20 December 2010

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(C-C) = 0.003$ Å; R factor = 0.048; wR factor = 0.123; data-to-parameter ratio = 23.1.

In the title compound, $[Ni(C_{24}H_{16}N_6)_2](NCS)_2 \cdot 2H_2O$, the central Ni^{II} ion is octahedrally coordinated by six N atoms of two tridentate 2,3,5,6-tetra-2-pyridylpyrazine ligands (tppz). Two thiocyanate anions act as counter-ions and two water molecules act as solvation agents. O—H...N hydrogen bonds are observed in the crystal structure.

Related literature

For related structures including $[M(II)(tppz)_2]^{2+}$ cations, see: Ruminski & Kiplinger (1990); Arana *et al.* (1992); Lainé *et al.* (1995); Allis *et al.* (2004); Burkholder & Zubieta (2004); Haines *et al.* (2000). For the application of a $[Co(II)(tppz)_2]^{2+}$ complex as a homogeneous catalyst, see: Königstein & Bauer (1994, 1997).



Experimental

Crystal data

$[Ni(C_{24}H_{16}N_6)_2](NCS)_2 \cdot 2H_2O$
 $M_r = 987.76$
 Monoclinic, $C2/c$
 $a = 17.9091$ (4) Å
 $b = 13.6851$ (2) Å
 $c = 19.4650$ (4) Å
 $\beta = 106.161$ (2)°

$V = 4582.11$ (15) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.57$ mm⁻¹
 $T = 293$ K
 $0.35 \times 0.26 \times 0.21$ mm

Data collection

Oxford Diffraction Xcalibur
 Sapphire2 diffractometer
 23806 measured reflections

7385 independent reflections
 4946 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.031$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.048$
 $wR(F^2) = 0.123$
 $S = 0.94$
 7385 reflections
 320 parameters
 4 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{max} = 0.56$ e Å⁻³
 $\Delta\rho_{min} = -0.43$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$O1W-H1W1 \cdots N9^i$	0.85 (2)	2.28 (3)	3.116 (4)	166 (3)
$O1W-H1W2 \cdots N1^{ii}$	0.85 (4)	2.20 (3)	3.044 (4)	173 (4)

Symmetry codes: (i) $-x, y, -z + \frac{1}{2}$; (ii) $-x + \frac{1}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *CrysAlis PRO* (Oxford Diffraction, 2007); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SIR97* (Altomare *et al.*, 1999); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *Mercury* (Macrae *et al.*, 2006); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

This work was supported by the Universidad del País Vasco (UPV 00169.125–13956/2004), the Basque Government (IT-282–07), and the Ministerio de Ciencia y Tecnología (CTQ2005–05778-PPQ). NDelaP thanks the UPV/EHU for financial support under the "Convocatoria para la concesión de ayudas de especialización para investigadores doctores en la UPV/EHU (2008)".

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FJ2379).

References

- Allis, D. G., Burkholder, E. & Zubieta, J. (2004). *Polyhedron*, **23**, 1145–1152.
 Altomare, A., Burla, M. C., Camalli, M., Cascarano, G. L., Giacovazzo, C., Guagliardi, A., Moliterni, A. G. G., Polidori, G. & Spagna, R. (1999). *J. Appl. Cryst.* **32**, 115–119.
 Arana, C., Yan, S., Keshavarz, K. M., Potes, K. T. & Abruña, H. D. (1992). *Inorg. Chem.* **31**, 3680–3682.
 Burkholder, E. & Zubieta, J. (2004). *Inorg. Chim. Acta*, **357**, 1229–1235.
 Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
 Haines, R. I., Hutchings, D. R. & Strickland, D. W. (2000). *Inorganic Reaction Mechanisms (Amsterdam)*, **2**, 223–233.
 Königstein, C. & Bauer, R. (1994). *Hydrogen Energy Prog. X, Proc. World Hydrogen Energy Conf.* 10th, **2**, 717–725.
 Königstein, C. & Bauer, R. (1997). *Int. Journal Hydrogen Energy*, **22**, 471–474.

- Lainé, P., Gourdon, A. & Launay, J. P. (1995). *Inorg. Chem.* **34**, 5156–5165.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- Oxford Diffraction (2007). *CrysAlis PRO*. Oxford Diffraction Ltd, Abingdon, England.
- Ruminski, R. R. & Kiplinger, J. L. (1990). *Inorg. Chem.* **29**, 4581–4584.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supporting information

Acta Cryst. (2011). E67, m129–m130 [https://doi.org/10.1107/S1600536810053419]

Bis(2,3,5,6-tetra-2-pyridylpyrazine- κ^3N^2,N^1,N^6)nickel(II) dithiocyanate dihydrate

Noelia De la Pinta, M. Luz Fidalgo, José M. Ezpeleta, Roberto Cortés and Gotzon Madariaga

S1. Comment

The combination of divalent cations of the second half of the transition series with the ligand tppz, gives coordination cations of the type $[M^{II}(\text{tppz})_2]^{2+}$, where the terminal nitrogen atoms of one extreme of the tppz ligand are directed towards the metallic atoms and the corresponding N atoms of the other extreme remain uncoordinated (Lainé *et al.*, 1995). These tppz cations are part of different coordination compounds (Allis *et al.*, 2004, Burkholder & Zubieta, 2004, Ruminski *et al.*, 1990, Haines *et al.*, 2000, Arana *et al.*, 1992, Köningstein *et al.*, 1997, 1994). We here report the crystal structure of a new compound, $[\text{Ni}(\text{C}_{24}\text{H}_{16}\text{N}_6)_2](\text{NCS})_2 \cdot 2\text{H}_2\text{O}$, which is made up of Ni^{II} cations coordinated to six nitrogen atoms of two tridentate tppz ligands. These monomeric entities reach the neutrality with two thiocyanate anions and two molecules of water act as solvent agents. The nitrogen atoms of each tppz, coordinated to the cation, are in the same plane. The Ni^{II} cation has a distorted octahedral environment, in which the bonds to the two pyrazine nitrogen atoms (Ni1—N3 and Ni1—N7) are significantly shorter than the bonds to the pyridyl nitrogen atoms (Ni1—N2 and Ni1—N6). The N—Ni1—N angles involving the atoms of the equatorial plane with respect to the short axis differ remarkably from the ideal octahedral values, the Ni1 atom not deviating significantly [0.0006 (2) Å] from the average plane.

The individual pyridyl rings are planar (maximum average displacement with respect to the plane of the ring, 0.011 Å), while the two pyrazine rings are significantly puckered. Nitrogen atoms of the non-coordinated pyridyl rings of the two tppz ligands point to the ligand metalated side instead of the free nitrogen atom of the pyrazine ring. These structural features of the coordinated tppz ligand would account for its tendency to adopt bis-chelation in this type of complexes. Weak π - π interactions appear to occur due to overlap between the pyridyl rings labelled by N5 and N9ⁱⁱⁱ (iii = $-x, -1 + y, 1/2 - z$) with a distance between the ring centroids of 3.7713 (12) Å.

The crystal packing of the bulky building block units leaves cavities of 116 Å³ where water molecules and NCS anions are located (Fig. 2). The low density of the material reflects this open structure. Most relevant H bonds are listed in Table 1.

S2. Experimental

Crystals of $[\text{Ni}(\text{C}_{24}\text{H}_{16}\text{N}_6)_2](\text{NCS})_2 \cdot 2\text{H}_2\text{O}$ were prepared by mixing an acetonitrile solution (10 ml) of $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$ (112 mg, 0.50 mmol) and another acetonitrile solution (10 ml) of 2,3,5,6-tetra-2-pyridylpyrazine (97.1 mg, 0.25 mmol). After a vigorous stirring of about 30 minutes at a temperature of 303 K, an aqua/acetonitrile (50%) solution (10 ml) of sodium thiocyanate (101.3 mg, 1.25 mmol) was added. The resultant solution was stirred at 313 K for 25 minutes and at room temperature for the following two days. Then the precipitate that did form was filtered off. Finally, the resulting solution was maintained at room temperature until prismatic green crystals were formed by slow evaporation. These crystals were found to be stable to X-ray exposure.

S3. Refinement

Structure solution by direct methods in the space group $C2/c$, followed by refinement, based on F^2 , of atomic coordinates and anisotropic displacement parameters, was performed using the programs *SIR97* (Altomare *et al.*, 1999) and *SHELXL97* (Sheldrick, 2008) successively. H atoms bonded to C atoms were found in successive difference Fourier maps and refined using a riding model, with $C-H = 0.93 \text{ \AA}$ and with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$. H atoms of O1W (see Fig. 1 for labelling) were approximately located in a difference Fourier map. During the refinement their positions were tightly restrained to the ideal geometry [$O-H=0.85(1) \text{ \AA}$, $H-O-H=107(3)^\circ$] with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$. Also the distances $H1W1-N9^i$ ($i = -x, y, -z + 1/2$ and $H1W2-N1^{ii}$ ($ii = -x + 1/2, y + 1/2, -z + 1/2$) were forced to be equal (within 0.02 \AA). The highest residual electron density is 0.02 \AA from atom Ni1 and the deepest hole is 0.65 \AA from atom S1.




Figure 1

Asymmetric unit of $[\text{Ni}(\text{C}_{24}\text{H}_{16}\text{N}_6)_2](\text{NCS})_2 \cdot 2\text{H}_2\text{O}$ showing the most relevant labels. H atoms excluded for clarity. Ellipsoids drawn at 50% probability level.




Figure 2

Projection of $[\text{Ni}(\text{C}_{24}\text{H}_{16}\text{N}_6)_2](\text{NCS})_2 \cdot 2\text{H}_2\text{O}$ packing along the **b** axis showing the link between pyridyl and NCS groups through the N1—O1W—N9 H-bonds.

Bis(2,3,5,6-tetra-2-pyridylpyrazine- $\kappa^3\text{N}^2, \text{N}^1, \text{N}^6$)nickel(II) dithiocyanate dihydrate

Crystal data

$[\text{Ni}(\text{C}_{24}\text{H}_{16}\text{N}_6)_2](\text{NCS})_2 \cdot 2\text{H}_2\text{O}$

$M_r = 987.76$

Monoclinic, $C2/c$

Hall symbol: $-C 2yc$

$a = 17.9091(4) \text{ \AA}$

$b = 13.6851(2) \text{ \AA}$

$c = 19.4650(4) \text{ \AA}$

$\beta = 106.161(2)^\circ$

$V = 4582.11(15) \text{ \AA}^3$

$Z = 4$

$F(000) = 2040$

$D_x = 1.432 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 9376 reflections

$\theta = 3.0\text{--}32.2^\circ$

$\mu = 0.57 \text{ mm}^{-1}$

$T = 293 \text{ K}$

Prism, green

$0.35 \times 0.26 \times 0.21 \text{ mm}$

Data collection

Oxford Diffraction Xcalibur Sapphire2 window
diffractometer

Radiation source: sealed X-ray tube

Graphite monochromator

Detector resolution: $8.3504 \text{ pixels mm}^{-1}$

ω scans

23806 measured reflections

7385 independent reflections

4946 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.031$ $\theta_{\text{max}} = 32.2^\circ$, $\theta_{\text{min}} = 3.0^\circ$ $h = -25 \rightarrow 21$ $k = -18 \rightarrow 20$ $l = -28 \rightarrow 29$ *Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.048$ $wR(F^2) = 0.123$ $S = 0.94$

7385 reflections

320 parameters

4 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sitesH atoms treated by a mixture of independent
and constrained refinement $w = 1/[\sigma^2(F_o^2) + (0.0771P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\text{max}} < 0.001$ $\Delta\rho_{\text{max}} = 0.56 \text{ e } \text{\AA}^{-3}$ $\Delta\rho_{\text{min}} = -0.43 \text{ e } \text{\AA}^{-3}$ *Special details*

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Ni1	0	0.310758 (18)	0.25	0.03035 (9)
N1	0.30210 (13)	0.08495 (16)	0.16393 (13)	0.0748 (6)
C1	0.24787 (15)	0.05532 (16)	0.17858 (12)	0.0613 (6)
S1	0.17309 (4)	0.01597 (5)	0.20104 (4)	0.07129 (18)
N2	0.09989 (8)	0.27794 (9)	0.33345 (7)	0.0357 (3)
N3	0	0.16447 (12)	0.25	0.0275 (4)
N4	0	-0.03295 (12)	0.25	0.0297 (4)
N5	0.09070 (11)	-0.00169 (11)	0.43264 (8)	0.0519 (4)
N6	0.07402 (8)	0.34349 (10)	0.18698 (7)	0.0346 (3)
N7	0	0.45723 (12)	0.25	0.0282 (4)
N8	0	0.65503 (12)	0.25	0.0287 (4)
N9	0.01006 (10)	0.61824 (11)	0.07513 (8)	0.0450 (4)
C2	0.11628 (9)	0.18165 (10)	0.34250 (8)	0.0318 (3)
C3	0.18842 (10)	0.14902 (13)	0.38158 (11)	0.0451 (4)
H3	0.1988	0.0824	0.3867	0.068*
C4	0.24504 (12)	0.21619 (16)	0.41296 (13)	0.0603 (6)
H4	0.2941	0.1955	0.4393	0.09*
C5	0.22787 (13)	0.31443 (15)	0.40478 (13)	0.0631 (6)
H5	0.2649	0.3611	0.426	0.095*
C6	0.15514 (12)	0.34205 (14)	0.36470 (11)	0.0500 (5)
H6	0.1438	0.4084	0.359	0.075*

C7	0.05337 (9)	0.11697 (10)	0.30076 (8)	0.0279 (3)
C8	0.04567 (8)	0.01515 (10)	0.30544 (8)	0.0284 (3)
C9	0.08377 (9)	-0.04437 (11)	0.36910 (8)	0.0327 (3)
C10	0.10704 (10)	-0.13895 (12)	0.36186 (10)	0.0399 (4)
H10	0.0997	-0.1664	0.3168	0.06*
C11	0.14136 (11)	-0.19211 (14)	0.42277 (12)	0.0538 (5)
H11	0.1583	-0.2557	0.4197	0.081*
C12	0.14968 (14)	-0.14892 (19)	0.48751 (13)	0.0686 (7)
H12	0.1729	-0.1827	0.5295	0.103*
C13	0.12351 (16)	-0.05524 (19)	0.49030 (11)	0.0702 (7)
H13	0.129	-0.0275	0.535	0.105*
C14	0.08424 (9)	0.43995 (10)	0.17709 (8)	0.0299 (3)
C15	0.14292 (10)	0.47369 (12)	0.15021 (10)	0.0405 (4)
H15	0.1486	0.5402	0.1433	0.061*
C16	0.19345 (11)	0.40681 (14)	0.13364 (11)	0.0494 (5)
H16	0.2333	0.4279	0.1151	0.074*
C17	0.18406 (11)	0.30917 (13)	0.14488 (10)	0.0452 (4)
H17	0.2184	0.2634	0.1355	0.068*
C18	0.12336 (11)	0.28013 (12)	0.17007 (9)	0.0413 (4)
H18	0.116	0.2137	0.1757	0.062*
C19	0.03442 (9)	0.50448 (10)	0.20708 (8)	0.0282 (3)
C20	0.02362 (9)	0.60616 (10)	0.20014 (8)	0.0284 (3)
C21	0.03384 (9)	0.66321 (11)	0.13860 (8)	0.0314 (3)
C22	0.06215 (11)	0.75722 (12)	0.14626 (10)	0.0443 (4)
H22	0.078	0.7865	0.1911	0.066*
C23	0.06656 (12)	0.80718 (14)	0.08575 (12)	0.0551 (5)
H23	0.0854	0.8708	0.0894	0.083*
C24	0.04322 (13)	0.76275 (17)	0.02107 (13)	0.0601 (6)
H24	0.0457	0.7952	-0.0202	0.09*
C25	0.01589 (13)	0.66902 (18)	0.01790 (11)	0.0581 (5)
H25	0.0004	0.6387	-0.0266	0.087*
O1W	0.15572 (18)	0.5266 (3)	0.47152 (14)	0.1363 (10)
H1W1	0.1090 (10)	0.546 (3)	0.465 (2)	0.204*
H1W2	0.171 (2)	0.545 (4)	0.4359 (16)	0.204*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ni1	0.04230 (17)	0.01832 (13)	0.03170 (15)	0	0.01241 (12)	0
N1	0.0666 (13)	0.0573 (12)	0.1034 (17)	0.0104 (10)	0.0287 (12)	0.0182 (11)
C1	0.0809 (16)	0.0431 (11)	0.0536 (12)	0.0199 (11)	0.0083 (11)	0.0009 (9)
S1	0.0804 (4)	0.0647 (4)	0.0714 (4)	0.0071 (3)	0.0255 (3)	0.0001 (3)
N2	0.0454 (8)	0.0242 (6)	0.0360 (7)	-0.0069 (5)	0.0086 (6)	-0.0026 (5)
N3	0.0302 (9)	0.0223 (8)	0.0298 (9)	0	0.0081 (7)	0
N4	0.0345 (9)	0.0217 (8)	0.0317 (9)	0	0.0073 (7)	0
N5	0.0743 (11)	0.0435 (9)	0.0326 (8)	-0.0023 (8)	0.0063 (8)	0.0012 (6)
N6	0.0471 (8)	0.0250 (6)	0.0346 (7)	0.0031 (5)	0.0160 (6)	-0.0022 (5)
N7	0.0346 (9)	0.0222 (8)	0.0300 (9)	0	0.0127 (7)	0

N8	0.0363 (9)	0.0219 (8)	0.0295 (9)	0	0.0117 (7)	0
N9	0.0563 (9)	0.0464 (9)	0.0335 (7)	-0.0011 (7)	0.0146 (7)	0.0023 (6)
C2	0.0375 (8)	0.0254 (7)	0.0310 (7)	-0.0036 (6)	0.0072 (6)	-0.0017 (6)
C3	0.0373 (9)	0.0359 (9)	0.0567 (11)	-0.0003 (7)	0.0039 (8)	-0.0016 (8)
C4	0.0395 (10)	0.0553 (12)	0.0736 (15)	-0.0079 (9)	-0.0048 (10)	-0.0044 (11)
C5	0.0536 (12)	0.0478 (12)	0.0753 (15)	-0.0209 (9)	-0.0029 (11)	-0.0068 (10)
C6	0.0578 (12)	0.0305 (8)	0.0548 (11)	-0.0128 (8)	0.0042 (9)	-0.0058 (8)
C7	0.0323 (7)	0.0235 (7)	0.0276 (7)	-0.0003 (5)	0.0078 (6)	-0.0007 (5)
C8	0.0315 (7)	0.0231 (7)	0.0296 (7)	0.0012 (5)	0.0070 (6)	0.0010 (5)
C9	0.0342 (8)	0.0276 (7)	0.0329 (8)	-0.0031 (6)	0.0038 (6)	0.0044 (6)
C10	0.0397 (9)	0.0313 (8)	0.0464 (10)	0.0011 (7)	0.0081 (7)	0.0075 (7)
C11	0.0410 (10)	0.0405 (10)	0.0753 (14)	0.0034 (8)	0.0087 (10)	0.0251 (10)
C12	0.0684 (14)	0.0713 (15)	0.0527 (13)	-0.0042 (12)	-0.0056 (11)	0.0324 (12)
C13	0.0956 (18)	0.0725 (16)	0.0327 (10)	-0.0073 (13)	0.0019 (11)	0.0079 (10)
C14	0.0380 (8)	0.0244 (7)	0.0287 (7)	0.0027 (6)	0.0117 (6)	-0.0017 (5)
C15	0.0464 (10)	0.0293 (8)	0.0521 (10)	-0.0007 (7)	0.0242 (8)	-0.0010 (7)
C16	0.0466 (10)	0.0456 (10)	0.0650 (12)	-0.0007 (8)	0.0306 (9)	-0.0082 (9)
C17	0.0485 (10)	0.0391 (9)	0.0523 (10)	0.0080 (7)	0.0208 (9)	-0.0096 (8)
C18	0.0550 (10)	0.0281 (8)	0.0444 (9)	0.0078 (7)	0.0197 (8)	-0.0051 (7)
C19	0.0347 (7)	0.0238 (7)	0.0281 (7)	0.0002 (5)	0.0121 (6)	0.0002 (5)
C20	0.0346 (7)	0.0222 (6)	0.0300 (7)	0.0002 (5)	0.0119 (6)	0.0021 (5)
C21	0.0361 (8)	0.0274 (7)	0.0344 (8)	0.0043 (6)	0.0158 (6)	0.0061 (6)
C22	0.0557 (11)	0.0321 (8)	0.0523 (11)	-0.0023 (7)	0.0267 (9)	0.0037 (7)
C23	0.0629 (13)	0.0405 (10)	0.0730 (14)	0.0026 (9)	0.0372 (11)	0.0207 (10)
C24	0.0595 (13)	0.0712 (14)	0.0579 (13)	0.0129 (10)	0.0300 (10)	0.0339 (11)
C25	0.0623 (13)	0.0775 (15)	0.0352 (10)	0.0049 (11)	0.0147 (9)	0.0107 (9)
O1W	0.144 (2)	0.167 (3)	0.108 (2)	0.035 (2)	0.0506 (18)	-0.0268 (18)

Geometric parameters (Å, °)

Ni1—N3	2.0019 (17)	C6—H6	0.93
Ni1—N7	2.0045 (17)	C7—C8	1.4057 (19)
Ni1—N6 ⁱ	2.0899 (14)	C8—C9	1.481 (2)
Ni1—N6	2.0899 (14)	C9—C10	1.379 (2)
Ni1—N2 ⁱ	2.1030 (14)	C10—C11	1.381 (2)
Ni1—N2	2.1030 (14)	C10—H10	0.93
N1—C1	1.159 (3)	C11—C12	1.362 (3)
C1—S1	1.613 (3)	C11—H11	0.93
N2—C6	1.336 (2)	C12—C13	1.371 (4)
N2—C2	1.3509 (19)	C12—H12	0.93
N3—C7 ⁱ	1.3366 (16)	C13—H13	0.93
N3—C7	1.3366 (16)	C14—C15	1.378 (2)
N4—C8	1.3325 (16)	C14—C19	1.485 (2)
N4—C8 ⁱ	1.3325 (16)	C15—C16	1.387 (2)
N5—C13	1.332 (3)	C15—H15	0.93
N5—C9	1.342 (2)	C16—C17	1.372 (3)
N6—C18	1.343 (2)	C16—H16	0.93
N6—C14	1.3538 (19)	C17—C18	1.370 (3)

N7—C19	1.3353 (16)	C17—H17	0.93
N7—C19 ⁱ	1.3353 (16)	C18—H18	0.93
N8—C20	1.3404 (17)	C19—C20	1.4062 (19)
N8—C20 ⁱ	1.3404 (17)	C20—C21	1.483 (2)
N9—C21	1.339 (2)	C21—C22	1.376 (2)
N9—C25	1.342 (2)	C22—C23	1.383 (3)
C2—C3	1.378 (2)	C22—H22	0.93
C2—C7	1.484 (2)	C23—C24	1.355 (3)
C3—C4	1.379 (3)	C23—H23	0.93
C3—H3	0.93	C24—C25	1.368 (3)
C4—C5	1.378 (3)	C24—H24	0.93
C4—H4	0.93	C25—H25	0.93
C5—C6	1.372 (3)	O1W—H1W1	0.85 (2)
C5—H5	0.93	O1W—H1W2	0.85 (4)
N3—Ni1—N7	180	N5—C9—C10	123.34 (15)
N3—Ni1—N6 ⁱ	102.38 (4)	N5—C9—C8	115.76 (14)
N7—Ni1—N6 ⁱ	77.62 (4)	C10—C9—C8	120.84 (14)
N3—Ni1—N6	102.38 (4)	C9—C10—C11	118.80 (18)
N7—Ni1—N6	77.62 (4)	C9—C10—H10	120.6
N6 ⁱ —Ni1—N6	155.25 (7)	C11—C10—H10	120.6
N3—Ni1—N2 ⁱ	77.67 (4)	C12—C11—C10	118.28 (19)
N7—Ni1—N2 ⁱ	102.33 (4)	C12—C11—H11	120.9
N6 ⁱ —Ni1—N2 ⁱ	87.53 (5)	C10—C11—H11	120.9
N6—Ni1—N2 ⁱ	97.74 (5)	C11—C12—C13	119.46 (18)
N3—Ni1—N2	77.67 (4)	C11—C12—H12	120.3
N7—Ni1—N2	102.33 (4)	C13—C12—H12	120.3
N6 ⁱ —Ni1—N2	97.74 (5)	N5—C13—C12	123.8 (2)
N6—Ni1—N2	87.53 (5)	N5—C13—H13	118.1
N2 ⁱ —Ni1—N2	155.34 (7)	C12—C13—H13	118.1
N1—C1—S1	178.4 (2)	N6—C14—C15	121.91 (14)
C6—N2—C2	118.34 (15)	N6—C14—C19	113.83 (13)
C6—N2—Ni1	125.08 (12)	C15—C14—C19	123.72 (14)
C2—N2—Ni1	114.60 (10)	C14—C15—C16	118.90 (16)
C7 ⁱ —N3—C7	121.80 (17)	C14—C15—H15	120.5
C7 ⁱ —N3—Ni1	119.10 (9)	C16—C15—H15	120.5
C7—N3—Ni1	119.10 (9)	C17—C16—C15	119.27 (17)
C8—N4—C8 ⁱ	120.79 (17)	C17—C16—H16	120.4
C13—N5—C9	116.30 (17)	C15—C16—H16	120.4
C18—N6—C14	118.07 (14)	C18—C17—C16	118.97 (16)
C18—N6—Ni1	125.01 (12)	C18—C17—H17	120.5
C14—N6—Ni1	115.18 (10)	C16—C17—H17	120.5
C19—N7—C19 ⁱ	122.07 (18)	N6—C18—C17	122.84 (16)
C19—N7—Ni1	118.97 (9)	N6—C18—H18	118.6
C19 ⁱ —N7—Ni1	118.97 (9)	C17—C18—H18	118.6
C20—N8—C20 ⁱ	120.13 (17)	N7—C19—C20	117.66 (13)
C21—N9—C25	116.73 (17)	N7—C19—C14	112.97 (13)
N2—C2—C3	121.62 (14)	C20—C19—C14	129.22 (13)

N2—C2—C7	113.95 (13)	N8—C20—C19	119.14 (13)
C3—C2—C7	124.12 (14)	N8—C20—C21	117.19 (13)
C2—C3—C4	119.29 (17)	C19—C20—C21	123.62 (13)
C2—C3—H3	120.4	N9—C21—C22	122.87 (15)
C4—C3—H3	120.4	N9—C21—C20	115.10 (14)
C5—C4—C3	119.08 (19)	C22—C21—C20	121.96 (15)
C5—C4—H4	120.5	C21—C22—C23	118.46 (18)
C3—C4—H4	120.5	C21—C22—H22	120.8
C6—C5—C4	118.71 (17)	C23—C22—H22	120.8
C6—C5—H5	120.6	C24—C23—C22	119.57 (19)
C4—C5—H5	120.6	C24—C23—H23	120.2
N2—C6—C5	122.94 (18)	C22—C23—H23	120.2
N2—C6—H6	118.5	C23—C24—C25	118.46 (18)
C5—C6—H6	118.5	C23—C24—H24	120.8
N3—C7—C8	117.85 (13)	C25—C24—H24	120.8
N3—C7—C2	113.09 (13)	N9—C25—C24	123.9 (2)
C8—C7—C2	128.96 (13)	N9—C25—H25	118
N4—C8—C7	119.10 (13)	C24—C25—H25	118
N4—C8—C9	116.22 (13)	H1W1—O1W—H1W2	108 (4)
C7—C8—C9	124.66 (13)		
N3—Ni1—N2—C6	167.18 (16)	C8 ⁱ —N4—C8—C7	-8.67 (10)
N7—Ni1—N2—C6	-12.82 (16)	C8 ⁱ —N4—C8—C9	170.04 (15)
N6 ⁱ —Ni1—N2—C6	-91.77 (16)	N3—C7—C8—N4	17.43 (19)
N6—Ni1—N2—C6	63.94 (16)	C2—C7—C8—N4	-158.78 (13)
N2 ⁱ —Ni1—N2—C6	167.18 (16)	N3—C7—C8—C9	-161.16 (13)
N3—Ni1—N2—C2	3.52 (10)	C2—C7—C8—C9	22.6 (2)
N7—Ni1—N2—C2	-176.48 (10)	C13—N5—C9—C10	1.5 (3)
N6 ⁱ —Ni1—N2—C2	104.58 (11)	C13—N5—C9—C8	178.73 (18)
N6—Ni1—N2—C2	-99.71 (11)	N4—C8—C9—N5	-143.30 (14)
N2 ⁱ —Ni1—N2—C2	3.52 (10)	C7—C8—C9—N5	35.3 (2)
N6 ⁱ —Ni1—N3—C7 ⁱ	89.60 (8)	N4—C8—C9—C10	34.0 (2)
N6—Ni1—N3—C7 ⁱ	-90.40 (8)	C7—C8—C9—C10	-147.32 (16)
N2 ⁱ —Ni1—N3—C7 ⁱ	4.94 (8)	N5—C9—C10—C11	-2.0 (3)
N2—Ni1—N3—C7 ⁱ	-175.06 (8)	C8—C9—C10—C11	-179.12 (16)
N6 ⁱ —Ni1—N3—C7	-90.40 (8)	C9—C10—C11—C12	0.9 (3)
N6—Ni1—N3—C7	89.60 (8)	C10—C11—C12—C13	0.5 (3)
N2 ⁱ —Ni1—N3—C7	-175.06 (8)	C9—N5—C13—C12	0.1 (4)
N2—Ni1—N3—C7	4.94 (8)	C11—C12—C13—N5	-1.0 (4)
N3—Ni1—N6—C18	-16.09 (14)	C18—N6—C14—C15	0.3 (2)
N7—Ni1—N6—C18	163.91 (14)	Ni1—N6—C14—C15	166.03 (13)
N6 ⁱ —Ni1—N6—C18	163.91 (14)	C18—N6—C14—C19	-171.54 (14)
N2 ⁱ —Ni1—N6—C18	-95.09 (14)	Ni1—N6—C14—C19	-5.83 (16)
N2—Ni1—N6—C18	60.72 (14)	N6—C14—C15—C16	-0.8 (3)
N3—Ni1—N6—C14	179.33 (10)	C19—C14—C15—C16	170.27 (16)
N7—Ni1—N6—C14	-0.67 (10)	C14—C15—C16—C17	-0.5 (3)
N6 ⁱ —Ni1—N6—C14	-0.67 (10)	C15—C16—C17—C18	2.1 (3)
N2 ⁱ —Ni1—N6—C14	100.33 (11)	C14—N6—C18—C17	1.4 (3)

N2—Ni1—N6—C14	-103.87 (11)	Ni1—N6—C18—C17	-162.77 (14)
N6 ⁱ —Ni1—N7—C19	-171.86 (8)	C16—C17—C18—N6	-2.6 (3)
N6—Ni1—N7—C19	8.14 (8)	C19 ⁱ —N7—C19—C20	-9.30 (10)
N2 ⁱ —Ni1—N7—C19	-87.20 (8)	Ni1—N7—C19—C20	170.70 (10)
N2—Ni1—N7—C19	92.80 (8)	C19 ⁱ —N7—C19—C14	166.59 (14)
N6 ⁱ —Ni1—N7—C19 ⁱ	8.14 (8)	Ni1—N7—C19—C14	-13.41 (14)
N6—Ni1—N7—C19 ⁱ	-171.86 (8)	N6—C14—C19—N7	12.20 (18)
N2 ⁱ —Ni1—N7—C19 ⁱ	92.80 (8)	C15—C14—C19—N7	-159.49 (14)
N2—Ni1—N7—C19 ⁱ	-87.20 (8)	N6—C14—C19—C20	-172.49 (15)
C6—N2—C2—C3	-1.4 (2)	C15—C14—C19—C20	15.8 (3)
Ni1—N2—C2—C3	163.47 (14)	C20 ⁱ —N8—C20—C19	-9.49 (10)
C6—N2—C2—C7	-175.20 (15)	C20 ⁱ —N8—C20—C21	168.10 (15)
Ni1—N2—C2—C7	-10.38 (17)	N7—C19—C20—N8	19.1 (2)
N2—C2—C3—C4	0.8 (3)	C14—C19—C20—N8	-156.03 (14)
C7—C2—C3—C4	174.00 (18)	N7—C19—C20—C21	-158.34 (13)
C2—C3—C4—C5	0.4 (3)	C14—C19—C20—C21	26.5 (2)
C3—C4—C5—C6	-0.9 (4)	C25—N9—C21—C22	0.8 (3)
C2—N2—C6—C5	0.8 (3)	C25—N9—C21—C20	177.73 (16)
Ni1—N2—C6—C5	-162.32 (18)	N8—C20—C21—N9	-140.33 (14)
C4—C5—C6—N2	0.4 (4)	C19—C20—C21—N9	37.1 (2)
C7 ⁱ —N3—C7—C8	-8.50 (10)	N8—C20—C21—C22	36.6 (2)
Ni1—N3—C7—C8	171.50 (10)	C19—C20—C21—C22	-145.93 (17)
C7 ⁱ —N3—C7—C2	168.30 (14)	N9—C21—C22—C23	-0.3 (3)
Ni1—N3—C7—C2	-11.70 (14)	C20—C21—C22—C23	-177.02 (16)
N2—C2—C7—N3	14.24 (18)	C21—C22—C23—C24	-0.1 (3)
C3—C2—C7—N3	-159.43 (15)	C22—C23—C24—C25	0.0 (3)
N2—C2—C7—C8	-169.40 (15)	C21—N9—C25—C24	-1.0 (3)
C3—C2—C7—C8	16.9 (3)	C23—C24—C25—N9	0.6 (3)

Symmetry code: (i) $-x, y, -z+1/2$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$O1W-H1W1\cdots N9^i$	0.85 (2)	2.28 (3)	3.116 (4)	166 (3)
$O1W-H1W2\cdots N1^{ii}$	0.85 (4)	2.20 (3)	3.044 (4)	173 (4)
$C3-H3\cdots N5$	0.93	2.61	3.045 (3)	109

Symmetry codes: (i) $-x, y, -z+1/2$; (ii) $-x+1/2, y+1/2, -z+1/2$.