

## Azido[1,2-bis(diphenylphosphanyl)-ethane- $\kappa^2P,P'$ ]( $\eta^5$ -indenyl)ruthenium(II)

Hui-Ling Sung,<sup>a\*</sup> Hsiu-Ling Hsu<sup>b</sup> and Ting-Shen Kuo<sup>c</sup>

<sup>a</sup>Department of Mathematics and Science (Pre-college), National Taiwan Normal University, Taiwan, <sup>b</sup>Department of Chemical and Materials Engineering, Lunghwa University of Science and Technology, Taiwan, and <sup>c</sup>Department of Chemistry, National Taiwan Normal University, 11677, Taiwan

Correspondence e-mail: hlsung@ntnu.edu.tw


Received 1 November 2010; accepted 17 December 2010

Key indicators: single-crystal X-ray study;  $T = 200\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.010\text{ \AA}$ ;  $R$  factor = 0.053;  $wR$  factor = 0.096; data-to-parameter ratio = 14.0.

Facile ligand substitution is observed when the ruthenium chloride complex  $[\text{Ru}(\eta^5\text{-C}_9\text{H}_7)\text{Cl}(\text{dppe})]$  (dppe is diphenylphosphanyl ethane) is treated with  $\text{NaN}_3$  in refluxing ethanol, yielding the title compound,  $[\text{Ru}(\eta^5\text{-C}_9\text{H}_7)(\text{N}_3)(\text{dppe})]$  or  $[\text{Ru}(\text{C}_9\text{H}_7)(\text{N}_3)(\text{C}_{26}\text{H}_{24}\text{P}_2)]$ . The Ru(II) atom has a typical piano-stool coordination. The Ru–P bond lengths are 2.284 (2) and 2.235 (2)  $\text{\AA}$ . NMR and MS analyses are in agreement with the structure of the title compound.

### Related literature

For the synthesis of the title compound, see: Singh *et al.* (2005). For the chemistry of organic azides, see: Labbe (1969); Patai (1971). For metal–azido complexes, see: Dori & Ziolo (1973); Frühauf (1997). Organic azides are particularly important for the synthesis of heterocyclic compounds by reaction with 1,3-dipole compounds, see: Padwa (1976). Metal–azido complexes have been reported to produce tetrazolates by reaction with nitrile and isonitriles, see: Beck & Schropp (1975); Ellis & Purcell (1982); Fehlhammer & Dahl (1972); Paul & Nag (1987); Treichel *et al.* (1971).



### Experimental

#### Crystal data

$[\text{Ru}(\text{C}_9\text{H}_7)(\text{N}_3)(\text{C}_{26}\text{H}_{24}\text{P}_2)]$	$V = 2934 (3)\text{ \AA}^3$
$M_r = 656.64$	$Z = 4$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 11.331 (6)\text{ \AA}$	$\mu = 0.67\text{ mm}^{-1}$
$b = 14.567 (9)\text{ \AA}$	$T = 200\text{ K}$
$c = 17.873 (11)\text{ \AA}$	$0.22 \times 0.10 \times 0.04\text{ mm}$
$\beta = 96.015 (19)^\circ$	

#### Data collection

Bruker Kappa APEXII CCD area-detector diffractometer	20374 measured reflections
Absorption correction: multi-scan ( <i>SADABS</i> ; Bruker, 2005)	5167 independent reflections
$T_{\min} = 0.866$ , $T_{\max} = 0.974$	2438 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.155$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.053$	370 parameters
$wR(F^2) = 0.096$	H-atom parameters constrained
$S = 0.75$	$\Delta\rho_{\max} = 0.54\text{ e \AA}^{-3}$
5167 reflections	$\Delta\rho_{\min} = -0.53\text{ e \AA}^{-3}$

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

We are grateful for financial support of this work by the National Science Council of the Republic of China (NSC Grant No. 97-2113-M-003-007-MY2) and the National Taiwan Normal University (99031012).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RN2075).

### References

- Beck, W. & Schropp, K. (1975). *Chem. Ber.* **108**, 3317–3325.
- Bruker (2005). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Dori, Z. & Ziolo, R. F. (1973). *Chem. Rev.* **73**, 247–254.
- Ellis, W. R. Jr & Purcell, W. L. (1982). *Inorg. Chem.* **21**, 834–837.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Fehlhammer, W. P. & Dahl, L. F. (1972). *J. Am. Chem. Soc.* **94**, 3370–3377.
- Frühauf, H. W. (1997). *Chem. Rev.* **97**, 523–596.
- Labbe, G. (1969). *Chem. Rev.* **69**, 345–363.
- Padwa, A. (1976). *Angew. Chem. Int. Ed. Engl.* **15**, 123–136.
- Patai, S. (1971). *The Chemistry of the Azido group*. New York: Interscience.
- Paul, P. & Nag, K. (1987). *Inorg. Chem.* **26**, 2969–2974.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Singh, K. S., Thöne, C. & Kollipara, M. R. (2005). *J. Organomet. Chem.* **690**, 4222–4231.
- Treichel, P. M., Knebel, W. J. & Hess, R. W. (1971). *J. Am. Chem. Soc.* **93**, 5424–5433.

# supporting information

*Acta Cryst.* (2011). E67, m124 [https://doi.org/10.1107/S1600536810053006]

## Azido[1,2-bis(diphenylphosphanyl)ethane- $\kappa^2P,P'$ ]( $\eta^5$ -indenyl)ruthenium(II)

Hui-Ling Sung, Hsiu-Ling Hsu and Ting-Shen Kuo

### S1. Comment

Organic azides are particularly important for synthesizing heterocyclic compounds by reaction with 1,3-dipole compounds (Padwa, 1976). Metal azido complexes have been reported to produce tetrazolates by reaction with nitrile (Paul & Nag, 1987; Ellis & Purcell, 1982) and isonitriles (Treichel *et al.*, 1971; Beck & Schropp, 1975; Fehlhammer & Dahl, 1972).


Treatment of the complex  $[\text{Ru}(\eta^5\text{-C}_9\text{H}_7)\text{Cl}(\text{dppe})]$  with sodium azide in ethanol afforded the title compound  $[\text{Ru}(\eta^5\text{-C}_9\text{H}_7)\text{N}_3(\text{dppe})]$  (Figure 1). In the crystal structure of the title compound, the azide groups are almost linear [ $\text{N}(3)=\text{N}(2)-\text{N}(1)=175.5$  (8) $^\circ$ ] and are coordinated to Ru with an Ru—N—N angle of 119.0 (5) $^\circ$ .

### S2. Experimental

To a solution of  $[\text{Ru}(\eta^5\text{-C}_9\text{H}_7)\text{Cl}(\text{dppe})]$  (0.1 g, 0.154 mmol) in ethanol (30 ml), an excess of  $\text{NaN}_3$  (0.05 g, 0.769 mmol) was added. The mixture was heated to reflux for 4 h and cooled to room temperature. The solvent was dried under vacuum and 10 ml of  $\text{CH}_2\text{Cl}_2$  was added to the residue. The product was dissolved in  $\text{CH}_2\text{Cl}_2$  and other salts such as  $\text{NaN}_3$  and  $\text{NaCl}$  precipitated. After filtration, the solvent of the mixture was concentrated to about 5 ml. The residue was then slowly added to 40 ml of vigorously stirred diethyl ether. The orange precipitate thus formed was filtered off, washed with diethyl ether and hexane and dried under vacuum to give the title compound  $[\text{Ru}(\eta^5\text{-C}_9\text{H}_7)\text{N}_3(\text{dppe})]$  (0.08 g, 0.122 mmol) in 79% yield. The orange crystals of the title compound for X-ray structure analysis were obtained by slow diffusion of diethyl ether into a  $\text{CH}_2\text{Cl}_2$  solution at room temperature for 3 days. Spectroscopic analysis:  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 298 K,  $\delta$ , p.p.m.): 7.44—7.20 (m, 24H, 20H of Ph group, 4H of indenyl ring), 4.91 (t, 1H,  $^3J_{\text{H}-\text{H}}=1.30$  Hz, H of indenyl ring), 5.51 (d, 2H,  $^3J_{\text{H}-\text{H}}=2.15$  Hz, H of indenyl ring), 2.38, 2.29 (m, 4H, 2 $\text{CH}_2$  of dppe).  $^{31}\text{P}\{^1\text{H}\}$  NMR ( $\text{CDCl}_3$ , 298 K,  $\delta$ , p.p.m.): 85.3.  $^{13}\text{C}\{^1\text{H}\}$  NMR ( $\text{CDCl}_3$ , 298 K,  $\delta$ , p.p.m.): 141—108 (Ph and indenyl group), 29.2 (t,  $J_{\text{C}-\text{P}}=22.64$  Hz,  $\text{CH}_2$  of dppe). HRMS (ESI,  $m/z$ ): 657.1 ( $M^+$ ), 615.3 ( $M^+-\text{N}_3$ ). Anal. Calcd for  $\text{C}_{35}\text{H}_{31}\text{N}_3\text{P}_2\text{Ru}$ : C, 64.02; H, 4.76; N, 6.40. Found: C, 64.16; H, 4.82; N, 6.28.

### S3. Refinement

All H atoms were initially located in a difference map, but were constrained to an idealized geometry. Constrained bond lengths and isotropic displacement parameters: C—H = 0.95 Å and  $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$  for aromatic H atoms, and C—H = 0.99 Å and  $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$  for methylene.

**Figure 1**

View of the title compound showing displacement ellipsoids at the 30% probability level. H atoms are omitted for clarity.

### Azido[1,2-bis(diphenylphosphanyl)ethane- $\kappa^2P,P'$ ]( $\eta^5$ -indenyl)ruthenium(II)

#### Crystal data



$M_r = 656.64$

Monoclinic,  $P2_1/c$

$a = 11.331 (6)$  Å

$b = 14.567 (9)$  Å

$c = 17.873 (11)$  Å

$\beta = 96.015 (19)^\circ$

$V = 2934 (3)$  Å<sup>3</sup>

$Z = 4$

$F(000) = 1344$

$D_x = 1.487 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 1167 reflections

$\theta = 2.6\text{--}19.4^\circ$

$\mu = 0.67 \text{ mm}^{-1}$

$T = 200$  K

Prism, orange-brown

$0.22 \times 0.10 \times 0.04$  mm

#### Data collection

Bruker Kappa APEXII CCD area-detector  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

Detector resolution: 9 pixels mm<sup>-1</sup>

CCD rotation images, thick slices scans

Absorption correction: multi-scan  
(SADABS; Bruker, 2005)

$T_{\min} = 0.866$ ,  $T_{\max} = 0.974$

20374 measured reflections

5167 independent reflections

2438 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.155$

$\theta_{\max} = 25.0^\circ$ ,  $\theta_{\min} = 1.8^\circ$

$h = -13 \rightarrow 10$

$k = -16 \rightarrow 17$

$l = -21 \rightarrow 20$

#### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.053$

$wR(F^2) = 0.096$

$S = 0.75$

5167 reflections

370 parameters

0 restraints

Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map

Hydrogen site location: inferred from  
neighbouring sites

H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0072P)^2]$   
 where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$   
 $\Delta\rho_{\max} = 0.54 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.53 \text{ e } \text{\AA}^{-3}$

### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.2833 (6)	-0.0987 (4)	0.6432 (4)	0.0325 (18)
C2	0.2129 (7)	-0.1041 (5)	0.7006 (5)	0.049 (2)
H2	0.2220	-0.0601	0.7401	0.059*
C3	0.1277 (7)	-0.1734 (6)	0.7021 (5)	0.059 (3)
H3	0.0776	-0.1764	0.7415	0.071*
C4	0.1185 (8)	-0.2362 (6)	0.6459 (6)	0.068 (3)
H4	0.0606	-0.2834	0.6457	0.082*
C5	0.1902 (8)	-0.2329 (5)	0.5900 (5)	0.064 (3)
H5	0.1836	-0.2790	0.5522	0.077*
C6	0.2728 (6)	-0.1636 (4)	0.5869 (4)	0.043 (2)
H6	0.3214	-0.1608	0.5467	0.052*
C7	0.3040 (6)	0.0764 (4)	0.5736 (3)	0.0283 (17)
C8	0.1933 (6)	0.0576 (5)	0.5384 (4)	0.0391 (18)
H8	0.1545	0.0022	0.5497	0.047*
C9	0.1373 (7)	0.1183 (5)	0.4863 (4)	0.051 (2)
H9	0.0611	0.1040	0.4616	0.062*
C10	0.1919 (7)	0.1986 (5)	0.4707 (4)	0.042 (2)
H10	0.1531	0.2403	0.4354	0.050*
C11	0.3001 (7)	0.2193 (4)	0.5050 (4)	0.042 (2)
H11	0.3376	0.2751	0.4932	0.050*
C12	0.3570 (6)	0.1599 (4)	0.5572 (4)	0.040 (2)
H12	0.4324	0.1758	0.5822	0.048*
C13	0.5126 (5)	-0.0399 (4)	0.5971 (4)	0.0340 (18)
H13A	0.4876	-0.0918	0.5633	0.041*
H13B	0.5410	0.0103	0.5663	0.041*
C14	0.6120 (5)	-0.0706 (4)	0.6553 (4)	0.0317 (17)
H14A	0.6859	-0.0807	0.6315	0.038*
H14B	0.5904	-0.1286	0.6794	0.038*
C15	0.7216 (6)	0.1045 (4)	0.6777 (4)	0.0324 (18)
C16	0.6805 (7)	0.1910 (4)	0.6570 (4)	0.039 (2)

H16	0.6040	0.2091	0.6688	0.047*
C17	0.7460 (7)	0.2512 (5)	0.6202 (4)	0.055 (2)
H17	0.7153	0.3103	0.6069	0.066*
C18	0.8549 (8)	0.2266 (6)	0.6024 (4)	0.057 (3)
H18	0.9004	0.2688	0.5768	0.069*
C19	0.9000 (7)	0.1414 (6)	0.6211 (4)	0.058 (2)
H19	0.9761	0.1244	0.6079	0.070*
C20	0.8349 (6)	0.0805 (5)	0.6592 (4)	0.045 (2)
H20	0.8669	0.0219	0.6730	0.053*
C21	0.7412 (6)	-0.0228 (4)	0.7992 (4)	0.0316 (18)
C22	0.7937 (6)	-0.1062 (4)	0.7984 (5)	0.046 (2)
H22	0.7785	-0.1444	0.7554	0.055*
C23	0.8685 (7)	-0.1366 (5)	0.8588 (5)	0.062 (3)
H23	0.9070	-0.1943	0.8565	0.075*
C24	0.8876 (6)	-0.0838 (5)	0.9224 (4)	0.050 (2)
H24	0.9376	-0.1055	0.9647	0.060*
C25	0.8347 (6)	-0.0003 (5)	0.9246 (4)	0.043 (2)
H25	0.8478	0.0368	0.9684	0.052*
C26	0.7628 (6)	0.0299 (4)	0.8635 (4)	0.041 (2)
H26	0.7266	0.0886	0.8653	0.049*
C27	0.3015 (7)	0.1547 (4)	0.7838 (4)	0.0366 (19)
C28	0.1798 (7)	0.1421 (4)	0.7569 (4)	0.043 (2)
H28	0.1481	0.1675	0.7101	0.051*
C29	0.1103 (7)	0.0932 (5)	0.7995 (5)	0.050 (2)
H29	0.0285	0.0858	0.7826	0.060*
C30	0.1562 (6)	0.0527 (5)	0.8687 (4)	0.045 (2)
H30	0.1052	0.0183	0.8970	0.054*
C31	0.2735 (6)	0.0626 (5)	0.8951 (4)	0.0415 (18)
H31	0.3040	0.0346	0.9411	0.050*
C32	0.3488 (6)	0.1149 (4)	0.8534 (4)	0.0294 (17)
C33	0.4714 (6)	0.1389 (4)	0.8655 (4)	0.0319 (18)
H33	0.5261	0.1246	0.9116	0.038*
C34	0.4981 (6)	0.1950 (4)	0.8056 (4)	0.035 (2)
H34	0.5756	0.2270	0.8028	0.042*
C35	0.4014 (7)	0.2019 (4)	0.7521 (4)	0.038 (2)
H35	0.3957	0.2409	0.7058	0.046*
N1	0.4613 (5)	-0.0796 (3)	0.8004 (3)	0.0340 (16)
N2	0.4855 (6)	-0.0931 (4)	0.8660 (4)	0.0495 (19)
N3	0.5048 (7)	-0.1117 (5)	0.9303 (4)	0.082 (3)
P1	0.38539 (16)	0.00036 (11)	0.64504 (10)	0.0282 (5)
P2	0.63247 (16)	0.02162 (11)	0.72512 (11)	0.0288 (5)
Ru1	0.45473 (5)	0.05865 (3)	0.75994 (3)	0.02632 (16)

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.027 (5)	0.030 (4)	0.037 (5)	-0.005 (3)	-0.011 (4)	0.006 (3)
C2	0.043 (6)	0.047 (5)	0.058 (6)	-0.004 (4)	0.001 (5)	-0.003 (4)

C3	0.044 (6)	0.069 (6)	0.062 (7)	-0.017 (5)	-0.007 (5)	0.029 (5)
C4	0.065 (7)	0.052 (6)	0.082 (9)	-0.038 (5)	-0.024 (6)	0.029 (5)
C5	0.082 (8)	0.045 (5)	0.061 (7)	-0.022 (5)	-0.018 (6)	0.005 (5)
C6	0.057 (6)	0.029 (4)	0.041 (6)	-0.003 (4)	-0.009 (4)	-0.001 (4)
C7	0.037 (5)	0.028 (4)	0.021 (4)	0.002 (3)	0.006 (4)	-0.002 (3)
C8	0.031 (5)	0.035 (4)	0.049 (5)	-0.004 (4)	-0.009 (4)	0.008 (4)
C9	0.045 (6)	0.051 (5)	0.057 (6)	-0.001 (4)	0.001 (5)	0.001 (4)
C10	0.047 (6)	0.042 (5)	0.034 (5)	0.017 (4)	-0.003 (4)	0.007 (4)
C11	0.061 (6)	0.024 (4)	0.038 (5)	0.001 (4)	0.001 (5)	0.008 (3)
C12	0.044 (5)	0.030 (4)	0.043 (5)	-0.003 (4)	-0.004 (4)	0.004 (4)
C13	0.032 (5)	0.031 (4)	0.040 (5)	-0.003 (3)	0.007 (4)	-0.006 (3)
C14	0.024 (4)	0.032 (4)	0.038 (5)	0.001 (3)	0.001 (4)	-0.007 (4)
C15	0.025 (5)	0.045 (4)	0.027 (5)	-0.012 (3)	0.006 (4)	-0.006 (3)
C16	0.048 (6)	0.039 (4)	0.030 (5)	-0.014 (4)	0.004 (4)	0.005 (4)
C17	0.049 (6)	0.062 (5)	0.053 (6)	-0.021 (5)	0.001 (5)	0.008 (4)
C18	0.045 (6)	0.079 (6)	0.045 (6)	-0.036 (5)	-0.009 (5)	0.020 (5)
C19	0.020 (5)	0.105 (7)	0.049 (6)	-0.018 (5)	0.006 (4)	-0.003 (5)
C20	0.032 (5)	0.055 (5)	0.048 (5)	-0.004 (4)	0.013 (4)	-0.007 (4)
C21	0.017 (4)	0.035 (4)	0.043 (5)	0.000 (3)	0.004 (4)	0.000 (3)
C22	0.034 (5)	0.032 (4)	0.066 (7)	0.004 (4)	-0.020 (5)	-0.012 (4)
C23	0.054 (6)	0.034 (5)	0.091 (8)	0.020 (4)	-0.027 (6)	-0.008 (5)
C24	0.040 (5)	0.058 (6)	0.048 (6)	-0.011 (4)	-0.010 (4)	-0.002 (4)
C25	0.033 (5)	0.043 (5)	0.051 (6)	0.004 (4)	-0.005 (4)	-0.005 (4)
C26	0.044 (5)	0.036 (4)	0.041 (5)	0.017 (4)	-0.001 (4)	0.001 (4)
C27	0.034 (5)	0.031 (4)	0.046 (6)	0.013 (4)	0.010 (4)	-0.007 (4)
C28	0.040 (6)	0.044 (5)	0.043 (6)	0.008 (4)	-0.001 (5)	-0.004 (4)
C29	0.021 (5)	0.065 (6)	0.062 (7)	0.000 (4)	0.000 (5)	-0.023 (5)
C30	0.035 (5)	0.054 (5)	0.050 (6)	-0.002 (4)	0.020 (4)	-0.012 (5)
C31	0.037 (5)	0.055 (4)	0.033 (5)	-0.001 (4)	0.006 (4)	-0.012 (4)
C32	0.031 (5)	0.037 (4)	0.019 (4)	0.005 (3)	-0.005 (4)	-0.003 (3)
C33	0.030 (5)	0.042 (4)	0.022 (5)	0.000 (3)	-0.006 (4)	-0.009 (3)
C34	0.026 (5)	0.029 (4)	0.050 (6)	-0.004 (3)	0.006 (4)	-0.018 (4)
C35	0.046 (5)	0.018 (4)	0.053 (6)	0.005 (3)	0.017 (5)	-0.002 (3)
N1	0.035 (4)	0.037 (4)	0.028 (4)	-0.001 (3)	-0.006 (3)	0.005 (3)
N2	0.056 (5)	0.036 (4)	0.056 (6)	0.003 (3)	0.003 (5)	0.013 (4)
N3	0.103 (7)	0.076 (5)	0.065 (6)	0.007 (4)	-0.004 (6)	0.023 (5)
P1	0.0299 (12)	0.0245 (10)	0.0298 (12)	0.0000 (8)	0.0008 (10)	0.0009 (8)
P2	0.0272 (12)	0.0289 (9)	0.0303 (13)	-0.0012 (8)	0.0024 (10)	-0.0014 (9)
Ru1	0.0255 (3)	0.0245 (3)	0.0284 (3)	-0.0003 (3)	0.0004 (2)	-0.0020 (3)

Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )

C1—C2	1.367 (8)	C19—H19	0.9500
C1—C6	1.376 (8)	C20—H20	0.9500
C1—P1	1.847 (6)	C21—C22	1.354 (8)
C2—C3	1.400 (9)	C21—C26	1.383 (8)
C2—H2	0.9500	C21—P2	1.830 (7)
C3—C4	1.354 (10)	C22—C23	1.374 (9)

C3—H3	0.9500	C22—H22	0.9500
C4—C5	1.354 (10)	C23—C24	1.371 (9)
C4—H4	0.9500	C23—H23	0.9500
C5—C6	1.382 (9)	C24—C25	1.358 (8)
C5—H5	0.9500	C24—H24	0.9500
C6—H6	0.9500	C25—C26	1.365 (9)
C7—C8	1.371 (8)	C25—H25	0.9500
C7—C12	1.400 (8)	C26—H26	0.9500
C7—P1	1.861 (7)	C27—C28	1.423 (9)
C8—C9	1.388 (9)	C27—C32	1.425 (9)
C8—H8	0.9500	C27—C35	1.487 (9)
C9—C10	1.366 (9)	C27—Ru1	2.304 (6)
C9—H9	0.9500	C28—C29	1.354 (9)
C10—C11	1.346 (9)	C28—H28	0.9500
C10—H10	0.9500	C29—C30	1.419 (9)
C11—C12	1.382 (8)	C29—H29	0.9500
C11—H11	0.9500	C30—C31	1.370 (9)
C12—H12	0.9500	C30—H30	0.9500
C13—C14	1.517 (8)	C31—C32	1.412 (8)
C13—P1	1.848 (6)	C31—H31	0.9500
C13—H13A	0.9900	C32—C33	1.426 (8)
C13—H13B	0.9900	C32—Ru1	2.307 (6)
C14—P2	1.832 (6)	C33—C34	1.406 (8)
C14—H14A	0.9900	C33—Ru1	2.210 (6)
C14—H14B	0.9900	C33—H33	1.0000
C15—C16	1.380 (8)	C34—C35	1.380 (9)
C15—C20	1.403 (8)	C34—Ru1	2.184 (6)
C15—P2	1.837 (6)	C34—H34	1.0000
C16—C17	1.362 (8)	C35—Ru1	2.173 (6)
C16—H16	0.9500	C35—H35	1.0000
C17—C18	1.354 (9)	N1—N2	1.192 (8)
C17—H17	0.9500	N1—Ru1	2.139 (5)
C18—C19	1.369 (9)	N2—N3	1.178 (8)
C18—H18	0.9500	P1—Ru1	2.284 (2)
C19—C20	1.379 (8)	P2—Ru1	2.235 (2)
C2—C1—C6	119.8 (7)	C28—C27—C32	120.3 (7)
C2—C1—P1	116.4 (5)	C28—C27—C35	132.9 (7)
C6—C1—P1	123.7 (6)	C32—C27—C35	106.8 (7)
C1—C2—C3	120.9 (8)	C28—C27—Ru1	125.5 (5)
C1—C2—H2	119.5	C32—C27—Ru1	72.1 (4)
C3—C2—H2	119.5	C35—C27—Ru1	65.9 (3)
C4—C3—C2	118.2 (8)	C29—C28—C27	118.6 (7)
C4—C3—H3	120.9	C29—C28—H28	120.7
C2—C3—H3	120.9	C27—C28—H28	120.7
C5—C4—C3	121.3 (8)	C28—C29—C30	121.8 (7)
C5—C4—H4	119.4	C28—C29—H29	119.1
C3—C4—H4	119.4	C30—C29—H29	119.1

C4—C5—C6	121.1 (8)	C31—C30—C29	120.7 (7)
C4—C5—H5	119.5	C31—C30—H30	119.7
C6—C5—H5	119.5	C29—C30—H30	119.7
C1—C6—C5	118.7 (7)	C30—C31—C32	119.5 (7)
C1—C6—H6	120.7	C30—C31—H31	120.2
C5—C6—H6	120.7	C32—C31—H31	120.2
C8—C7—C12	118.0 (6)	C31—C32—C27	119.2 (7)
C8—C7—P1	124.0 (5)	C31—C32—C33	133.2 (7)
C12—C7—P1	117.9 (5)	C27—C32—C33	107.6 (6)
C7—C8—C9	120.8 (7)	C31—C32—Ru1	125.4 (4)
C7—C8—H8	119.6	C27—C32—Ru1	71.9 (4)
C9—C8—H8	119.6	C33—C32—Ru1	67.9 (4)
C10—C9—C8	119.8 (8)	C34—C33—C32	108.2 (6)
C10—C9—H9	120.1	C34—C33—Ru1	70.4 (4)
C8—C9—H9	120.1	C32—C33—Ru1	75.4 (4)
C11—C10—C9	120.6 (7)	C34—C33—H33	125.7
C11—C10—H10	119.7	C32—C33—H33	125.7
C9—C10—H10	119.7	Ru1—C33—H33	125.7
C10—C11—C12	120.3 (7)	C35—C34—C33	110.5 (7)
C10—C11—H11	119.8	C35—C34—Ru1	71.1 (4)
C12—C11—H11	119.8	C33—C34—Ru1	72.3 (4)
C11—C12—C7	120.3 (7)	C35—C34—H34	124.7
C11—C12—H12	119.8	C33—C34—H34	124.7
C7—C12—H12	119.8	Ru1—C34—H34	124.7
C14—C13—P1	109.6 (4)	C34—C35—C27	106.5 (7)
C14—C13—H13A	109.8	C34—C35—Ru1	72.0 (4)
P1—C13—H13A	109.8	C27—C35—Ru1	75.5 (3)
C14—C13—H13B	109.8	C34—C35—H35	126.3
P1—C13—H13B	109.8	C27—C35—H35	126.3
H13A—C13—H13B	108.2	Ru1—C35—H35	126.3
C13—C14—P2	106.4 (4)	N2—N1—Ru1	119.0 (5)
C13—C14—H14A	110.4	N3—N2—N1	175.5 (8)
P2—C14—H14A	110.4	C13—P1—C1	105.1 (3)
C13—C14—H14B	110.4	C13—P1—C7	103.0 (3)
P2—C14—H14B	110.4	C1—P1—C7	100.8 (3)
H14A—C14—H14B	108.6	C13—P1—Ru1	108.8 (2)
C16—C15—C20	117.1 (6)	C1—P1—Ru1	117.5 (2)
C16—C15—P2	122.5 (5)	C7—P1—Ru1	119.7 (2)
C20—C15—P2	120.4 (5)	C21—P2—C14	105.1 (3)
C17—C16—C15	122.1 (7)	C21—P2—C15	101.8 (3)
C17—C16—H16	119.0	C14—P2—C15	101.8 (3)
C15—C16—H16	119.0	C21—P2—Ru1	116.2 (2)
C18—C17—C16	119.9 (8)	C14—P2—Ru1	108.2 (2)
C18—C17—H17	120.0	C15—P2—Ru1	121.7 (2)
C16—C17—H17	120.0	N1—Ru1—C35	157.8 (2)
C17—C18—C19	120.6 (7)	N1—Ru1—C34	137.1 (3)
C17—C18—H18	119.7	C35—Ru1—C34	36.9 (2)
C19—C18—H18	119.7	N1—Ru1—C33	102.3 (2)

C18—C19—C20	119.8 (7)	C35—Ru1—C33	63.0 (3)
C18—C19—H19	120.1	C34—Ru1—C33	37.3 (2)
C20—C19—H19	120.1	N1—Ru1—P2	82.33 (15)
C19—C20—C15	120.4 (7)	C35—Ru1—P2	117.78 (19)
C19—C20—H20	119.8	C34—Ru1—P2	98.59 (19)
C15—C20—H20	119.8	C33—Ru1—P2	111.50 (18)
C22—C21—C26	117.9 (7)	N1—Ru1—P1	87.17 (16)
C22—C21—P2	125.0 (6)	C35—Ru1—P1	103.4 (2)
C26—C21—P2	116.9 (5)	C34—Ru1—P1	135.8 (2)
C21—C22—C23	121.1 (7)	C33—Ru1—P1	162.63 (18)
C21—C22—H22	119.4	P2—Ru1—P1	83.97 (7)
C23—C22—H22	119.4	N1—Ru1—C27	120.6 (2)
C22—C23—C24	120.1 (7)	C35—Ru1—C27	38.6 (2)
C22—C23—H23	119.9	C34—Ru1—C27	61.5 (2)
C24—C23—H23	119.9	C33—Ru1—C27	61.3 (3)
C25—C24—C23	119.6 (8)	P2—Ru1—C27	156.39 (18)
C25—C24—H24	120.2	P1—Ru1—C27	101.4 (2)
C23—C24—H24	120.2	N1—Ru1—C32	95.3 (2)
C24—C25—C26	119.7 (7)	C35—Ru1—C32	62.8 (2)
C24—C25—H25	120.2	C34—Ru1—C32	61.3 (2)
C26—C25—H25	120.2	C33—Ru1—C32	36.7 (2)
C25—C26—C21	121.6 (6)	P2—Ru1—C32	147.10 (19)
C25—C26—H26	119.2	P1—Ru1—C32	128.82 (19)
C21—C26—H26	119.2	C27—Ru1—C32	36.0 (2)