

Bis(*N*-nitroso-*N*-phenylhydroxylaminato- κ^2O,O')(1,10-phenanthroline- κ^2N,N')-lead(II)

Ezzatollah Najafi,^a Mostafa M. Amini^a and Seik Weng Ng^{b*}

^aDepartment of Chemistry, General Campus, Shahid Beheshti University, Tehran 1983963113, Iran, and ^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: seikweng@um.edu.my


Received 20 February 2011; accepted 22 February 2011

Key indicators: single-crystal X-ray study; $T = 100\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.018; wR factor = 0.047; data-to-parameter ratio = 16.6.

The two cupferronate ions and the *N*-heterocycle in the mononuclear title compound, $[\text{Pb}(\text{C}_6\text{H}_5\text{N}_2\text{O}_2)_2(\text{C}_{12}\text{H}_8\text{N}_2)]$, O,O' - and N,N' -chelate to the Pb^{II} atom, the geometry of which is a distorted Ψ -pentagonal bipyramid.

Related literature

For the structure of dinuclear $[\text{Pb}(\text{C}_6\text{H}_5\text{N}_2\text{O}_2)_2]_2$, see: Najafi *et al.* (2011).



Experimental

Crystal data

$[\text{Pb}(\text{C}_6\text{H}_5\text{N}_2\text{O}_2)_2(\text{C}_{12}\text{H}_8\text{N}_2)]$	$V = 2285.7 (2)\text{ \AA}^3$
$M_r = 661.63$	$Z = 4$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 7.7033 (4)\text{ \AA}$	$\mu = 7.43\text{ mm}^{-1}$
$b = 15.9948 (8)\text{ \AA}$	$T = 100\text{ K}$
$c = 18.8929 (10)\text{ \AA}$	$0.20 \times 0.10 \times 0.10\text{ mm}$
$\beta = 100.919 (1)^{\circ}$	

Data collection

Bruker SMART APEX	21418 measured reflections
diffractometer	5233 independent reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 1996)	4676 reflections with $I > 2\sigma(I)$
$T_{\min} = 0.318$, $T_{\max} = 0.524$	$R_{\text{int}} = 0.028$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.018$	316 parameters
$wR(F^2) = 0.047$	H-atom parameters constrained
$S = 0.91$	$\Delta\rho_{\max} = 0.74\text{ e \AA}^{-3}$
5233 reflections	$\Delta\rho_{\min} = -0.53\text{ e \AA}^{-3}$

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2010).

We thank Shahid Beheshti University and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5481).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Bruker (2009). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Najafi, E., Amini, M. M. & Ng, S. W. (2011). *Acta Cryst. E67*, m377.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2011). E67, m378 [doi:10.1107/S1600536811006787]

Bis(*N*-nitroso-*N*-phenylhydroxylaminato- κ^2O,O')(1,10-phenanthroline- κ^2N,N')lead(II)

Ezzatollah Najafi, Mostafa M. Amini and Seik Weng Ng

S1. Comment

The cupferronate ion is a common ion used for the complexation of metals; the lead(II) derivative exists as a dinuclear compound, the four cupferronate ions in dinuclear $[Pb(C_6H_5N_2O_2)_2]_2$ O,O' -chelate to the lead(II) atom, and two of the four nitroso O atoms are also involved in bridging. The geometry of both five-coordinate lead atoms is Ψ -octahedral; if another longer intermolecular $Pb\cdots O$ interactions (approx. 3.0 Å) are considered, the geometry is a Ψ -square-antiprism (Najafi *et al.*, 2011). The 1,10-phenanthroline adduct is monomeric (Scheme I, Fig. 1). The two cupferronate ions and the N-heterocycle in mononuclear $Pb(C_{12}H_8N_2)(C_6H_5N_2O_2)_2$ chelate to the lead(II) atom; the geometry of the lead atom is a Ψ -pentagonal bipyramid.


S2. Experimental

Lead(II) nitrate (0.33 g, 1 mmol) dissolved in ethanol (20 ml) was added to the cupferron ligand (0.31 g, 2 mmol) and 1,10-phenanthroline hydrate (0.40, 2 mmol) dissolved in ethanol (20 ml). The mixture was stirred and then set aside for the growth of brown colored crystals.

S3. Refinement

Hydrogen atoms were placed in calculated positions (C–H 0.95 Å) and were included in the refinement in the riding model approximation, with $U(H)$ set to $1.2U_{eq}(C)$.


Omitted from the refinement were the following reflections owing to bad disagreement between the observed and calculated F^2 values: (0 0 1), (0 1 2), (1 0 1), (0 0 2), (11 4 7), (-9 - 11 5), (11 3 8), (11 5 6), (-4 - 9 10), (-9 - 9 2) and (3 - 2 14).

**Figure 1**

Anisotropic displacement ellipsoid plot (Barbour, 2001) of $\text{Pb}(\text{C}_{12}\text{H}_8\text{N}_2)(\text{C}_6\text{H}_5\text{N}_2\text{O}_2)_2$ at the 70% probability level. Hydrogen atoms are drawn as spheres of arbitrary radius.

Bis(*N*-nitroso-*N*-phenylhydroxylaminato- $\kappa^2\text{O},\text{O}'$)(1,10-phenanthroline- $\kappa^2\text{N},\text{N}'$)lead(II)

Crystal data



$M_r = 661.63$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 7.7033$ (4) Å

$b = 15.9948$ (8) Å

$c = 18.8929$ (10) Å

$\beta = 100.919$ (1)°

$V = 2285.7$ (2) Å³

$Z = 4$

$F(000) = 1272$

$D_x = 1.923 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 9896 reflections

$\theta = 2.2\text{--}28.3$ °

$\mu = 7.43 \text{ mm}^{-1}$

$T = 100$ K

Prism, brown

0.20 × 0.10 × 0.10 mm

Data collection

Bruker SMART APEX
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

ω scans

Absorption correction: multi-scan
(SADABS; Sheldrick, 1996)

$T_{\min} = 0.318$, $T_{\max} = 0.524$

21418 measured reflections

5233 independent reflections

4676 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.028$
 $\theta_{\text{max}} = 27.5^\circ$, $\theta_{\text{min}} = 2.2^\circ$

$h = -9 \rightarrow 10$
 $k = -20 \rightarrow 18$
 $l = -24 \rightarrow 24$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.018$
 $wR(F^2) = 0.047$
 $S = 0.91$
5233 reflections
316 parameters
0 restraints
Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map
Hydrogen site location: inferred from neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0315P)^2 + 1.0382P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.001$
 $\Delta\rho_{\text{max}} = 0.74 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.53 \text{ e } \text{\AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Pb1	0.545412 (12)	0.564925 (6)	0.707994 (5)	0.01407 (4)
O1	0.6518 (3)	0.57008 (11)	0.83910 (10)	0.0215 (4)
O2	0.6004 (3)	0.43217 (11)	0.77043 (10)	0.0221 (4)
O3	0.3666 (2)	0.67625 (11)	0.74664 (10)	0.0171 (4)
O4	0.2682 (2)	0.52341 (11)	0.72411 (10)	0.0168 (4)
N1	0.6651 (3)	0.49786 (14)	0.87218 (11)	0.0167 (4)
N2	0.6420 (3)	0.42654 (14)	0.84043 (12)	0.0207 (5)
N3	0.2048 (3)	0.65180 (13)	0.74731 (10)	0.0123 (4)
N4	0.1479 (3)	0.57674 (12)	0.73517 (11)	0.0139 (4)
N5	0.4379 (3)	0.44824 (13)	0.60458 (12)	0.0156 (4)
N6	0.3223 (3)	0.61102 (14)	0.58176 (11)	0.0160 (4)
C1	0.7119 (4)	0.49851 (17)	0.94969 (14)	0.0189 (5)
C2	0.6692 (4)	0.43069 (17)	0.98924 (15)	0.0228 (6)
H2	0.6111	0.3831	0.9657	0.027*
C3	0.7135 (5)	0.43430 (18)	1.06373 (16)	0.0298 (7)
H3	0.6877	0.3882	1.0917	0.036*
C4	0.7955 (5)	0.5050 (2)	1.09786 (15)	0.0355 (8)
H4	0.8229	0.5075	1.1490	0.043*
C5	0.8371 (5)	0.57181 (19)	1.05735 (16)	0.0345 (8)
H5	0.8947	0.6196	1.0808	0.041*
C6	0.7950 (4)	0.56902 (18)	0.98277 (15)	0.0254 (6)
H6	0.8225	0.6147	0.9548	0.031*
C7	0.0758 (3)	0.71352 (15)	0.75744 (13)	0.0132 (5)
C8	-0.0737 (3)	0.68857 (16)	0.78329 (13)	0.0159 (5)
H8	-0.0865	0.6325	0.7980	0.019*
C9	-0.2042 (4)	0.74779 (17)	0.78711 (14)	0.0193 (5)
H9	-0.3089	0.7319	0.8034	0.023*
C10	-0.1817 (4)	0.83009 (17)	0.76720 (14)	0.0211 (6)
H10	-0.2722	0.8701	0.7686	0.025*
C11	-0.0270 (4)	0.85391 (16)	0.74523 (14)	0.0187 (5)
H11	-0.0099	0.9108	0.7338	0.022*

C12	0.1030 (3)	0.79543 (15)	0.73976 (13)	0.0156 (5)
H12	0.2084	0.8115	0.7242	0.019*
C13	0.4890 (4)	0.36929 (17)	0.61474 (14)	0.0195 (6)
H13	0.5640	0.3550	0.6590	0.023*
C14	0.4390 (4)	0.30593 (17)	0.56419 (14)	0.0213 (6)
H14	0.4786	0.2502	0.5743	0.026*
C15	0.3320 (4)	0.32538 (17)	0.49995 (14)	0.0202 (6)
H15	0.2974	0.2832	0.4647	0.024*
C16	0.2737 (3)	0.40803 (17)	0.48634 (14)	0.0174 (5)
C17	0.3300 (3)	0.46863 (17)	0.54094 (13)	0.0153 (5)
C18	0.1582 (4)	0.43202 (17)	0.42116 (14)	0.0204 (6)
H18	0.1205	0.3912	0.3850	0.024*
C19	0.1015 (4)	0.51195 (18)	0.41007 (14)	0.0198 (6)
H19	0.0243	0.5263	0.3664	0.024*
C20	0.1563 (4)	0.57524 (16)	0.46329 (14)	0.0177 (5)
C21	0.2698 (3)	0.55380 (16)	0.52921 (13)	0.0156 (5)
C22	0.0991 (4)	0.65833 (18)	0.45407 (14)	0.0210 (6)
H22	0.0237	0.6750	0.4106	0.025*
C23	0.1520 (4)	0.71536 (18)	0.50770 (14)	0.0217 (6)
H23	0.1136	0.7718	0.5021	0.026*
C24	0.2643 (3)	0.68889 (17)	0.57130 (14)	0.0194 (5)
H24	0.3002	0.7287	0.6085	0.023*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Pb1	0.01178 (6)	0.01599 (6)	0.01461 (5)	-0.00027 (4)	0.00298 (4)	-0.00125 (3)
O1	0.0316 (12)	0.0155 (10)	0.0164 (9)	-0.0056 (8)	0.0021 (8)	0.0018 (7)
O2	0.0287 (12)	0.0188 (10)	0.0172 (9)	0.0091 (8)	0.0003 (8)	-0.0034 (7)
O3	0.0097 (8)	0.0152 (9)	0.0267 (9)	-0.0040 (7)	0.0043 (7)	-0.0032 (7)
O4	0.0155 (9)	0.0112 (9)	0.0250 (9)	0.0015 (7)	0.0073 (7)	-0.0010 (7)
N1	0.0163 (11)	0.0166 (11)	0.0173 (10)	0.0003 (9)	0.0031 (8)	0.0008 (9)
N2	0.0243 (13)	0.0184 (12)	0.0181 (11)	0.0058 (9)	0.0008 (9)	-0.0022 (9)
N3	0.0109 (10)	0.0117 (10)	0.0137 (9)	0.0006 (8)	0.0006 (8)	0.0003 (8)
N4	0.0140 (11)	0.0102 (10)	0.0175 (10)	-0.0011 (8)	0.0033 (8)	0.0001 (8)
N5	0.0143 (11)	0.0172 (11)	0.0162 (10)	0.0009 (9)	0.0049 (9)	0.0002 (8)
N6	0.0139 (11)	0.0187 (11)	0.0160 (10)	0.0012 (9)	0.0047 (8)	0.0001 (9)
C1	0.0192 (14)	0.0211 (14)	0.0157 (12)	0.0002 (11)	0.0015 (10)	0.0006 (10)
C2	0.0258 (16)	0.0186 (14)	0.0225 (14)	-0.0020 (11)	0.0004 (11)	0.0013 (11)
C3	0.043 (2)	0.0256 (16)	0.0197 (14)	-0.0043 (14)	0.0019 (13)	0.0061 (11)
C4	0.058 (2)	0.0329 (18)	0.0131 (13)	-0.0125 (16)	-0.0011 (13)	0.0020 (12)
C5	0.053 (2)	0.0283 (17)	0.0179 (14)	-0.0126 (15)	-0.0028 (14)	-0.0020 (12)
C6	0.0319 (17)	0.0239 (16)	0.0197 (13)	-0.0081 (12)	0.0031 (12)	0.0041 (11)
C7	0.0126 (12)	0.0131 (12)	0.0128 (11)	0.0004 (9)	-0.0002 (9)	-0.0021 (9)
C8	0.0152 (13)	0.0117 (12)	0.0204 (12)	-0.0026 (10)	0.0025 (10)	-0.0019 (10)
C9	0.0158 (13)	0.0186 (14)	0.0237 (13)	-0.0008 (11)	0.0041 (11)	-0.0066 (10)
C10	0.0202 (14)	0.0175 (13)	0.0249 (13)	0.0068 (11)	0.0023 (11)	-0.0047 (11)
C11	0.0245 (15)	0.0108 (12)	0.0193 (12)	0.0002 (11)	0.0007 (11)	-0.0009 (10)

C12	0.0167 (13)	0.0130 (12)	0.0163 (11)	-0.0010 (10)	0.0010 (10)	-0.0005 (10)
C13	0.0178 (14)	0.0215 (14)	0.0202 (13)	0.0035 (11)	0.0059 (11)	-0.0002 (11)
C14	0.0225 (14)	0.0148 (13)	0.0288 (14)	0.0008 (11)	0.0100 (11)	-0.0006 (11)
C15	0.0197 (14)	0.0196 (14)	0.0230 (13)	-0.0040 (11)	0.0082 (11)	-0.0043 (11)
C16	0.0140 (13)	0.0218 (13)	0.0186 (12)	-0.0045 (11)	0.0085 (10)	-0.0022 (10)
C17	0.0101 (12)	0.0196 (13)	0.0175 (12)	-0.0046 (10)	0.0059 (10)	-0.0019 (10)
C18	0.0192 (14)	0.0249 (15)	0.0180 (12)	-0.0082 (11)	0.0060 (11)	-0.0036 (11)
C19	0.0163 (13)	0.0294 (15)	0.0135 (12)	-0.0051 (11)	0.0020 (10)	0.0003 (10)
C20	0.0145 (13)	0.0254 (15)	0.0146 (12)	-0.0024 (11)	0.0062 (10)	0.0024 (10)
C21	0.0127 (13)	0.0206 (14)	0.0152 (12)	-0.0025 (10)	0.0068 (10)	-0.0008 (10)
C22	0.0143 (13)	0.0293 (15)	0.0188 (12)	0.0008 (11)	0.0018 (10)	0.0078 (11)
C23	0.0207 (14)	0.0219 (15)	0.0238 (13)	0.0004 (11)	0.0073 (11)	0.0044 (11)
C24	0.0198 (13)	0.0187 (13)	0.0212 (12)	0.0003 (11)	0.0080 (10)	-0.0002 (10)

Geometric parameters (\AA , ^\circ)

Pb1—O4	2.3106 (18)	C7—C8	1.392 (3)
Pb1—O2	2.4270 (18)	C8—C9	1.393 (4)
Pb1—O3	2.4457 (17)	C8—H8	0.9500
Pb1—O1	2.4586 (19)	C9—C10	1.389 (4)
Pb1—N5	2.715 (2)	C9—H9	0.9500
Pb1—N6	2.763 (2)	C10—C11	1.387 (4)
O1—N1	1.308 (3)	C10—H10	0.9500
O2—N2	1.304 (3)	C11—C12	1.389 (4)
O3—N3	1.309 (3)	C11—H11	0.9500
O4—N4	1.305 (3)	C12—H12	0.9500
N1—N2	1.285 (3)	C13—C14	1.396 (4)
N1—C1	1.440 (3)	C13—H13	0.9500
N3—N4	1.284 (3)	C14—C15	1.367 (4)
N3—C7	1.439 (3)	C14—H14	0.9500
N5—C13	1.326 (3)	C15—C16	1.404 (4)
N5—C17	1.365 (3)	C15—H15	0.9500
N6—C24	1.325 (3)	C16—C17	1.422 (4)
N6—C21	1.354 (3)	C16—C18	1.428 (4)
C1—C6	1.386 (4)	C17—C21	1.443 (4)
C1—C2	1.392 (4)	C18—C19	1.354 (4)
C2—C3	1.385 (4)	C18—H18	0.9500
C2—H2	0.9500	C19—C20	1.433 (4)
C3—C4	1.392 (4)	C19—H19	0.9500
C3—H3	0.9500	C20—C22	1.401 (4)
C4—C5	1.387 (4)	C20—C21	1.421 (4)
C4—H4	0.9500	C22—C23	1.367 (4)
C5—C6	1.385 (4)	C22—H22	0.9500
C5—H5	0.9500	C23—C24	1.406 (4)
C6—H6	0.9500	C23—H23	0.9500
C7—C12	1.378 (3)	C24—H24	0.9500
O4—Pb1—O2		C8—C7—N3	
76.42 (7)		119.2 (2)	

O4—Pb1—O3	65.37 (6)	C7—C8—C9	118.5 (2)
O2—Pb1—O3	123.25 (6)	C7—C8—H8	120.8
O4—Pb1—O1	90.97 (7)	C9—C8—H8	120.8
O2—Pb1—O1	62.97 (6)	C10—C9—C8	120.1 (2)
O3—Pb1—O1	76.93 (6)	C10—C9—H9	119.9
O4—Pb1—N5	74.56 (6)	C8—C9—H9	119.9
O2—Pb1—N5	75.51 (6)	C11—C10—C9	120.0 (2)
O3—Pb1—N5	127.08 (6)	C11—C10—H10	120.0
O1—Pb1—N5	138.25 (6)	C9—C10—H10	120.0
O4—Pb1—N6	75.59 (6)	C10—C11—C12	120.7 (2)
O2—Pb1—N6	132.52 (7)	C10—C11—H11	119.7
O3—Pb1—N6	76.70 (6)	C12—C11—H11	119.7
O1—Pb1—N6	153.50 (7)	C7—C12—C11	118.5 (2)
N5—Pb1—N6	60.47 (6)	C7—C12—H12	120.8
N1—O1—Pb1	115.64 (14)	C11—C12—H12	120.8
N2—O2—Pb1	122.70 (14)	N5—C13—C14	123.8 (3)
N3—O3—Pb1	112.12 (13)	N5—C13—H13	118.1
N4—O4—Pb1	122.37 (14)	C14—C13—H13	118.1
N2—N1—O1	124.7 (2)	C15—C14—C13	118.9 (3)
N2—N1—C1	117.8 (2)	C15—C14—H14	120.5
O1—N1—C1	117.5 (2)	C13—C14—H14	120.5
N1—N2—O2	113.4 (2)	C14—C15—C16	119.7 (2)
N4—N3—O3	124.9 (2)	C14—C15—H15	120.2
N4—N3—C7	116.4 (2)	C16—C15—H15	120.2
O3—N3—C7	118.64 (19)	C15—C16—C17	117.8 (2)
N3—N4—O4	114.2 (2)	C15—C16—C18	122.4 (2)
C13—N5—C17	118.0 (2)	C17—C16—C18	119.8 (3)
C13—N5—Pb1	120.57 (17)	N5—C17—C16	121.8 (2)
C17—N5—Pb1	121.42 (16)	N5—C17—C21	119.0 (2)
C24—N6—C21	118.7 (2)	C16—C17—C21	119.2 (2)
C24—N6—Pb1	121.10 (17)	C19—C18—C16	121.1 (2)
C21—N6—Pb1	120.19 (16)	C19—C18—H18	119.4
C6—C1—C2	121.9 (2)	C16—C18—H18	119.4
C6—C1—N1	118.0 (2)	C18—C19—C20	121.0 (2)
C2—C1—N1	120.1 (2)	C18—C19—H19	119.5
C3—C2—C1	118.4 (3)	C20—C19—H19	119.5
C3—C2—H2	120.8	C22—C20—C21	117.7 (2)
C1—C2—H2	120.8	C22—C20—C19	122.6 (2)
C2—C3—C4	120.5 (3)	C21—C20—C19	119.7 (2)
C2—C3—H3	119.8	N6—C21—C20	121.9 (2)
C4—C3—H3	119.8	N6—C21—C17	118.9 (2)
C5—C4—C3	120.2 (3)	C20—C21—C17	119.3 (2)
C5—C4—H4	119.9	C23—C22—C20	119.9 (3)
C3—C4—H4	119.9	C23—C22—H22	120.0
C6—C5—C4	120.1 (3)	C20—C22—H22	120.0
C6—C5—H5	119.9	C22—C23—C24	118.8 (3)
C4—C5—H5	119.9	C22—C23—H23	120.6
C1—C6—C5	119.0 (3)	C24—C23—H23	120.6

C1—C6—H6	120.5	N6—C24—C23	123.1 (3)
C5—C6—H6	120.5	N6—C24—H24	118.5
C12—C7—C8	122.1 (2)	C23—C24—H24	118.5
C12—C7—N3	118.6 (2)		
O4—Pb1—O1—N1	68.29 (18)	C1—C2—C3—C4	1.3 (5)
O2—Pb1—O1—N1	-5.91 (16)	C2—C3—C4—C5	-1.4 (6)
O3—Pb1—O1—N1	132.71 (18)	C3—C4—C5—C6	1.0 (6)
N5—Pb1—O1—N1	0.7 (2)	C2—C1—C6—C5	0.3 (5)
N6—Pb1—O1—N1	126.70 (18)	N1—C1—C6—C5	178.4 (3)
O4—Pb1—O2—N2	-92.0 (2)	C4—C5—C6—C1	-0.4 (5)
O3—Pb1—O2—N2	-44.2 (2)	N4—N3—C7—C12	-152.2 (2)
O1—Pb1—O2—N2	6.17 (19)	O3—N3—C7—C12	23.7 (3)
N5—Pb1—O2—N2	-169.3 (2)	N4—N3—C7—C8	26.5 (3)
N6—Pb1—O2—N2	-147.37 (18)	O3—N3—C7—C8	-157.6 (2)
O4—Pb1—O3—N3	-7.02 (13)	C12—C7—C8—C9	3.9 (4)
O2—Pb1—O3—N3	-59.47 (16)	N3—C7—C8—C9	-174.7 (2)
O1—Pb1—O3—N3	-104.23 (15)	C7—C8—C9—C10	-1.7 (4)
N5—Pb1—O3—N3	37.44 (17)	C8—C9—C10—C11	-1.7 (4)
N6—Pb1—O3—N3	73.02 (14)	C9—C10—C11—C12	2.9 (4)
O2—Pb1—O4—N4	146.36 (18)	C8—C7—C12—C11	-2.7 (4)
O3—Pb1—O4—N4	9.37 (16)	N3—C7—C12—C11	176.0 (2)
O1—Pb1—O4—N4	84.50 (17)	C10—C11—C12—C7	-0.8 (4)
N5—Pb1—O4—N4	-135.20 (18)	C17—N5—C13—C14	0.0 (4)
N6—Pb1—O4—N4	-72.39 (17)	Pb1—N5—C13—C14	179.90 (19)
Pb1—O1—N1—N2	6.5 (3)	N5—C13—C14—C15	0.5 (4)
Pb1—O1—N1—C1	-174.66 (17)	C13—C14—C15—C16	-0.5 (4)
O1—N1—N2—O2	-1.0 (4)	C14—C15—C16—C17	0.2 (4)
C1—N1—N2—O2	-179.9 (2)	C14—C15—C16—C18	-178.3 (2)
Pb1—O2—N2—N1	-5.4 (3)	C13—N5—C17—C16	-0.4 (4)
Pb1—O3—N3—N4	5.5 (3)	Pb1—N5—C17—C16	179.72 (17)
Pb1—O3—N3—C7	-170.04 (15)	C13—N5—C17—C21	178.8 (2)
O3—N3—N4—O4	2.4 (3)	Pb1—N5—C17—C21	-1.1 (3)
C7—N3—N4—O4	177.98 (19)	C15—C16—C17—N5	0.3 (4)
Pb1—O4—N4—N3	-10.2 (3)	C18—C16—C17—N5	178.9 (2)
O4—Pb1—N5—C13	-96.9 (2)	C15—C16—C17—C21	-178.9 (2)
O2—Pb1—N5—C13	-17.25 (19)	C18—C16—C17—C21	-0.3 (4)
O3—Pb1—N5—C13	-138.21 (18)	C15—C16—C18—C19	178.8 (3)
O1—Pb1—N5—C13	-23.3 (2)	C17—C16—C18—C19	0.3 (4)
N6—Pb1—N5—C13	-178.8 (2)	C16—C18—C19—C20	0.3 (4)
O4—Pb1—N5—C17	83.03 (18)	C18—C19—C20—C22	-179.3 (3)
O2—Pb1—N5—C17	162.6 (2)	C18—C19—C20—C21	-0.9 (4)
O3—Pb1—N5—C17	41.7 (2)	C24—N6—C21—C20	0.6 (4)
O1—Pb1—N5—C17	156.59 (17)	Pb1—N6—C21—C20	-179.48 (18)
N6—Pb1—N5—C17	1.10 (17)	C24—N6—C21—C17	-178.8 (2)
O4—Pb1—N6—C24	98.62 (19)	Pb1—N6—C21—C17	1.0 (3)
O2—Pb1—N6—C24	154.25 (17)	C22—C20—C21—N6	-0.1 (4)
O3—Pb1—N6—C24	31.04 (18)	C19—C20—C21—N6	-178.6 (2)

O1—Pb1—N6—C24	37.1 (3)	C22—C20—C21—C17	179.4 (2)
N5—Pb1—N6—C24	178.8 (2)	C19—C20—C21—C17	0.9 (4)
O4—Pb1—N6—C21	-81.27 (18)	N5—C17—C21—N6	0.0 (3)
O2—Pb1—N6—C21	-25.6 (2)	C16—C17—C21—N6	179.2 (2)
O3—Pb1—N6—C21	-148.84 (19)	N5—C17—C21—C20	-179.5 (2)
O1—Pb1—N6—C21	-142.83 (17)	C16—C17—C21—C20	-0.3 (4)
N5—Pb1—N6—C21	-1.08 (17)	C21—C20—C22—C23	-0.4 (4)
N2—N1—C1—C6	158.2 (3)	C19—C20—C22—C23	178.1 (2)
O1—N1—C1—C6	-20.8 (4)	C20—C22—C23—C24	0.3 (4)
N2—N1—C1—C2	-23.7 (4)	C21—N6—C24—C23	-0.7 (4)
O1—N1—C1—C2	157.4 (3)	Pb1—N6—C24—C23	179.37 (18)
C6—C1—C2—C3	-0.7 (5)	C22—C23—C24—N6	0.3 (4)
N1—C1—C2—C3	-178.8 (3)		