

1-Dimethylamino-9,10-anthraquinone

Paweł Niedziałkowski, Joanna Narloch, Damian Trzybiński* and Tadeusz Ossowski

Faculty of Chemistry, University of Gdańsk, J. Sobieskiego 18, 80-952 Gdańsk, Poland

Correspondence e-mail: trzybinski@chem.univ.gda.pl


Received 14 February 2011; accepted 22 February 2011

Key indicators: single-crystal X-ray study; $T = 295$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.037; wR factor = 0.079; data-to-parameter ratio = 7.2.

In the crystal structure of the title compound, $\text{C}_{16}\text{H}_{13}\text{NO}_2$, adjacent molecules are linked through $\text{C}-\text{H}\cdots\pi$ and $\pi-\pi$ [centroid-centroid distances = $3.844(2)$ Å] contacts. The anthracene ring system and dimethylamino group are oriented at a dihedral angle of $38.4(1)^\circ$. In the crystal, the mean planes of adjacent anthracene units are inclined at angles of $59.3(1)$, $75.7(1)$ and $76.0(1)^\circ$.

Related literature

For general background to anthraquinones, see: Arai *et al.* (1985); Dalliya *et al.* (2007); Gatto *et al.* (1996); Kowalczyk *et al.* (2010); Mori *et al.* (1990); Ossowski *et al.* (2005); Zoń *et al.* (2003). For a related structure, see: Yatsenko *et al.* (2000). For molecular interactions, see: Hunter *et al.* (2001); Spek (2009); Takahashi *et al.* (2001).



Experimental

Crystal data

$\text{C}_{16}\text{H}_{13}\text{NO}_2$	$V = 1216.82(11)$ Å ³
$M_r = 251.27$	$Z = 4$
Orthorhombic, $P2_12_12_1$	Mo $K\alpha$ radiation
$a = 7.2823(3)$ Å	$\mu = 0.09$ mm ⁻¹
$b = 11.1519(7)$ Å	$T = 295$ K
$c = 14.9834(7)$ Å	$0.45 \times 0.20 \times 0.18$ mm

Data collection

Oxford Diffraction Gemini R ULTRA Ruby CCD diffractometer	1258 independent reflections
4683 measured reflections	918 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.033$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.037$	174 parameters
$wR(F^2) = 0.079$	H-atom parameters constrained
$S = 0.96$	$\Delta\rho_{\text{max}} = 0.12$ e Å ⁻³
1258 reflections	$\Delta\rho_{\text{min}} = -0.18$ e Å ⁻³

Table 1

Hydrogen-bond geometry (Å, °).

Cg_1 and Cg_2 are the centroids of the C1–C4/C11/C12 and C5–C8/C13/C14 rings respectively.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$C2-H2\cdots Cg_1^{i1}$	0.93	2.99	3.724 (3)	137
$C4-H4\cdots Cg_2^{ii}$	0.93	2.81	3.678 (3)	156

Symmetry codes: (i) $-x, y + \frac{3}{2}, -z + \frac{3}{2}$; (ii) $x + \frac{3}{2}, -y + \frac{1}{2}, -z + 1$.

Data collection: *CrysAlis CCD* (Oxford Diffraction, 2008); cell refinement: *CrysAlis RED* (Oxford Diffraction, 2008); data reduction: *CrysAlis RED*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97* and *PLATON* (Spek, 2009).

This study was financed by the State Funds for Scientific Research (grant DS/8210-4-0177-11).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NG5119).

References

- Arai, S., Kato, S. & Hida, M. (1985). *Bull. Chem. Soc. Jpn.* **58**, 1458–1463.
- Dalliya, P., Maity, D. K., Nayak, S. K., Mukherjee, T. & Pal, H. (2007). *Photochem. Photobiol. A Chem.* **186**, 218–228.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Gatto, B., Zagotto, G., Sissi, C., Cera, C., Uriarte, E., Palu, G., Capranico, G. & Palumbo, M. (1996). *J. Med. Chem.* **39**, 3114–3122.
- Hunter, C. A., Lawson, K. R., Perkins, J. & Urch, C. J. (2001). *J. Chem. Soc. Perkin Trans. 2*, pp. 651–669.
- Kowalczyk, A., Nowicka, A. M., Jurczakowski, R., Niedziałkowski, P., Ossowski, T. & Stojek, Z. (2010). *Electroanalysis*, **22**, 49–59.
- Mori, H., Yoshimi, N., Iwata, H., Mori, Y., Hara, A., Tanaka, T. & Kawai, K. (1990). *Carcinogenesis*, **11**, 799–802.
- Ossowski, T., Zarzeckańska, D., Zalewski, L., Niedziałkowski, P., Majewski, R. & Szymańska, A. (2005). *Tetrahedron Lett.* **46**, 1735–1738.
- Oxford Diffraction. (2008). *CrysAlis CCD* and *CrysAlis RED*. Oxford Diffraction Ltd, Abingdon, England.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Takahashi, O., Kohno, Y., Iwasaki, S., Saito, K., Iwaoka, M., Tomada, S., Umezawa, Y., Tsuboyama, S. & Nishio, M. (2001). *Bull. Chem. Soc. Jpn.* **74**, 2421–2430.
- Yatsenko, A. V., Paseshnichenko, K. A. & Popov, S. I. (2000). *Z. Kristallogr.* **215**, 542–546.
- Zoń, A., Palys, M., Stojek, Z., Sulowska, H. & Ossowski, T. (2003). *Electroanalysis*, **15**, 579–585.

supporting information

Acta Cryst. (2011). E67, o723 [doi:10.1107/S1600536811006829]

1-Dimethylamino-9,10-anthraquinone

Paweł Niedziałkowski, Joanna Narloch, Damian Trzybiński and Tadeusz Ossowski

S1. Comment

Anthraquinones, the largest group of naturally occurring quinones, present in bacteria, fungi and many higher plant families contain π -electrons, reducible *p*-quinone system and are redoxactive (Zoń *et al.*, 2003). That is the reason why those found many practical applications (Kowalczyk *et al.*, 2010; Ossowski *et al.*, 2005). Both natural and synthetic derivatives have been used as colourants in food, cosmetics, textiles and hair dyes (Mori *et al.*, 1990). In medicine they are known as antitumor drugs and antibacterial or anti-inflammatory agents (Gatto *et al.*, 1996). Among anthraquinones, the amino-derivatives, due to the possibility of their chemical modification, reveal greatest potential of application. Here, we present the crystal structure of the 1-(dimethylamino)-9,10-anthraquinone – compound with interesting photophysical properties (Arai *et al.*, 1985; Dalliya *et al.*, 2007).

In the molecule of the title compound (Fig. 1), likewise in the 1-(methyl(phenyl)amino)anthraquinone (Yatsenko *et al.*, 2000), relatively strong deviation of planarity of the anthraquinone skeleton is observed. In case of the title compound, such distortion (0.1274 (3) Å) is directly caused by the steric effect of the bulky $-\text{N}(\text{CH}_3)_2$ group (Dalliya *et al.*, 2007). The dimethylamino group is twisted at an angle of 38.4 (1)° relative to the anthracene fragment. The neighboring anthracene moieties are inclined at an angle of 59.3 (1)°, 75.7 (1)° and 76.0 (1)° in the crystal lattice.

In the crystal structure, the adjacent molecules are linked by C–H \cdots π (Table 2, Fig. 2) and π – π [centroid-centroid distances = 3.844 (2) Å] (Table 3, Fig. 3) contacts. All interactions demonstrated were found by *PLATON* (Spek, 2009). The C–H \cdots π interactions should be of an attractive nature (Takahashi *et al.*, 2001), like the π – π (Hunter *et al.*, 2001) interactions.

S2. Experimental

1-(Dimethylamino)-9,10-anthraquinone was synthesized according to the procedure described below. The solution of 40% dimethylamine in water (2,21 ml, 12.36 mmol) was added to 1-chloro-9,10-anthraquinone (1 g, 4.12 mmol) in 15 ml toluene. The mixture was stirred at 130° for 4 h. The progress of the reaction was monitored by TLC (SiO₂, dichloromethane) until the completion of reaction. The resulting mixture was concentrated to remove the solvent and dissolved in 100 ml of dichloromethane. The solution was washed with water (100 ml), the organic phase was dried over MgSO₄ and concentrated. The resultant solid was purified by column chromatography using dichloromethane as a solvent obtaining the title compound as a red solid (921 mg, 89%). The product was recrystallized by slow evaporation from methanol to give red crystals suitable for X-ray diffraction (m.p. 137.5–137.9°C). Spectral data: IR (KBr): 3584, 2916, 2806, 1662, 1637, 1551, 1499, 1374, 1311, 1270, 1180, 1024, 935, 793, 731, 704 cm⁻¹; ¹H NMR (CDCl₃, 400 MHz): 3.03 (6H, s, CH₃), 7.34–7.36 (1H, d, J₁ = 8.8 Hz, H-2-Ar), 7.54–7.58 (1H, t, J₁ = J₂ = 8.0 Hz, H-3-Ar), 7.69–7.72 (1H, t, J₁ = 7.6 Hz, J₂ = 6.8 Hz, J₃ = 7.2 Hz, H-6-Ar), 7.74–7.76 (1H, d, J₁ = 7.6 Hz, H-4-Ar), 7.78–7.82 (1H, t, J₁ = 7.6 Hz, J₂ = 8.4 Hz, J₃ = 8.0 Hz, H-7-Ar), 8.22–8.24 (1H, t, J₁ = 7.2 Hz, H-8-Ar). Elemental analysis: calculated for C₁₆H₁₃NO₂: C 76.48, H 5.21, N 5.57; found: C 76.52, H 5.28, N 5.51.

S3. Refinement

1063 Friedel pairs were merged. H atoms were positioned geometrically, with C—H = 0.93 Å and 0.96 Å for the aromatic and methyl H atoms, respectively, and constrained to ride on their parent atoms with $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C})$, where $x = 1.2$ for the aromatic and $x = 1.5$ for the methyl H atoms.





Figure 1

The molecular structure of the title compound showing the atom-labeling scheme. Displacement ellipsoids are drawn at the 25% probability level, and H atoms are shown as small spheres of arbitrary radius. Cg1 and Cg2 are the centroids of the C1—C4/C11/C12 and C5—C8/C13/C14 rings respectively.

**Figure 2**

The arrangement of the molecules in the crystal structure viewed approximately along *a* direction. The C–H... π interactions are represented by dotted lines. H atoms not involved in interactions have been omitted. [Symmetry codes: (i) $-x, y + 3/2, -z + 3/2$; (ii) $x + 3/2, -y + 1/2, -z + 1$.]




Figure 3

The arrangement of the molecules in the crystal structure viewed approximately along b direction. The π - π interactions are represented by dotted lines. H atoms have been omitted for clarity. [Symmetry codes: (iii) $x + 1, y, z$; (iv) $x - 1, y + 1, z + 1$.]

1-Dimethylamino-9,10-anthraquinone

Crystal data

$C_{16}H_{13}NO_2$

$M_r = 251.27$

Orthorhombic, $P2_12_12_1$

Hall symbol: P 2ac 2ab

$a = 7.2823$ (3) Å

$b = 11.1519$ (7) Å

$c = 14.9834$ (7) Å

$V = 1216.82$ (11) Å³

$Z = 4$

$F(000) = 528$

$D_x = 1.372$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 1944 reflections

$\theta = 3.1$ – 29.0°

$\mu = 0.09 \text{ mm}^{-1}$
 $T = 295 \text{ K}$

Prism, red
 $0.45 \times 0.20 \times 0.18 \text{ mm}$

Data collection

Oxford Diffraction Gemini R ULTRA Ruby
 CCD
 diffractometer
 Radiation source: Enhance (Mo) X-ray Source
 Graphite monochromator
 Detector resolution: $10.4002 \text{ pixels mm}^{-1}$
 ω scans
 4683 measured reflections

1258 independent reflections
 918 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.033$
 $\theta_{\text{max}} = 25.1^\circ$, $\theta_{\text{min}} = 3.1^\circ$
 $h = -8 \rightarrow 6$
 $k = -12 \rightarrow 13$
 $l = -17 \rightarrow 13$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.037$
 $wR(F^2) = 0.079$
 $S = 0.96$
 1258 reflections
 174 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0492P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} < 0.001$
 $\Delta\rho_{\text{max}} = 0.12 \text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.18 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.7457 (3)	0.5675 (2)	0.53415 (14)	0.0349 (6)
C2	0.8768 (3)	0.6609 (3)	0.53396 (17)	0.0437 (7)
H2	0.9738	0.6578	0.4937	0.052*
C3	0.8647 (4)	0.7554 (3)	0.59116 (18)	0.0486 (7)
H3	0.9530	0.8155	0.5892	0.058*
C4	0.7235 (3)	0.7630 (2)	0.65179 (17)	0.0422 (6)
H4	0.7206	0.8255	0.6928	0.051*
C5	0.1522 (4)	0.6067 (3)	0.78267 (16)	0.0459 (7)
H5	0.1608	0.6622	0.8288	0.055*
C6	0.0081 (4)	0.5277 (3)	0.78088 (18)	0.0487 (7)
H6	-0.0795	0.5290	0.8260	0.058*
C7	-0.0067 (3)	0.4466 (3)	0.71212 (17)	0.0465 (7)
H7	-0.1065	0.3946	0.7099	0.056*
C8	0.1262 (3)	0.4422 (2)	0.64640 (17)	0.0409 (6)
H8	0.1164	0.3863	0.6006	0.049*
C9	0.4157 (3)	0.5148 (2)	0.57716 (15)	0.0334 (6)
C10	0.4341 (4)	0.6937 (2)	0.71603 (16)	0.0394 (7)
C11	0.5878 (3)	0.5814 (2)	0.58994 (14)	0.0325 (6)

C12	0.5869 (3)	0.6781 (2)	0.65160 (16)	0.0335 (6)
C13	0.2744 (3)	0.5206 (2)	0.64829 (15)	0.0335 (6)
C14	0.2856 (3)	0.6048 (2)	0.71661 (14)	0.0354 (6)
N15	0.7757 (3)	0.4671 (2)	0.48311 (13)	0.0421 (6)
C16	0.7173 (4)	0.3491 (2)	0.51109 (19)	0.0528 (8)
H16A	0.6745	0.3525	0.5716	0.079*
H16B	0.8189	0.2945	0.5072	0.079*
H16C	0.6198	0.3219	0.4730	0.079*
C17	0.9195 (3)	0.4667 (3)	0.41526 (17)	0.0552 (8)
H17A	0.9068	0.5362	0.3780	0.083*
H17B	0.9086	0.3957	0.3794	0.083*
H17C	1.0376	0.4679	0.4437	0.083*
O18	0.3823 (2)	0.45881 (18)	0.50827 (11)	0.0484 (5)
O19	0.4317 (3)	0.77920 (19)	0.76763 (15)	0.0644 (6)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0328 (13)	0.0405 (16)	0.0315 (12)	0.0014 (14)	-0.0002 (12)	0.0029 (12)
C2	0.0356 (14)	0.0555 (19)	0.0400 (14)	-0.0071 (15)	0.0045 (12)	0.0082 (14)
C3	0.0453 (14)	0.0471 (17)	0.0532 (16)	-0.0149 (15)	-0.0037 (15)	0.0040 (17)
C4	0.0427 (14)	0.0392 (16)	0.0448 (14)	-0.0052 (14)	-0.0053 (13)	-0.0071 (14)
C5	0.0478 (15)	0.0464 (18)	0.0435 (14)	0.0079 (15)	0.0064 (13)	-0.0049 (14)
C6	0.0377 (14)	0.057 (2)	0.0512 (15)	0.0046 (16)	0.0143 (12)	0.0078 (17)
C7	0.0345 (14)	0.0513 (19)	0.0538 (16)	-0.0061 (14)	-0.0005 (13)	0.0095 (17)
C8	0.0348 (13)	0.0461 (16)	0.0416 (13)	-0.0012 (13)	-0.0033 (12)	0.0038 (14)
C9	0.0339 (12)	0.0348 (15)	0.0314 (12)	0.0014 (12)	-0.0027 (11)	0.0011 (12)
C10	0.0434 (15)	0.0391 (16)	0.0358 (14)	0.0024 (13)	-0.0017 (13)	-0.0039 (13)
C11	0.0296 (12)	0.0372 (15)	0.0306 (12)	0.0010 (12)	-0.0026 (11)	0.0048 (12)
C12	0.0317 (12)	0.0346 (14)	0.0342 (12)	0.0011 (13)	-0.0051 (12)	0.0036 (12)
C13	0.0284 (12)	0.0364 (15)	0.0358 (12)	0.0050 (12)	-0.0062 (11)	0.0035 (12)
C14	0.0319 (13)	0.0378 (15)	0.0364 (12)	0.0035 (12)	-0.0001 (12)	0.0014 (13)
N15	0.0347 (11)	0.0497 (14)	0.0418 (11)	-0.0001 (12)	0.0084 (10)	-0.0042 (12)
C16	0.0510 (16)	0.0460 (18)	0.0614 (17)	0.0072 (16)	0.0043 (15)	-0.0056 (16)
C17	0.0407 (14)	0.076 (2)	0.0489 (15)	0.0057 (17)	0.0081 (13)	-0.0116 (16)
O18	0.0404 (10)	0.0629 (12)	0.0418 (10)	-0.0084 (10)	0.0001 (8)	-0.0145 (10)
O19	0.0661 (13)	0.0604 (14)	0.0665 (13)	-0.0108 (11)	0.0160 (11)	-0.0280 (13)

Geometric parameters (Å, °)

C1—N15	1.373 (3)	C8—H8	0.9300
C1—C2	1.413 (4)	C9—O18	1.230 (3)
C1—C11	1.430 (3)	C9—C11	1.469 (3)
C2—C3	1.360 (4)	C9—C13	1.483 (3)
C2—H2	0.9300	C10—O19	1.227 (3)
C3—C4	1.374 (4)	C10—C14	1.467 (4)
C3—H3	0.9300	C10—C12	1.483 (3)
C4—C12	1.374 (3)	C11—C12	1.420 (3)

C4—H4	0.9300	C13—C14	1.392 (3)
C5—C6	1.370 (4)	N15—C16	1.446 (3)
C5—C14	1.387 (3)	N15—C17	1.459 (3)
C5—H5	0.9300	C16—H16A	0.9600
C6—C7	1.375 (4)	C16—H16B	0.9600
C6—H6	0.9300	C16—H16C	0.9600
C7—C8	1.382 (3)	C17—H17A	0.9600
C7—H7	0.9300	C17—H17B	0.9600
C8—C13	1.389 (3)	C17—H17C	0.9600
N15—C1—C2	119.5 (2)	C14—C10—C12	118.5 (2)
N15—C1—C11	122.8 (2)	C12—C11—C1	117.8 (2)
C2—C1—C11	117.7 (2)	C12—C11—C9	117.7 (2)
C3—C2—C1	121.7 (2)	C1—C11—C9	123.7 (2)
C3—C2—H2	119.1	C4—C12—C11	121.4 (2)
C1—C2—H2	119.1	C4—C12—C10	117.5 (2)
C2—C3—C4	120.8 (3)	C11—C12—C10	121.1 (2)
C2—C3—H3	119.6	C8—C13—C14	119.0 (2)
C4—C3—H3	119.6	C8—C13—C9	119.8 (2)
C12—C4—C3	119.9 (2)	C14—C13—C9	121.1 (2)
C12—C4—H4	120.1	C5—C14—C13	119.6 (2)
C3—C4—H4	120.1	C5—C14—C10	120.7 (2)
C6—C5—C14	120.9 (2)	C13—C14—C10	119.7 (2)
C6—C5—H5	119.6	C1—N15—C16	122.26 (19)
C14—C5—H5	119.6	C1—N15—C17	120.3 (2)
C5—C6—C7	119.8 (2)	C16—N15—C17	114.2 (2)
C5—C6—H6	120.1	N15—C16—H16A	109.5
C7—C6—H6	120.1	N15—C16—H16B	109.5
C6—C7—C8	120.2 (2)	H16A—C16—H16B	109.5
C6—C7—H7	119.9	N15—C16—H16C	109.5
C8—C7—H7	119.9	H16A—C16—H16C	109.5
C7—C8—C13	120.5 (2)	H16B—C16—H16C	109.5
C7—C8—H8	119.8	N15—C17—H17A	109.5
C13—C8—H8	119.8	N15—C17—H17B	109.5
O18—C9—C11	122.3 (2)	H17A—C17—H17B	109.5
O18—C9—C13	119.2 (2)	N15—C17—H17C	109.5
C11—C9—C13	118.4 (2)	H17A—C17—H17C	109.5
O19—C10—C14	120.7 (2)	H17B—C17—H17C	109.5
O19—C10—C12	120.8 (2)		
N15—C1—C2—C3	-172.0 (2)	O19—C10—C12—C11	-178.1 (2)
C11—C1—C2—C3	6.6 (3)	C14—C10—C12—C11	1.8 (3)
C1—C2—C3—C4	0.2 (4)	C7—C8—C13—C14	-0.9 (3)
C2—C3—C4—C12	-3.7 (4)	C7—C8—C13—C9	-179.9 (2)
C14—C5—C6—C7	-1.0 (4)	O18—C9—C13—C8	14.8 (3)
C5—C6—C7—C8	1.9 (4)	C11—C9—C13—C8	-168.1 (2)
C6—C7—C8—C13	-1.0 (4)	O18—C9—C13—C14	-164.1 (2)
N15—C1—C11—C12	168.8 (2)	C11—C9—C13—C14	12.9 (3)

C2—C1—C11—C12	-9.8 (3)	C6—C5—C14—C13	-0.9 (4)
N15—C1—C11—C9	-21.7 (3)	C6—C5—C14—C10	176.6 (2)
C2—C1—C11—C9	159.7 (2)	C8—C13—C14—C5	1.9 (3)
O18—C9—C11—C12	155.6 (2)	C9—C13—C14—C5	-179.2 (2)
C13—C9—C11—C12	-21.3 (3)	C8—C13—C14—C10	-175.7 (2)
O18—C9—C11—C1	-13.8 (4)	C9—C13—C14—C10	3.3 (3)
C13—C9—C11—C1	169.2 (2)	O19—C10—C14—C5	-8.3 (4)
C3—C4—C12—C11	0.2 (4)	C12—C10—C14—C5	171.8 (2)
C3—C4—C12—C10	-177.5 (2)	O19—C10—C14—C13	169.2 (2)
C1—C11—C12—C4	6.6 (3)	C12—C10—C14—C13	-10.7 (3)
C9—C11—C12—C4	-163.5 (2)	C2—C1—N15—C16	145.7 (2)
C1—C11—C12—C10	-175.8 (2)	C11—C1—N15—C16	-32.9 (3)
C9—C11—C12—C10	14.1 (3)	C2—C1—N15—C17	-12.9 (3)
O19—C10—C12—C4	-0.4 (4)	C11—C1—N15—C17	168.5 (2)
C14—C10—C12—C4	179.5 (2)		

Hydrogen-bond geometry (Å, °)

Cg1 and *Cg2* are the centroids of the C1—C4/C11/C12 and C5—C8/C13/C14 rings respectively.

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C2—H2... <i>Cg1</i> ⁱ	0.93	2.99	3.724 (3)	137
C4—H4... <i>Cg2</i> ⁱⁱ	0.93	2.81	3.678 (3)	156

Symmetry codes: (i) $-x, y+3/2, -z+3/2$; (ii) $x+3/2, -y+1/2, -z+1$.