

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

4-Nitrophenyl 2-bromo-2-methylpropanoate

Corrado Rizzoli,^{a*} Elda Marku^b and Lucedio Greci^c

^aDipartimento di Chimica Generale ed Inorganica, Chimica Analitica, Chimica Fisica, Università degli Studi di Parma, Viale G. P. Usberti 17/A, I-43124 Parma, Italy, ^bFakulteti i Shkencave të Natyrës, Departamenti i Kimisë, Universiteti i Tiranës, Bulevardi "Zogu I", Tirana, Albania, and ^cDipartimento ISAC, Università Politecnica delle Marche, Via Breccie Bianche, I-60131 Ancona, Italy
Correspondence e-mail: corrado.rizzoli@unipr.it


Received 16 February 2011; accepted 17 February 2011

Key indicators: single-crystal X-ray study; $T = 295$ K; mean $\sigma(\text{C}-\text{C}) = 0.006$ Å; R factor = 0.046; wR factor = 0.132; data-to-parameter ratio = 15.0.

In the title compound, $\text{C}_{10}\text{H}_{10}\text{BrNO}_4$, the planes of the carboxylate and nitro groups are rotated by 60.53 (13) and 6.4 (3)°, respectively, to the benzene ring. In the crystal, intermolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds link the molecules into zigzag chains parallel to the c axis.

Related literature

For the synthesis and biological properties of the title compound and analogues, see: Bischoff (1907); Kaeriyama *et al.* (1976). For the use of the title compound in organic synthesis, see: Haddleton & Waterson (1999); Edeleva *et al.* (2009); Guillaneuf *et al.* (2007).



Experimental

Crystal data

 $\text{C}_{10}\text{H}_{10}\text{BrNO}_4$ $M_r = 288.10$ Orthorhombic, *Pbcn* $a = 11.4128$ (16) Å $b = 14.450$ (2) Å $c = 14.539$ (2) Å $V = 2397.7$ (6) Å³ $Z = 8$ Mo $K\alpha$ radiation $\mu = 3.43$ mm⁻¹ $T = 295$ K $0.45 \times 0.15 \times 0.14$ mm

Data collection

Bruker APEXII CCD diffractometer
Absorption correction: multi-scan (SADABS; Bruker, 2008)
 $T_{\min} = 0.625$, $T_{\max} = 0.720$

18926 measured reflections
2177 independent reflections
1086 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.049$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.046$
 $wR(F^2) = 0.132$
 $S = 1.00$
2177 reflections

145 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.84$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.51$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C6}-\text{H6}\cdots\text{O3}^i$	0.93	2.59	3.385 (4)	144

Symmetry code: (i) $x, -y + 1, z - \frac{1}{2}$.

Data collection: APEX2 (Bruker, 2008); cell refinement: SAINT (Bruker, 2008); data reduction: SAINT; program(s) used to solve structure: SIR97 (Altomare *et al.*, 1999); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 for Windows (Farrugia, 1997) and SCHAKAL97 (Keller, 1997); software used to prepare material for publication: SHELXL97 and PARST95 (Nardelli, 1995).

Financial support from the Università Politecnica delle Marche and the Università degli Studi di Parma is gratefully acknowledged.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK2722).

References

- Altomare, A., Burla, M. C., Camalli, M., Casciarano, G. L., Giacovazzo, C., Guagliardi, A., Moliterni, A. G. G., Polidori, G. & Spagna, R. (1999). *J. Appl. Cryst.* **32**, 115–119.
- Bischoff, C. A. (1907). *Chem. Ber.* **39**, 3854–3861.
- Bruker (2008). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Edeleva, M. V., Kirilyuk, I. A., Zubenko, D. P., Zhurko, I. F., Marque, S. R. A., Gignes, D., Guillaneuf, Y. & Bagryanskaya, E. G. (2009). *J. Polym. Sci. Part A Polym. Chem.* **47**, 6579–6595.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Guillaneuf, Y., Gignes, D., Marque, S. R. A., Astolfi, P., Greci, L., Tordo, P. & Bertin, D. (2007). *Macromolecules*, **40**, 3108–3114.
- Haddleton, D. M. & Waterson, C. (1999). *Macromolecules*, **32**, 8732–8739.
- Kaeriyama, M., Sato, M., Kumita, I., Okuma, K., Ando, M., Hashimoto, S. & Yoneda, H. (1976). Jpn Kokai Tokyo Koho Patent JP 51146427.
- Keller, E. (1997). SCHAKAL97. University of Freiburg, Germany.
- Nardelli, M. (1995). *J. Appl. Cryst.* **28**, 659.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supporting information

Acta Cryst. (2011). E67, o698 [doi:10.1107/S1600536811005988]

4-Nitrophenyl 2-bromo-2-methylpropanoate

Corrado Rizzoli, Elda Marku and Lucedio Greci

S1. Comment

The synthesis of the title compound and analogues was originally reported in the early part of last century (Bischoff, 1907), and in the seventies several of these compounds were found to possess useful miticidal, insecticidal, nematocidal or fungicidal activities (Kaeriyama *et al.*, 1976). More recently, the title compound was prepared (Haddleton & Waterson, 1999) and used as initiator for H-atom transfer polymerization. The same compound was also used in the preparation of new alkoxyamines derived from imadaziline-, imidazoline- and pyrrolidine-1-oxyl nitroxides (Edeleva *et al.*, 2009) and, within our group, for the synthesis of phenyl- and 4-nitrophenyl- 2-(2,2-diphenyl-3-(phenylimino)-indolin-1-yloxy)-2-methylpropionate (Guillaneuf *et al.*, 2007). In order to obtain structural parameters for molecular mechanics calculations for the above mentioned alkoxyamines, the X-ray crystal structure of the title compound has been determined and the results are reported herein.

In the molecule of the title compound (Fig. 1), the plane of the nitro group is approximately coplanar with the benzene ring (dihedral angle 6.4 (3) °), whereas the plane of the carboxylic group is tilted by 60.53 (13) °. All bond lengths and angles are unexceptional. In the crystal structure (Fig. 2), the molecules are linked by intermolecular C—H···O hydrogen bonds (Table 1) into zigzag chains running parallel to the *c* axis.

S2. Experimental

The title compound was prepared according to the literature method (Haddleton & Waterson, 1999). Crystals suitable for X-ray analysis were obtained by slow evaporation of its *n*-pentane solution (m. p. 342–343 K). IR data, ν , cm⁻¹: 1753 (C=O), 1615 and 1592 (benzene ring), 1521 (NO₂). ¹H NMR spectrum, δ in CDCl₃: 2.08 (s, 6H); 7.3 (2H, d, *J* = 9.2 Hz); 8.31 (2H, d, *J* = 9.2 Hz). The ESI-MS obtained with 3200 QTRAP spectrometer does not give the molecular ion peak. The melting point was measured by an electrothermal apparatus and is uncorrected. The ¹H NMR spectrum was recorded with a Varian 400 MHz spectrometer. The IR spectrum was recorded with a Perkin-Elmer MGX1 spectrophotometer.

S3. Refinement

All H atoms were placed in geometrically idealized positions and treated as riding atoms, with C—H = 0.93–0.96 Å, and $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$ or $1.5 U_{\text{eq}}(\text{C})$ for methyl H atoms.




Figure 1

The molecular structure of the title compound, with displacement ellipsoids drawn at the 40% probability level.




Figure 2

Partial crystal packing of the title compound viewed approximately along the *a* axis. Intermolecular C—H...O hydrogen bonds are shown as dashed lines.

4-Nitrophenyl 2-bromo-2-methylpropanoate

Crystal data

$C_{10}H_{10}BrNO_4$

$M_r = 288.10$

Orthorhombic, *Pbcn*

Hall symbol: $-P\ 2n\ 2ab$

$a = 11.4128(16)\ \text{\AA}$

$b = 14.450(2)\ \text{\AA}$

$c = 14.539(2)\ \text{\AA}$

$V = 2397.7(6)\ \text{\AA}^3$

$Z = 8$

$F(000) = 1152$

$D_x = 1.596\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 877 reflections

$\theta = 5.6\text{--}21.3^\circ$

$\mu = 3.43 \text{ mm}^{-1}$
 $T = 295 \text{ K}$

Needle, colourless
 $0.45 \times 0.15 \times 0.14 \text{ mm}$

Data collection

Bruker APEXII CCD
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 ω scans
 Absorption correction: multi-scan
 (SADABS; Bruker, 2008)
 $T_{\min} = 0.625$, $T_{\max} = 0.720$

18926 measured reflections
 2177 independent reflections
 1086 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.049$
 $\theta_{\max} = 25.3^\circ$, $\theta_{\min} = 2.3^\circ$
 $h = -13 \rightarrow 13$
 $k = -15 \rightarrow 17$
 $l = -17 \rightarrow 17$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.046$
 $wR(F^2) = 0.132$
 $S = 1.00$
 2177 reflections
 145 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0685P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.84 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.51 \text{ e } \text{\AA}^{-3}$

Special details

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	-0.00330 (6)	0.10818 (4)	0.62160 (4)	0.1233 (3)
O1	-0.0531 (2)	0.33291 (18)	0.53800 (18)	0.0757 (8)
O2	0.1275 (2)	0.30016 (16)	0.58917 (16)	0.0639 (7)
O3	0.1600 (3)	0.6175 (2)	0.8793 (2)	0.1042 (11)
O4	0.1583 (4)	0.6983 (2)	0.7567 (2)	0.1188 (13)
N1	0.1583 (3)	0.6243 (3)	0.7961 (3)	0.0724 (9)
C1	0.0266 (3)	0.1820 (2)	0.5074 (2)	0.0603 (10)
C2	0.1417 (3)	0.1512 (2)	0.4701 (3)	0.0783 (12)
H2A	0.2016	0.1614	0.5154	0.117*
H2B	0.1597	0.1859	0.4156	0.117*
H2C	0.1380	0.0865	0.4554	0.117*
C3	-0.0733 (4)	0.1656 (3)	0.4429 (4)	0.1049 (17)
H3A	-0.1451	0.1849	0.4714	0.157*
H3B	-0.0777	0.1009	0.4283	0.157*
H3C	-0.0612	0.2005	0.3875	0.157*
C4	0.0267 (3)	0.2799 (3)	0.5452 (2)	0.0603 (10)
C5	0.1339 (3)	0.3821 (2)	0.6389 (3)	0.0551 (9)

C6	0.1295 (3)	0.4663 (3)	0.5954 (2)	0.0628 (10)
H6	0.1207	0.4695	0.5319	0.075*
C8	0.1531 (3)	0.5392 (3)	0.7411 (2)	0.0556 (9)
C7	0.1381 (3)	0.5461 (3)	0.6467 (3)	0.0626 (10)
H7	0.1339	0.6038	0.6184	0.075*
C9	0.1598 (3)	0.4549 (3)	0.7842 (2)	0.0618 (10)
H9	0.1699	0.4515	0.8476	0.074*
C10	0.1515 (3)	0.3756 (3)	0.7324 (3)	0.0623 (10)
H10	0.1576	0.3179	0.7604	0.075*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.1760 (7)	0.0961 (5)	0.0979 (5)	-0.0278 (3)	0.0319 (3)	0.0105 (3)
O1	0.0642 (16)	0.0717 (18)	0.091 (2)	0.0147 (14)	-0.0123 (15)	-0.0147 (15)
O2	0.0549 (15)	0.0642 (17)	0.0727 (16)	0.0074 (13)	-0.0079 (13)	-0.0145 (14)
O3	0.153 (3)	0.095 (2)	0.064 (2)	-0.023 (2)	-0.0036 (18)	-0.0151 (16)
O4	0.198 (4)	0.061 (2)	0.097 (2)	-0.017 (2)	-0.004 (2)	-0.0031 (19)
N1	0.084 (2)	0.071 (3)	0.062 (2)	-0.0081 (19)	0.0088 (19)	-0.009 (2)
C1	0.058 (2)	0.062 (2)	0.061 (2)	0.0014 (18)	-0.0044 (18)	-0.0028 (19)
C2	0.080 (3)	0.062 (3)	0.093 (3)	0.014 (2)	0.012 (2)	-0.005 (2)
C3	0.097 (3)	0.087 (3)	0.131 (4)	0.018 (3)	-0.043 (3)	-0.047 (3)
C4	0.053 (2)	0.070 (3)	0.058 (2)	0.006 (2)	-0.0084 (19)	-0.007 (2)
C5	0.0414 (19)	0.060 (2)	0.064 (2)	0.0041 (17)	0.0010 (17)	-0.006 (2)
C6	0.069 (2)	0.075 (3)	0.045 (2)	-0.001 (2)	0.0076 (18)	-0.001 (2)
C8	0.046 (2)	0.064 (3)	0.056 (2)	-0.0086 (18)	0.0046 (16)	-0.007 (2)
C7	0.071 (2)	0.059 (3)	0.057 (2)	-0.003 (2)	0.0135 (19)	0.003 (2)
C9	0.062 (2)	0.073 (3)	0.050 (2)	-0.002 (2)	-0.0102 (18)	0.005 (2)
C10	0.058 (2)	0.055 (2)	0.074 (3)	0.0015 (19)	-0.0109 (19)	0.0033 (19)

Geometric parameters (Å, °)

Br1—C1	2.002 (4)	C3—H3A	0.9600
O1—C4	1.195 (4)	C3—H3B	0.9600
O2—C4	1.348 (4)	C3—H3C	0.9600
O2—C5	1.390 (4)	C5—C6	1.372 (5)
O3—N1	1.214 (4)	C5—C10	1.378 (5)
O4—N1	1.214 (4)	C6—C7	1.376 (5)
N1—C8	1.467 (5)	C6—H6	0.9300
C1—C2	1.489 (5)	C8—C9	1.372 (5)
C1—C3	1.496 (5)	C8—C7	1.388 (5)
C1—C4	1.518 (5)	C7—H7	0.9300
C2—H2A	0.9600	C9—C10	1.375 (5)
C2—H2B	0.9600	C9—H9	0.9300
C2—H2C	0.9600	C10—H10	0.9300
C4—O2—C5	118.5 (3)	O1—C4—O2	123.6 (4)
O4—N1—O3	122.8 (4)	O1—C4—C1	124.4 (3)

O4—N1—C8	118.8 (3)	O2—C4—C1	112.0 (3)
O3—N1—C8	118.4 (4)	C6—C5—C10	121.4 (4)
C2—C1—C3	113.4 (3)	C6—C5—O2	121.0 (3)
C2—C1—C4	114.2 (3)	C10—C5—O2	117.6 (3)
C3—C1—C4	112.0 (3)	C5—C6—C7	119.4 (3)
C2—C1—Br1	107.1 (2)	C5—C6—H6	120.3
C3—C1—Br1	107.8 (3)	C7—C6—H6	120.3
C4—C1—Br1	101.3 (2)	C9—C8—C7	121.5 (3)
C1—C2—H2A	109.5	C9—C8—N1	119.5 (4)
C1—C2—H2B	109.5	C7—C8—N1	118.9 (3)
H2A—C2—H2B	109.5	C6—C7—C8	119.0 (4)
C1—C2—H2C	109.5	C6—C7—H7	120.5
H2A—C2—H2C	109.5	C8—C7—H7	120.5
H2B—C2—H2C	109.5	C8—C9—C10	119.1 (4)
C1—C3—H3A	109.5	C8—C9—H9	120.4
C1—C3—H3B	109.5	C10—C9—H9	120.4
H3A—C3—H3B	109.5	C9—C10—C5	119.5 (4)
C1—C3—H3C	109.5	C9—C10—H10	120.2
H3A—C3—H3C	109.5	C5—C10—H10	120.2
H3B—C3—H3C	109.5		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C6—H6...O3 ⁱ	0.93	2.59	3.385 (4)	144

Symmetry code: (i) *x*, $-\gamma+1$, $z-1/2$.