

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

1,4-Bis[(2-pyridylethyl)iminomethyl]-benzene

 Haleden Chiririwa,^a John R. Moss,^a Hong Su,^{a*} Denver Hendricks^b and Reinout Meijboom^c
^aDepartment of Chemistry, University of Cape Town, Private Bag, Rondebosch 7707, South Africa, ^bDivision of Medical Biochemistry, Faculty of Health Sciences, Private Bag X3, Observatory 7935, South Africa, and ^cDepartment of Chemistry University of Johannesburg, PO Box 524, Auckland Park, Johannesburg 2006, South Africa.


Correspondence e-mail: harrychiririwa@yahoo.com

Received 23 February 2011; accepted 15 March 2011

 Key indicators: single-crystal X-ray study; $T = 173$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.040; wR factor = 0.110; data-to-parameter ratio = 19.4.

In the title compound, $\text{C}_{22}\text{H}_{22}\text{N}_4$, the centroid of the benzene ring is located on an inversion centre. The dihedral angle between the benzene and pyridine rings is $10.94(5)^\circ$. The crystal structure displays weak intermolecular $\text{C}-\text{H}\cdots\text{N}$ hydrogen bonding and $\text{C}-\text{H}\cdots\pi$ interactions.

Related literature

 For related compounds, see: Chakraborty *et al.* (1999); Haga *et al.* (1985).


Experimental

Crystal data

 $\text{C}_{22}\text{H}_{22}\text{N}_4$
 $M_r = 342.44$
 Monoclinic, $P2_1/n$
 $a = 6.0078(6)$ Å
 $b = 26.023(3)$ Å
 $c = 6.1319(7)$ Å

 $\beta = 106.009(2)^\circ$
 $V = 921.47(17)$ Å³
 $Z = 2$
 Mo $K\alpha$ radiation

 $\mu = 0.08$ mm⁻¹
 $T = 173$ K
 $0.26 \times 0.24 \times 0.17$ mm

Data collection

 Bruker Kappa DUO APEXII
 diffractometer
 11941 measured reflections

 2288 independent reflections
 1945 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.024$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.040$
 $wR(F^2) = 0.110$
 $S = 1.06$
 2288 reflections

 118 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.28$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.20$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

 Cg1 and Cg2 are the centroids of the $\text{C1}-\text{C5}/\text{N1}$ and $\text{C9}-\text{C11}/\text{C9}'-\text{C11}'$ rings, respectively.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C3}-\text{H3}\cdots\text{N1}^{\text{i}}$	0.95	2.74	3.544 (3)	143 (3)
$\text{C4}-\text{H4}\cdots\text{N2}^{\text{ii}}$	0.95	2.69	3.593 (2)	159 (4)
$\text{C7}-\text{H7A}\cdots\text{N1}^{\text{iii}}$	0.99	2.87	3.847 (2)	171 (5)
$\text{C2}-\text{H2}\cdots\text{Cg1}^{\text{iii}}$	0.95	2.88	3.826 (4)	172 (5)
$\text{C6}-\text{H6A}\cdots\text{Cg2}^{\text{iv}}$	0.99	2.90	3.508 (3)	120 (2)

 Symmetry codes: (i) $x, y, z + 1$; (ii) $x - 1, y, z$; (iii) $x + \frac{1}{2}, -y + \frac{1}{2}, z + \frac{1}{2}$; (iv) $x + 1, y, z + 1$.

Data collection: *APEX2* (Bruker, 2006); cell refinement: *SAINTE* (Bruker, 2006); data reduction: *SAINTE*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *pubCIF* (Westrip, 2010).

We gratefully acknowledge Mintek and Project AuTEK for funding this project.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: GO2006).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
 Bruker (2006). *APEX2* and *SAINTE*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Chakraborty, S., Munshi, P. & Lahiri, G. K. (1999). *Polyhedron*, **18**, 1437–1444.
 Haga, M. & Koizumi, K. (1985). *Inorg. Chim. Acta*, **104**, 47–50.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2011). E67, o921 [doi:10.1107/S1600536811009809]

1,4-Bis[(2-pyridylethyl)iminomethyl]benzene

Haleden Chiririwa, John R. Moss, Hong Su, Denver Hendricks and Reinout Meijboom

S1. Comment


This work originates from our interest in developing a new class of tetradentate ligands. To the best of our knowledge, this work demonstrates the first example of neutral pyridinyldimine-based bridging ligand. The title compound might be expected to behave as a tetradentate chelating agent, in which both of the N atoms from the imine might coordinate, along with the two pyridinyl N atoms. Chakraborty *et al.* (1999) reported coordination of similar ligands to ruthenium whilst Haga and Koizumi (1985) reported their coordination to molybdenum. The structure of the title compound crystallized in space group $P2_1/n$ with $Z = 2$. The molecule, shown in Fig. 1, has a center of inversion at the centroid of the benzene ring and was located in special positions at Wyckoff position a . The conformation of the molecule is best described by the dihedral angle of the central ring and pyridyl ring of $10.94(5)^\circ$. The structure is stabilized by weak hydrogen bonds of the type $C-H\cdots N$ and $C-H\cdots\pi$, the metrics of which are given in Table 1. The $C-H\cdots N$ intermolecular interactions, as well as $C6-H6A\cdots$ Ring 1 (of $C10-C9-C11-C10'-C9'-C11'$), connect the parallel neighbouring molecules into 2-dimensional layers. And these layers are then linked along the b axis into 3-dimensional herringbone packing *via* $C2-H2\cdots$ Ring 2 (of $C1-C2-C3-C4-C5-N1$) interactions, as shown in Fig.2.

S2. Experimental


The title compound was synthesized as follows: a solution of benzene 1,4-dicarboxaldehyde (0.50 g, 3.73 mmol) in methanol (10 ml) was added dropwise to a stirred solution of 2-(pyridin-2-yl)ethanamine (0.91 g, 7.42 mmol) in methanol (10 ml). The mixture was stirred at room temperature for *ca* 16 h. The precipitate was filtered off and washed with diethyl ether and dried under vacuum for 4 h affording a fine shiny white powder in 85% yield. M.p.: does not melt below 260 °C. Recrystallization by slow diffusion of Et_2O into a concentrated CH_2Cl_2 of the solution gave colorless crystals suitable for X-ray structure analysis.

S3. Refinement

All non-hydrogen atoms were refined anisotropically and all hydrogen atoms were placed in idealized positions and refined with a riding model with U_{iso} set at 1.2 or 1.5 times U_{eq} of their parent atoms and fixed C—H bond lengths.

**Figure 1**

Molecular structure of titled compound showing the atomic numbering scheme. All non-hydrogen atoms were presented with ellipsoidal model with probability level 40%. Half of the molecule without atomic labels was generated *via* centre of symmetry (symmetry code: $-x, -y, -z$).

**Figure 2**

Projection viewed along $[100]$ showing 3-D herringbone packing. Only the hydrogen atoms that involved in $C-H\cdots N$ and $C-H\cdots\pi$ intermolecular interactions (see the list in Table 1) are shown and labelled. The red dotted lines represent the weak interactions.

1,4-Bis[(2-pyridylethyl)iminomethyl]benzene*Crystal data*C₂₂H₂₂N₄ $M_r = 342.44$ Monoclinic, $P2_1/n$

Hall symbol: -P 2yn

 $a = 6.0078$ (6) Å $b = 26.023$ (3) Å $c = 6.1319$ (7) Å $\beta = 106.009$ (2)° $V = 921.47$ (17) Å³ $Z = 2$ $F(000) = 364$ $D_x = 1.234$ Mg m⁻³Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 11941 reflections

 $\theta = 3.1$ – 28.3 ° $\mu = 0.08$ mm⁻¹ $T = 173$ K

Plate, colourless

 $0.26 \times 0.24 \times 0.17$ mm*Data collection*

Bruker Kappa DUO APEXII

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 0.5° φ scans and ω

11941 measured reflections

2288 independent reflections

1945 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.024$ $\theta_{\text{max}} = 28.3^\circ$, $\theta_{\text{min}} = 3.1^\circ$ $h = -8 \rightarrow 8$ $k = -34 \rightarrow 33$ $l = -8 \rightarrow 8$ *Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.040$ $wR(F^2) = 0.110$ $S = 1.06$

2288 reflections

118 parameters

0 restraints

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0506P)^2 + 0.2363P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\text{max}} < 0.001$ $\Delta\rho_{\text{max}} = 0.28$ e Å⁻³ $\Delta\rho_{\text{min}} = -0.20$ e Å⁻³*Special details***Experimental.** Data for (I):

IR (KBr): 1610 cm⁻¹ (C=N, imine) ¹HNMR:(CDCl₃) δ_{H} 8.55(ddd, 2H, $J = 0.8$ Hz, $J = 1.7$ Hz, $J = 4.8$ Hz) 8.21 (t, 2H, $J = 1.3$ Hz) 7.69 (s, 2H) 7.55 (dt, 2H, $J = 1.9$ Hz, $J = 7.7$ Hz) 7.11 (m, 2H) 4.03 (dt, 8H, $J = 1.2$ Hz, $J = 7.2$ Hz) 3.19(t, 4H, $J = 7.2$ Hz); ¹³CNMR: (400 MHz, CDCl₃) δ 161.05, 159.45, 149.37, 138.88, 136.13, 128.21, 123.67, 121.24, 61.18, 39.61; Analysis calculated for C₂₂H₂₂N₄: C, 77.16%; H, 6.48%; N, 16.36%; Found: C, 77.19%; H, 6.22%; N, 16.52%; EI—MS: m/z 249.90[M - C₇H₆N]⁺.

Half sphere of data collected using *SAINTE* strategy (Bruker, 2006). Crystal to detector distance = 50 mm; combination of φ and ω scans of 0.5° , 40 s per $^\circ$, 2 iterations.

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry.

An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	1.01974 (18)	0.16910 (4)	0.82099 (17)	0.0365 (3)
C1	1.1434 (2)	0.20288 (5)	0.9711 (2)	0.0431 (3)
H1	1.2675	0.2201	0.9339	0.052*
N2	0.40565 (16)	0.09448 (4)	0.35620 (15)	0.0282 (2)
C2	1.1022 (2)	0.21422 (5)	1.1750 (2)	0.0406 (3)
H2	1.1936	0.2389	1.2748	0.049*
C3	0.9248 (2)	0.18885 (5)	1.2306 (2)	0.0381 (3)
H3	0.8928	0.1952	1.3715	0.046*
C4	0.7937 (2)	0.15394 (4)	1.07884 (19)	0.0312 (3)
H4	0.6689	0.1363	1.1130	0.037*
C5	0.84667 (18)	0.14496 (4)	0.87569 (18)	0.0252 (2)
C6	0.7121 (2)	0.10708 (4)	0.7047 (2)	0.0327 (3)
H6A	0.6662	0.0778	0.7860	0.039*
H6B	0.8134	0.0936	0.6152	0.039*
C7	0.49617 (19)	0.13015 (4)	0.54368 (19)	0.0283 (2)
H7A	0.3778	0.1365	0.6254	0.034*
H7B	0.5346	0.1634	0.4844	0.034*
C8	0.21489 (18)	0.07288 (4)	0.34628 (17)	0.0251 (2)
H8	0.1384	0.0810	0.4584	0.030*
C9	0.10603 (17)	0.03556 (4)	0.16772 (17)	0.0231 (2)
C10	-0.10072 (18)	0.01157 (4)	0.17194 (18)	0.0259 (2)
H10	-0.1702	0.0196	0.2894	0.031*
C11	0.20546 (18)	0.02380 (4)	-0.00674 (18)	0.0254 (2)
H11	0.3453	0.0401	-0.0122	0.030*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0364 (6)	0.0409 (6)	0.0340 (5)	-0.0050 (4)	0.0129 (4)	0.0008 (4)
C1	0.0345 (6)	0.0402 (7)	0.0532 (8)	-0.0114 (5)	0.0095 (6)	0.0025 (6)
N2	0.0260 (5)	0.0295 (5)	0.0272 (5)	-0.0033 (4)	0.0044 (4)	-0.0064 (4)
C2	0.0428 (7)	0.0273 (6)	0.0400 (7)	-0.0030 (5)	-0.0081 (5)	-0.0037 (5)
C3	0.0552 (8)	0.0312 (6)	0.0266 (6)	0.0043 (5)	0.0089 (5)	-0.0037 (5)
C4	0.0354 (6)	0.0288 (5)	0.0317 (6)	-0.0010 (5)	0.0131 (5)	-0.0008 (4)
C5	0.0249 (5)	0.0240 (5)	0.0242 (5)	0.0031 (4)	0.0025 (4)	0.0002 (4)
C6	0.0342 (6)	0.0275 (6)	0.0311 (6)	0.0025 (5)	0.0000 (5)	-0.0061 (4)
C7	0.0257 (5)	0.0278 (5)	0.0296 (5)	-0.0019 (4)	0.0045 (4)	-0.0074 (4)
C8	0.0244 (5)	0.0252 (5)	0.0250 (5)	0.0007 (4)	0.0056 (4)	-0.0027 (4)
C9	0.0219 (5)	0.0217 (5)	0.0239 (5)	0.0004 (4)	0.0034 (4)	-0.0009 (4)
C10	0.0252 (5)	0.0284 (5)	0.0252 (5)	-0.0010 (4)	0.0088 (4)	-0.0026 (4)
C11	0.0214 (5)	0.0262 (5)	0.0289 (5)	-0.0030 (4)	0.0073 (4)	-0.0014 (4)

Geometric parameters (Å, °)

N1—C5	1.3344 (15)	C6—C7	1.5204 (16)
N1—C1	1.3395 (17)	C6—H6A	0.9900
C1—C2	1.372 (2)	C6—H6B	0.9900
C1—H1	0.9500	C7—H7A	0.9900
N2—C8	1.2628 (14)	C7—H7B	0.9900
N2—C7	1.4609 (13)	C8—C9	1.4748 (14)
C2—C3	1.3739 (19)	C8—H8	0.9500
C2—H2	0.9500	C9—C11	1.3954 (14)
C3—C4	1.3811 (17)	C9—C10	1.3968 (14)
C3—H3	0.9500	C10—C11 ⁱ	1.3848 (14)
C4—C5	1.3880 (15)	C10—H10	0.9500
C4—H4	0.9500	C11—C10 ⁱ	1.3848 (14)
C5—C6	1.5023 (15)	C11—H11	0.9500
C5—N1—C1	117.33 (10)	C7—C6—H6B	109.0
N1—C1—C2	124.19 (12)	H6A—C6—H6B	107.8
N1—C1—H1	117.9	N2—C7—C6	109.03 (9)
C2—C1—H1	117.9	N2—C7—H7A	109.9
C8—N2—C7	117.17 (9)	C6—C7—H7A	109.9
C1—C2—C3	118.02 (11)	N2—C7—H7B	109.9
C1—C2—H2	121.0	C6—C7—H7B	109.9
C3—C2—H2	121.0	H7A—C7—H7B	108.3
C2—C3—C4	119.10 (11)	N2—C8—C9	122.66 (9)
C2—C3—H3	120.4	N2—C8—H8	118.7
C4—C3—H3	120.4	C9—C8—H8	118.7
C3—C4—C5	119.07 (11)	C11—C9—C10	119.15 (9)
C3—C4—H4	120.5	C11—C9—C8	121.17 (9)
C5—C4—H4	120.5	C10—C9—C8	119.68 (9)
N1—C5—C4	122.27 (10)	C11 ⁱ —C10—C9	120.70 (9)
N1—C5—C6	116.12 (10)	C11 ⁱ —C10—H10	119.6
C4—C5—C6	121.60 (10)	C9—C10—H10	119.6
C5—C6—C7	113.13 (9)	C10 ⁱ —C11—C9	120.15 (9)
C5—C6—H6A	109.0	C10 ⁱ —C11—H11	119.9
C7—C6—H6A	109.0	C9—C11—H11	119.9
C5—C6—H6B	109.0		
C5—N1—C1—C2	-0.4 (2)	C8—N2—C7—C6	111.41 (11)
N1—C1—C2—C3	0.9 (2)	C5—C6—C7—N2	167.84 (9)
C1—C2—C3—C4	-1.10 (19)	C7—N2—C8—C9	-179.14 (9)
C2—C3—C4—C5	0.83 (18)	N2—C8—C9—C11	-2.76 (16)
C1—N1—C5—C4	0.06 (17)	N2—C8—C9—C10	177.48 (10)
C1—N1—C5—C6	-179.64 (10)	C11—C9—C10—C11 ⁱ	0.57 (17)
C3—C4—C5—N1	-0.30 (17)	C8—C9—C10—C11 ⁱ	-179.67 (9)
C3—C4—C5—C6	179.39 (10)	C10—C9—C11—C10 ⁱ	-0.57 (17)

N1—C5—C6—C7	-94.33 (12)	C8—C9—C11—C10 ⁱ	179.68 (9)
C4—C5—C6—C7	85.97 (13)		

Symmetry code: (i) $-x, -y, -z$.

Hydrogen-bond geometry (Å, °)

Cg1 and Cg2 are the centroids of the C1—C5/N1 and C9—C11/C9'—C11' rings, respectively.

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
C3—H3...N1 ⁱⁱ	0.95	2.74	3.544 (3)	143 (3)
C4—H4...N2 ⁱⁱ	0.95	2.69	3.593 (2)	159 (4)
C7—H7A...N1 ⁱⁱⁱ	0.99	2.87	3.847 (2)	171 (5)
C2—H2...Cg1 ^{iv}	0.95	2.88	3.826 (4)	172 (5)
C6—H6A...Cg2 ^v	0.99	2.90	3.508 (3)	120 (2)

Symmetry codes: (ii) $x, y, z+1$; (iii) $x-1, y, z$; (iv) $x+1/2, -y+1/2, z+1/2$; (v) $x+1, y, z+1$.