

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

N-(2-Fluorobenzoyl)-N',N''-bis(4-methylphenyl)phosphoric triamide

Mehrdad Pourayoubi,^{a*} Atekeh Tarahhomi,^a Arnold L. Rheingold^b and James A. Golen^b

^aDepartment of Chemistry, Ferdowsi University of Mashhad, Mashhad 91779, Iran,and ^bDepartment of Chemistry, University of California, San Diego, 9500 Gilman Drive, La Jolla, CA 92093, USA

Correspondence e-mail: mehrdad_pourayoubi@yahoo.com


Received 13 February 2011; accepted 14 March 2011

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.046; wR factor = 0.127; data-to-parameter ratio = 17.2.

The P atom in the title compound, $\text{C}_{21}\text{H}_{21}\text{FN}_3\text{O}_2\text{P}$, is in a tetrahedral coordination environment and the environment of each N atom is essentially planar (sums of angles = 359.7, 359.9 and 358.4°). The phosphoryl and carbonyl groups adopt *anti* orientations with respect to each other. In the crystal, adjacent molecules are linked *via* $\text{N}-\text{H}\cdots\text{O}=\text{P}$ and two $\text{N}-\text{H}\cdots\text{O}=\text{C}$ hydrogen bonds into an extended chain parallel to the *a* axis.

Related literature

For a phosphorus ligand having a $\text{C}(\text{O})\text{NHP}(\text{O})$ skeleton, see: Gholivand *et al.* (2010). For a related structure, see: Pourayoubi *et al.* (2010). For bond lengths in related structures, see: Sabbaghi *et al.* (2010) and references cited therein.



Experimental

Crystal data

 $\text{C}_{21}\text{H}_{21}\text{FN}_3\text{O}_2\text{P}$ $M_r = 397.38$ Monoclinic, $P2_1/n$ $a = 9.7697$ (9) Å $b = 10.2197$ (9) Å $c = 20.2404$ (18) Å $\beta = 96.605$ (1)° $V = 2007.5$ (3) Å³ $Z = 4$ Mo $K\alpha$ radiation $\mu = 0.17$ mm⁻¹ $T = 100$ K $0.25 \times 0.15 \times 0.15$ mm

Data collection

Bruker SMART CCD area-detector diffractometer

Absorption correction: multi-scan (SADABS; Bruker, 2005)

 $T_{\min} = 0.959$, $T_{\max} = 0.975$

15865 measured reflections

4551 independent reflections

3411 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.035$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.046$ $wR(F^2) = 0.127$ $S = 1.05$

4551 reflections

264 parameters

3 restraints

H atoms treated by a mixture of independent and constrained refinement

 $\Delta\rho_{\text{max}} = 0.41$ e Å⁻³ $\Delta\rho_{\text{min}} = -0.26$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1N}\cdots\text{O2}^{\text{i}}$	0.87 (1)	1.92 (1)	2.780 (2)	171 (2)
$\text{N2}-\text{H2N}\cdots\text{O1}^{\text{ii}}$	0.86 (1)	2.08 (1)	2.886 (2)	156 (2)
$\text{N3}-\text{H3N}\cdots\text{O1}^{\text{ii}}$	0.86 (1)	2.24 (2)	2.945 (2)	139 (2)

Symmetry codes: (i) $-x + 1, -y + 2, -z + 1$; (ii) $-x + 2, -y + 2, -z + 1$.

Data collection: SMART (Bruker, 2005); cell refinement: SAINT (Bruker, 2005); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL and enCIFer (Allen *et al.*, 2004).

Support of this investigation by Ferdowsi University of Mashhad is gratefully acknowledged. The authors wish to thank Bruker AXS Inc. for the use of one of their SMART X2S benchtop instruments.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NC2222).

References

- Allen, F. H., Johnson, O., Shields, G. P., Smith, B. R. & Towler, M. (2004). *J. Appl. Cryst.* **37**, 335–338.
- Bruker (2005). *SADABS*, *SMART* and *SAINTE*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Gholivand, K., Mahzouni, H. R., Pourayoubi, M. & Amiri, S. (2010). *Inorg. Chim. Acta*, **363**, 2318–2324.
- Pourayoubi, M., Tarahhomi, A., Rheingold, A. L. & Golen, J. A. (2010). *Acta Cryst.* **E66**, o2524.
- Sabbaghi, F., Pourayoubi, M., Toghrace, M. & Divjakovic, V. (2010). *Acta Cryst.* **E66**, o344.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supporting information

Acta Cryst. (2011). E67, o934 [doi:10.1107/S1600536811009640]

***N*-(2-Fluorobenzoyl)-*N',N''*-bis(4-methylphenyl)phosphoric triamide**

Mehrdad Pourayoubi, Atekeh Tarahhomi, Arnold L. Rheingold and James A. Golen

S1. Comment

Carbacylamidophosphates with a C(O)NHP(O) skeleton have attracted attention because of their roles as the *O,O'*-donor ligands for metal complexation (Gholivand *et al.*, 2010). Following the previous works about carbacylamidophosphates such as P(O)[NHC(O)C₆H₃(2,6-F₂)] [N(CH₃)(CH₂C₆H₅)₂] (Pourayoubi *et al.*, 2010), here, we report on the synthesis and crystal structure of the title compound, P(O)[NHC(O)C₆H₄(2-F)] [NH—C₆H₄-4-CH₃]₂.

In the crystal structure of the title compound the phosphoryl and carbonyl groups adopt *anti* positions to each other. The P atom has a slightly distorted tetrahedral configuration (Fig. 1). The bond angles around the P atom are in the range of 101.09 (8)° to 116.67 (8)°. The P1—N2 and P1—N3 bonds (1.6361 (15) Å and 1.6291 (17) Å) are shorter than the P1—N1 bond (1.6872 (15) Å). The environment of the nitrogen atoms is essentially planar. The P=O bond length of 1.4723 (13) Å is comparable to those in similar compounds *e.g.* in P(O)[NHC(O)C₆H₄(4-NO₂)] [NHC₆H₁₁]₂ (Sabbaghi *et al.*, 2010).


In the crystal structure, adjacent molecules are linked *via* N_{C(O)NHP(O)}—H···O=P and two N_{amide}—H···O=C hydrogen bonds (see Table 1), into an extended chain parallel to the *a* axis.

S2. Experimental

2-F—C₆H₄C(O)NHP(O)Cl₂ has been synthesized from the reaction between phosphorus pentachloride (4.0 g, 19.2 mmol) and 2-fluorobenzamide (2.671 g, 19.2 mmol) in dry CCl₄ at 358 K (3 h) and then the treatment of formic acid (0.884 g, 19.2 mmol) at ice bath temperature. To a solution of 2-F—C₆H₄C(O)NHP(O)Cl₂ (0.3 g, 1.17 mmol) in dry chloroform (30 ml), a mixture of *p*-toluidine (0.251 g, 2.34 mmol) and triethylamine (0.237 g, 2.34 mmol) in dry chloroform (10 ml) was added at 273 K. After 4 h stirring, the solvent was removed and the product was washed with distilled water and recrystallized from methanol/chloroform at room temperature. IR (KBr, cm⁻¹): 3308 (NH), 3030 (NH), 2896, 2627, 1639 (C=O), 1457, 1220, 1061, 944, 795.

S3. Refinement

Hydrogen atoms H1N, H2N, and H3N were located in Fourier difference map and were refined with *DFIX* 0.88 (0.01) for the N—H bond lengths and isotropic displacement parameter of 1.2 times *U*_{eq} of the parent N atoms. All other hydrogen atoms were placed in their calculated positions with atom—H lengths of 0.95 Å (CH) and 0.98 Å (CH₃) and isotropic displacement parameters for these atoms were set to 1.20 times (CH) and 1.50 times (CH₃) *U*_{eq} of the parent C atom.

**Figure 1**

An ORTEP-style plot of title compound with labeling. Displacement ellipsoids are drawn at the 50% probability level.

***N*-{bis[(4-methylphenyl)amino]phosphoryl}-2-fluorobenzamide**

Crystal data

$C_{21}H_{21}FN_3O_2P$

$M_r = 397.38$

Monoclinic, $P2_1/n$

$a = 9.7697$ (9) Å

$b = 10.2197$ (9) Å

$c = 20.2404$ (18) Å

$\beta = 96.605$ (1)°

$V = 2007.5$ (3) Å³

$Z = 4$

$F(000) = 832$

$D_x = 1.315$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 5744 reflections

$\theta = 2.2$ – 27.5 °

$\mu = 0.17$ mm⁻¹

$T = 100$ K

Block, colourless

$0.25 \times 0.15 \times 0.15$ mm

Data collection

Bruker SMART CCD area-detector
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

φ and ω scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2005)

$T_{\min} = 0.959$, $T_{\max} = 0.975$

15865 measured reflections

4551 independent reflections

3411 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.035$

$\theta_{\text{max}} = 27.9$ °, $\theta_{\text{min}} = 2.0$ °

$h = -12 \rightarrow 12$

$k = -11 \rightarrow 13$

$l = -25 \rightarrow 25$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.046$
 $wR(F^2) = 0.127$
 $S = 1.05$
 4551 reflections
 264 parameters
 3 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H atoms treated by a mixture of independent
 and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0669P)^2 + 0.498P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.007$
 $\Delta\rho_{\max} = 0.41 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.26 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
P1	0.72255 (5)	0.92119 (4)	0.50855 (2)	0.02136 (14)
F1	0.92737 (14)	1.40039 (12)	0.55287 (6)	0.0441 (3)
O1	0.91560 (13)	1.14443 (13)	0.50431 (6)	0.0293 (3)
O2	0.58757 (12)	0.86140 (12)	0.51487 (6)	0.0250 (3)
N1	0.69300 (16)	1.08365 (14)	0.50258 (7)	0.0210 (3)
H1N	0.6074 (11)	1.1095 (19)	0.4983 (9)	0.025*
N2	0.79772 (16)	0.87866 (15)	0.44363 (8)	0.0240 (3)
H2N	0.8811 (12)	0.8507 (19)	0.4525 (10)	0.029*
N3	0.84260 (16)	0.88686 (16)	0.56874 (8)	0.0267 (4)
H3N	0.9270 (12)	0.905 (2)	0.5636 (10)	0.032*
C1	0.8139 (2)	1.41838 (19)	0.50898 (9)	0.0296 (4)
C2	0.7701 (2)	1.5451 (2)	0.49516 (11)	0.0381 (5)
H2C	0.8194	1.6173	0.5157	0.046*
C3	0.6533 (2)	1.5647 (2)	0.45090 (11)	0.0379 (5)
H3B	0.6219	1.6511	0.4408	0.045*
C4	0.5822 (2)	1.4596 (2)	0.42132 (11)	0.0344 (5)
H4A	0.5029	1.4737	0.3903	0.041*
C5	0.6265 (2)	1.33331 (19)	0.43680 (9)	0.0278 (4)
H5A	0.5765	1.2612	0.4166	0.033*
C6	0.74372 (19)	1.31052 (17)	0.48164 (9)	0.0230 (4)
C7	0.79305 (18)	1.17460 (18)	0.49728 (8)	0.0220 (4)
C8	0.7441 (2)	0.89104 (17)	0.37612 (9)	0.0244 (4)
C9	0.6203 (2)	0.9545 (2)	0.35614 (10)	0.0319 (5)
H9A	0.5685	0.9914	0.3884	0.038*

C10	0.5725 (2)	0.9640 (2)	0.28921 (10)	0.0360 (5)
H10A	0.4886	1.0089	0.2762	0.043*
C11	0.6439 (2)	0.9096 (2)	0.24070 (10)	0.0404 (5)
C12	0.7657 (2)	0.8443 (2)	0.26118 (10)	0.0419 (6)
H12A	0.8157	0.8051	0.2288	0.050*
C13	0.8164 (2)	0.8347 (2)	0.32801 (10)	0.0341 (5)
H13A	0.9004	0.7898	0.3409	0.041*
C14	0.5884 (3)	0.9202 (3)	0.16774 (11)	0.0561 (7)
H14A	0.6098	0.8398	0.1446	0.084*
H14B	0.6313	0.9950	0.1479	0.084*
H14C	0.4884	0.9326	0.1636	0.084*
C15	0.8269 (2)	0.85706 (18)	0.63606 (9)	0.0260 (4)
C16	0.7217 (2)	0.7756 (2)	0.65213 (10)	0.0406 (5)
H16A	0.6570	0.7408	0.6180	0.049*
C17	0.7115 (2)	0.7452 (3)	0.71817 (11)	0.0506 (7)
H17A	0.6367	0.6929	0.7288	0.061*
C18	0.8072 (3)	0.7891 (2)	0.76906 (10)	0.0435 (6)
C19	0.9162 (3)	0.8633 (2)	0.75138 (11)	0.0507 (7)
H19A	0.9862	0.8909	0.7850	0.061*
C20	0.9253 (3)	0.8980 (2)	0.68584 (11)	0.0427 (6)
H20A	1.0000	0.9504	0.6752	0.051*
C21	0.7964 (3)	0.7510 (3)	0.84055 (11)	0.0657 (8)
H21A	0.8791	0.7801	0.8687	0.099*
H21B	0.7879	0.6557	0.8437	0.099*
H21C	0.7150	0.7926	0.8556	0.099*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
P1	0.0143 (3)	0.0263 (2)	0.0236 (2)	0.00103 (18)	0.00229 (18)	0.00314 (18)
F1	0.0442 (8)	0.0440 (7)	0.0404 (7)	-0.0137 (6)	-0.0116 (6)	-0.0023 (5)
O1	0.0150 (7)	0.0379 (8)	0.0351 (8)	-0.0004 (6)	0.0030 (6)	0.0005 (6)
O2	0.0142 (7)	0.0278 (7)	0.0330 (7)	-0.0001 (5)	0.0025 (5)	0.0038 (5)
N1	0.0143 (8)	0.0244 (8)	0.0245 (8)	0.0002 (6)	0.0038 (6)	0.0007 (6)
N2	0.0158 (8)	0.0297 (8)	0.0262 (8)	0.0048 (6)	0.0015 (6)	-0.0003 (6)
N3	0.0133 (8)	0.0388 (9)	0.0279 (9)	0.0000 (7)	0.0018 (7)	0.0086 (6)
C1	0.0288 (11)	0.0346 (11)	0.0252 (10)	-0.0079 (9)	0.0024 (8)	-0.0004 (8)
C2	0.0497 (15)	0.0292 (10)	0.0368 (12)	-0.0104 (10)	0.0106 (10)	-0.0051 (9)
C3	0.0433 (14)	0.0280 (11)	0.0443 (12)	0.0005 (9)	0.0135 (11)	0.0062 (9)
C4	0.0271 (11)	0.0353 (11)	0.0411 (12)	0.0002 (9)	0.0058 (9)	0.0084 (9)
C5	0.0231 (10)	0.0292 (10)	0.0317 (10)	-0.0033 (8)	0.0057 (8)	0.0011 (8)
C6	0.0202 (10)	0.0273 (9)	0.0227 (9)	-0.0035 (7)	0.0073 (7)	-0.0015 (7)
C7	0.0167 (10)	0.0303 (9)	0.0192 (9)	-0.0013 (7)	0.0033 (7)	-0.0019 (7)
C8	0.0212 (10)	0.0274 (9)	0.0247 (9)	-0.0034 (8)	0.0032 (8)	0.0010 (7)
C9	0.0270 (11)	0.0404 (11)	0.0278 (10)	0.0020 (9)	0.0019 (8)	-0.0030 (8)
C10	0.0290 (12)	0.0461 (12)	0.0313 (11)	0.0001 (10)	-0.0038 (9)	0.0029 (9)
C11	0.0345 (13)	0.0601 (15)	0.0262 (11)	-0.0110 (11)	0.0015 (9)	0.0024 (9)
C12	0.0351 (13)	0.0624 (15)	0.0300 (12)	-0.0047 (11)	0.0109 (10)	-0.0062 (10)

C13	0.0254 (11)	0.0452 (12)	0.0324 (11)	0.0013 (9)	0.0060 (9)	-0.0017 (9)
C14	0.0465 (16)	0.095 (2)	0.0262 (12)	-0.0108 (14)	0.0008 (11)	0.0069 (12)
C15	0.0234 (10)	0.0299 (10)	0.0248 (10)	0.0057 (8)	0.0040 (8)	0.0025 (7)
C16	0.0266 (12)	0.0627 (15)	0.0318 (11)	-0.0063 (10)	0.0001 (9)	0.0129 (10)
C17	0.0322 (13)	0.0782 (18)	0.0421 (14)	-0.0019 (12)	0.0075 (11)	0.0229 (12)
C18	0.0496 (15)	0.0558 (14)	0.0263 (11)	0.0136 (12)	0.0098 (10)	0.0039 (9)
C19	0.0709 (19)	0.0504 (14)	0.0284 (12)	-0.0100 (13)	-0.0048 (12)	-0.0038 (10)
C20	0.0499 (15)	0.0433 (13)	0.0336 (12)	-0.0166 (11)	-0.0006 (10)	0.0005 (9)
C21	0.074 (2)	0.095 (2)	0.0310 (13)	0.0190 (17)	0.0163 (13)	0.0111 (13)

Geometric parameters (Å, °)

P1—O2	1.4723 (13)	C9—H9A	0.9500
P1—N3	1.6291 (17)	C10—C11	1.385 (3)
P1—N2	1.6361 (15)	C10—H10A	0.9500
P1—N1	1.6872 (15)	C11—C12	1.386 (3)
F1—C1	1.351 (2)	C11—C14	1.517 (3)
O1—C7	1.229 (2)	C12—C13	1.389 (3)
N1—C7	1.362 (2)	C12—H12A	0.9500
N1—H1N	0.872 (9)	C13—H13A	0.9500
N2—C8	1.412 (2)	C14—H14A	0.9800
N2—H2N	0.862 (9)	C14—H14B	0.9800
N3—C15	1.422 (2)	C14—H14C	0.9800
N3—H3N	0.863 (9)	C15—C20	1.375 (3)
C1—C6	1.380 (3)	C15—C16	1.390 (3)
C1—C2	1.382 (3)	C16—C17	1.387 (3)
C2—C3	1.382 (3)	C16—H16A	0.9500
C2—H2C	0.9500	C17—C18	1.384 (3)
C3—C4	1.378 (3)	C17—H17A	0.9500
C3—H3B	0.9500	C18—C19	1.387 (3)
C4—C5	1.385 (3)	C18—C21	1.514 (3)
C4—H4A	0.9500	C19—C20	1.386 (3)
C5—C6	1.396 (3)	C19—H19A	0.9500
C5—H5A	0.9500	C20—H20A	0.9500
C6—C7	1.492 (3)	C21—H21A	0.9800
C8—C9	1.391 (3)	C21—H21B	0.9800
C8—C13	1.392 (3)	C21—H21C	0.9800
C9—C10	1.384 (3)		
O2—P1—N3	114.88 (8)	C9—C10—C11	121.7 (2)
O2—P1—N2	116.67 (8)	C9—C10—H10A	119.2
N3—P1—N2	101.09 (8)	C11—C10—H10A	119.2
O2—P1—N1	105.49 (8)	C10—C11—C12	117.8 (2)
N3—P1—N1	111.59 (8)	C10—C11—C14	120.6 (2)
N2—P1—N1	107.03 (8)	C12—C11—C14	121.6 (2)
C7—N1—P1	123.97 (13)	C11—C12—C13	121.5 (2)
C7—N1—H1N	118.3 (13)	C11—C12—H12A	119.2
P1—N1—H1N	117.4 (13)	C13—C12—H12A	119.2

C8—N2—P1	127.06 (13)	C12—C13—C8	120.0 (2)
C8—N2—H2N	117.9 (13)	C12—C13—H13A	120.0
P1—N2—H2N	114.9 (13)	C8—C13—H13A	120.0
C15—N3—P1	127.97 (13)	C11—C14—H14A	109.5
C15—N3—H3N	111.7 (14)	C11—C14—H14B	109.5
P1—N3—H3N	118.7 (14)	H14A—C14—H14B	109.5
F1—C1—C6	119.15 (18)	C11—C14—H14C	109.5
F1—C1—C2	118.14 (18)	H14A—C14—H14C	109.5
C6—C1—C2	122.7 (2)	H14B—C14—H14C	109.5
C3—C2—C1	118.7 (2)	C20—C15—C16	118.94 (18)
C3—C2—H2C	120.7	C20—C15—N3	119.60 (18)
C1—C2—H2C	120.7	C16—C15—N3	121.11 (18)
C4—C3—C2	120.4 (2)	C17—C16—C15	119.8 (2)
C4—C3—H3B	119.8	C17—C16—H16A	120.1
C2—C3—H3B	119.8	C15—C16—H16A	120.1
C3—C4—C5	119.9 (2)	C18—C17—C16	121.8 (2)
C3—C4—H4A	120.0	C18—C17—H17A	119.1
C5—C4—H4A	120.0	C16—C17—H17A	119.1
C4—C5—C6	120.93 (19)	C17—C18—C19	117.3 (2)
C4—C5—H5A	119.5	C17—C18—C21	120.8 (2)
C6—C5—H5A	119.5	C19—C18—C21	121.8 (2)
C1—C6—C5	117.37 (18)	C20—C19—C18	121.5 (2)
C1—C6—C7	121.68 (18)	C20—C19—H19A	119.2
C5—C6—C7	120.92 (16)	C18—C19—H19A	119.2
O1—C7—N1	121.22 (17)	C15—C20—C19	120.5 (2)
O1—C7—C6	123.01 (16)	C15—C20—H20A	119.8
N1—C7—C6	115.77 (15)	C19—C20—H20A	119.8
C9—C8—C13	118.93 (18)	C18—C21—H21A	109.5
C9—C8—N2	122.47 (16)	C18—C21—H21B	109.5
C13—C8—N2	118.58 (17)	H21A—C21—H21B	109.5
C10—C9—C8	120.08 (18)	C18—C21—H21C	109.5
C10—C9—H9A	120.0	H21A—C21—H21C	109.5
C8—C9—H9A	120.0	H21B—C21—H21C	109.5
O2—P1—N1—C7	178.66 (14)	C5—C6—C7—N1	-39.0 (2)
N3—P1—N1—C7	53.27 (16)	P1—N2—C8—C9	6.1 (3)
N2—P1—N1—C7	-56.46 (16)	P1—N2—C8—C13	-172.26 (15)
O2—P1—N2—C8	57.27 (18)	C13—C8—C9—C10	-1.7 (3)
N3—P1—N2—C8	-177.42 (15)	N2—C8—C9—C10	179.93 (18)
N1—P1—N2—C8	-60.53 (17)	C8—C9—C10—C11	1.1 (3)
O2—P1—N3—C15	-28.7 (2)	C9—C10—C11—C12	0.3 (3)
N2—P1—N3—C15	-155.21 (16)	C9—C10—C11—C14	179.3 (2)
N1—P1—N3—C15	91.30 (17)	C10—C11—C12—C13	-0.9 (3)
F1—C1—C2—C3	-179.31 (17)	C14—C11—C12—C13	180.0 (2)
C6—C1—C2—C3	-1.7 (3)	C11—C12—C13—C8	0.3 (3)
C1—C2—C3—C4	0.1 (3)	C9—C8—C13—C12	1.0 (3)
C2—C3—C4—C5	1.1 (3)	N2—C8—C13—C12	179.47 (19)
C3—C4—C5—C6	-0.7 (3)	P1—N3—C15—C20	-144.91 (18)

F1—C1—C6—C5	179.64 (15)	P1—N3—C15—C16	41.9 (3)
C2—C1—C6—C5	2.0 (3)	C20—C15—C16—C17	4.9 (3)
F1—C1—C6—C7	-2.4 (3)	N3—C15—C16—C17	178.1 (2)
C2—C1—C6—C7	179.95 (17)	C15—C16—C17—C18	-3.0 (4)
C4—C5—C6—C1	-0.8 (3)	C16—C17—C18—C19	-1.2 (4)
C4—C5—C6—C7	-178.77 (17)	C16—C17—C18—C21	-178.0 (2)
P1—N1—C7—O1	-8.6 (2)	C17—C18—C19—C20	3.3 (4)
P1—N1—C7—C6	170.55 (12)	C21—C18—C19—C20	-179.9 (2)
C1—C6—C7—O1	-37.7 (3)	C16—C15—C20—C19	-2.8 (3)
C5—C6—C7—O1	140.17 (18)	N3—C15—C20—C19	-176.1 (2)
C1—C6—C7—N1	143.13 (17)	C18—C19—C20—C15	-1.4 (4)

Hydrogen-bond geometry (Å, °)

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
N1—H1N...O2 ⁱ	0.87 (1)	1.92 (1)	2.780 (2)	171 (2)
N2—H2N...O1 ⁱⁱ	0.86 (1)	2.08 (1)	2.886 (2)	156 (2)
N3—H3N...O1 ⁱⁱ	0.86 (1)	2.24 (2)	2.945 (2)	139 (2)

Symmetry codes: (i) $-x+1, -y+2, -z+1$; (ii) $-x+2, -y+2, -z+1$.