

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

1-(4-Bromophenyl)-2-ethylsulfinyl-2-(phenylselanyl)ethanone monohydrate

Julio Zukerman-Schpector,^{a*} Carlos A. De Simone,^b
Paulo R. Olivato,^c Carlos R. Cerqueira Jr^c and
Edward R. T. Tiekink^d

^aDepartment of Chemistry, Universidade Federal de São Carlos, 13565-905 São Carlos, SP, Brazil, ^bInstituto de Química e Biotecnologia, Universidade Federal de Alagoas, 57072-970 Maceió, AL, Brazil, ^cChemistry Institute, Universidade de São Paulo, 05508-000 São Paulo-SP, Brazil, and ^dDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
Correspondence e-mail: julio@power.ufscar.br


Received 3 April 2011; accepted 5 April 2011

Key indicators: single-crystal X-ray study; $T = 290$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.037; wR factor = 0.095; data-to-parameter ratio = 18.7.

In the title hydrate, $\text{C}_{16}\text{H}_{15}\text{BrO}_2\text{SSe}\cdot\text{H}_2\text{O}$, the sulfinyl O atom lies on the opposite side of the molecule to the Se and carbonyl O atoms. The benzene rings form a dihedral angle of 51.66 (17) $^\circ$ and are splayed with respect to each other. The observed conformation allows the water molecules to bridge sulfinyl O atoms *via* $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds, generating a linear supramolecular chain along the b axis; the chain is further stabilized by $\text{C}-\text{H}\cdots\text{O}$ contacts. The chains are held in place in the crystal structure by $\text{C}\cdots\text{H}\cdots\pi$ and $\text{C}-\text{Br}\cdots\pi$ interactions.

Related literature

For background to β,β -bis-substituted-carbonyl compounds, see: Reis *et al.* (2006). For related structures, see: Olivato *et al.* (2004); Zukerman-Schpector *et al.* (2009, 2010). For details of the synthetic protocols, see: Long (1946); Leonard & Johnson (1962); Zoretic & Soja (1976).



Experimental

Crystal data

$\text{C}_{16}\text{H}_{15}\text{BrO}_2\text{SSe}\cdot\text{H}_2\text{O}$
 $M_r = 448.23$
Monoclinic, $P2_1/c$
 $a = 14.6942$ (2) Å
 $b = 6.1103$ (1) Å
 $c = 21.5717$ (4) Å
 $\beta = 113.714$ (1) $^\circ$
 $V = 1773.30$ (5) Å³
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 4.50$ mm⁻¹
 $T = 290$ K
 $0.36 \times 0.19 \times 0.16$ mm

Data collection

Nonius KappaCCD diffractometer
Absorption correction: multi-scan
(*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.291$, $T_{\max} = 0.734$
32063 measured reflections
3734 independent reflections
3177 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.076$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.037$
 $wR(F^2) = 0.095$
 $S = 1.03$
3734 reflections
200 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.80$ e Å⁻³
 $\Delta\rho_{\min} = -0.55$ e Å⁻³

Table 1

 Hydrogen-bond geometry (Å, $^\circ$).

Cg1 and Cg2 are the centroids of the $\text{C5}-\text{C10}$ and $\text{C11}-\text{C16}$ rings, respectively.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O1w}-\text{H1w}\cdots\text{O2}^i$	0.85	1.95	2.788 (4)	169
$\text{O1w}-\text{H2w}\cdots\text{O2}$	0.84	1.99	2.810 (4)	165
$\text{C2}-\text{H2}\cdots\text{O1w}^i$	0.98	2.40	3.334 (4)	159
$\text{C3}-\text{H3b}\cdots\text{O1w}^i$	0.97	2.54	3.434 (4)	153
$\text{C9}-\text{H9}\cdots\text{O1w}^{ii}$	0.93	2.55	3.320 (4)	141
$\text{C10}-\text{H10}\cdots\text{O2}^{ii}$	0.93	2.58	3.456 (4)	157
$\text{C14}-\text{H14}\cdots\text{Cg1}^{iii}$	0.93	2.96	3.793 (5)	149
$\text{C8}-\text{Br}\cdots\text{Cg2}^{iv}$	1.90 (1)	3.49 (1)	5.349 (3)	165 (1)

Symmetry codes: (i) $-x + 1, y - \frac{1}{2}, -z + \frac{1}{2}$; (ii) $x, y - 1, z$; (iii) $-x, y + \frac{1}{2}, -z + \frac{1}{2}$; (iv) $x, -y - \frac{1}{2}, z - \frac{1}{2}$.

Data collection: *COLLECT* (Nonius, 1999); cell refinement: *SCALEPACK* (Otwinowski & Minor, 1997); data reduction: *DENZO* (Otwinowski & Minor, 1997) and *SCALEPACK*; program(s) used to solve structure: *SIR97* (Altomare *et al.*, 1999); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *DIAMOND* (Brandenburg, 2006); software used to prepare material for publication: *MarvinSketch* (Chemaxon, 2010) and *pubCIF* (Westrip, 2010).

We thank the Brazilian agencies FAPESP, CNPq (fellowships to JZS and PRO) and CAPES (808/2009 to JZS) for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HG5022).

References

- Altomare, A., Burla, M. C., Camalli, M., Cascarano, G. L., Giacovazzo, C., Guagliardi, A., Moliterni, A. G. G., Polidori, G. & Spagna, R. (1999). *J. Appl. Cryst.* **32**, 115–119.
Brandenburg, K. (2006). *DIAMOND*. Crystal Impact GbR, Bonn, Germany. Chemaxon (2010). *MarvinSketch*. <http://www.chemaxon.com>.
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
Leonard, N. J. & Johnson, C. R. (1962). *J. Org. Chem.* **27**, 282–284.

- Long, L. M. (1946). *J. Am. Chem. Soc.* **68**, 2159–2161.
- Nonius (1999). *COLLECT*. Nonius BV, Delft, The Netherlands.
- Olivato, P. R., Reis, A. K. C. A., Rodrigues, A., Zukerman-Schpector, J., Tormena, C. F., Rittner, R. & Dal Colle, M. (2004). *J. Mol. Struct.* **707**, 199–210.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Reis, A. K. C. A., Olivato, P. R., Zukerman-Schpector, J., Tormena, C. F. J., Rittner, R. & Dal Colle, M. (2006). *J. Mol. Struct.* **798**, 57–63.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.
- Zoretic, P. A. & Soja, P. (1976). *J. Org. Chem.* **41**, 3587–3589.
- Zukerman-Schpector, J., De Simone, C. A., Olivato, P. R., Cerqueira, C. R., Santos, J. M. M. & Tiekink, E. R. T. (2010). *Acta Cryst.* **E66**, o1863.
- Zukerman-Schpector, J., Vinhato, E., Olivato, P. R., Rodrigues, A., Dal Colle, M., Cerqueira, C. R. Jr, Arman, H. D. & Tiekink, E. R. T. (2009). *Z. Kristallogr.* **224**, 484–492.

supporting information

Acta Cryst. (2011). E67, o1099–o1100 [doi:10.1107/S1600536811012712]

1-(4-Bromophenyl)-2-ethylsulfinyl-2-(phenylselanyl)ethanone monohydrate

Julio Zukerman-Schpector, Carlos A. De Simone, Paulo R. Olivato, Carlos R. Cerqueira and Edward R. T. Tiekink

S1. Comment

As part of our on-going research on the conformational and electronic interactions in some β,β -substituted-carbonyl compounds, *e.g.* 4'-substituted 2-(bromo)-2-(ethylsulfonyl)- and 4'-substituted 2-(methylthio)-2-(diethoxyphosphoryl)]-acetophenones, and 3,3-bis[(4'-chlorophenyl)thio]-1-methylpiperidin-2-one, using theoretical, spectroscopic and X-ray diffraction methods (Olivato *et al.*, 2004; Reis *et al.*, 2006; Zukerman-Schpector *et al.*, 2009; Zukerman-Schpector *et al.*, 2010), the title hydrate, (I), was synthesized and its crystal structure determined, Fig. 1.

With reference to the pyramidal-S atom, the sulfinyl-O lies to the opposite side of the molecule to each of the Se and carbonyl-O atoms. This conformation allows for the formation of supramolecular chains mediated by the sulfinyl-O and water molecules, see below. The benzene rings are splayed with respect to each other as seen in the value of the C1—C2—Se—C11 torsion angle of $-27.7(2)^\circ$; the dihedral angle formed between the rings is $51.66(17)^\circ$.

In the crystal packing, the water molecules bridge sulfinyl-O atoms *via* O—H \cdots O hydrogen bonds to form a linear supramolecular chain along the *b* axis, Fig. 2 and Table 1. Chains are stabilized by a series of C—H \cdots O interactions, Table 1, and are held in place by C—H $\cdots\pi$ (aryl-Br) and C—Br $\cdots\pi$ (aryl-Se) interactions, Fig. 3 and Table 1.

S2. Experimental

Following the procedure of Long (1946), a solution of potassium hydroxide (400 mg, 7.2 mmol) and ethanethiol (0.5 ml, 7.2 mmol) in ethanol (10 ml) was added to a solution of 2-bromo-4'-bromoacetophenone (2.0 g, 7.2 mmol) in ethanol, to give 2-ethylthio-4'-bromoacetophenone (1.6 g, yield = 86%). The product was isolated and oxidized with 12 ml of an aqueous solution of sodium periodate (0.5 M) in acetonitrile (16 ml), after Leonard & Johnson (1962), to give 2-ethylsulfinyl-4'-bromoacetophenone that was extracted with dichloromethane and dried over anhydrous magnesium sulfate. 2-Ethylsulfinyl-4'-bromoacetophenone (730 mg, 2.6 mmol) was added drop-wise to a cooled (195 K) solution of diisopropylamine (0.4 ml, 2.6 mmol) and butyllithium (2.3 ml, 2.6 mmol) in THF (20 ml). After 20 minutes, phenylselenenylbromide (610 mg, 2.6 mmol) dissolved in THF (10 ml) was added drop-wise to the enolate solution (Zoretic and Soja, 1976). After stirring for 3 h at 195 K, water (50 ml) was added at room temperature and extraction with chloroform was performed. The organic layer was dried over anhydrous magnesium sulfate. After evaporation of solvent, a crude solid was obtained. Purification through flash chromatography with a solution of hexane and ethyl acetate in a 1:1 ratio gave a mixture of the two possible diastereoisomers (500 mg, yield = 45%). One of the diastereoisomers was separated by recrystallization at low temperature (283 K) from chloroform. Suitable crystals for X-ray analysis were obtained by vapour diffusion of *n*-hexane into its chloroform solution at 283 K; *M.pt.* 366–367 K. IR (cm⁻¹): $\nu(\text{C=O})$ 1670, $\nu(\text{S=O})$ 993. NMR (CDCl₃, p.p.m.): δ 1.42–1.45 (3H, t, $^3J = 7.5$ Hz), 2.92–2.99 (1H, dq, $^2J = 13$ Hz, $^3J = 7.5$ Hz), 3.32–3.25 (1H, dq, $^2J = 13$ Hz, $^3J = 7.5$ Hz), 5.44 (1H, s), 7.29–7.33 (2H, m, Aryl-H), 7.38–7.41 (1H, m, Aryl-H), 7.52–7.55 (2H, m, Aryl-H), 7.59–7.62 (2H, m, Aryl-H), 7.75–7.73 (2H, m, Aryl-H). Analysis found: C 42.76, H 3.84%. C₁₆H₁₅BrO₂SSe.H₂O

requires: C 42.87, H 3.82%.

S3. Refinement

The H atoms were geometrically placed ($C-H = 0.93-0.98 \text{ \AA}$) and refined as riding with $U_{iso}(H) = 1.2-1.5U_{eq}(C)$. Those of the water molecule were found in a difference map, fixed in those positions and refined with $U_{iso}(H) = 1.2U_{eq}(O)$; see Table 1 for distances.






Figure 1

The molecular structure of (I) showing atom labelling scheme and displacement ellipsoids at the 35% probability level (arbitrary spheres for the H atoms).

**Figure 2**

Supramolecular linear chain along the *b* axis in (I) mediated by O—H...O hydrogen bonding (orange dashed lines).

**Figure 3**

View of the unit-cell contents in projection down the *b* axis in (I). Chains shown in Fig. 2, sustained by O—H...O hydrogen bonding (orange dashed lines), are held in place by C—H... π and C—Br... π contacts, shown as blue and purple dashed lines, respectively.

1-(4-Bromophenyl)-2-ethylsulfinyl-2-(phenylselanyl)ethanone monohydrate

Crystal data

C₁₆H₁₅BrO₂SSe·H₂O $M_r = 448.23$ Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

 $a = 14.6942 (2) \text{ \AA}$ $b = 6.1103 (1) \text{ \AA}$ $c = 21.5717 (4) \text{ \AA}$ $\beta = 113.714 (1)^\circ$ $V = 1773.30 (5) \text{ \AA}^3$ $Z = 4$ $F(000) = 888$ $D_x = 1.679 \text{ Mg m}^{-3}$ Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 23524 reflections

 $\theta = 2.6\text{--}26.7^\circ$ $\mu = 4.50 \text{ mm}^{-1}$ $T = 290 \text{ K}$

Plate, colourless

 $0.36 \times 0.19 \times 0.16 \text{ mm}$

Data collection

Nonius KappaCCD

diffractometer

Radiation source: sealed tube

Graphite monochromator

CCD rotation images scans

Absorption correction: multi-scan
(SADABS; Sheldrick, 1996) $T_{\min} = 0.291$, $T_{\max} = 0.734$

32063 measured reflections

3734 independent reflections

3177 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.076$ $\theta_{\max} = 26.7^\circ$, $\theta_{\min} = 3.6^\circ$ $h = -18 \rightarrow 18$ $k = -7 \rightarrow 7$ $l = -27 \rightarrow 25$

Refinement

Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.037$ $wR(F^2) = 0.095$ $S = 1.03$

3734 reflections

200 parameters

0 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0428P)^2 + 1.5141P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} < 0.001$ $\Delta\rho_{\max} = 0.80 \text{ e \AA}^{-3}$ $\Delta\rho_{\min} = -0.55 \text{ e \AA}^{-3}$

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.27316 (19)	0.2209 (5)	0.25810 (14)	0.0424 (6)
C2	0.36493 (19)	0.1572 (5)	0.32017 (13)	0.0419 (6)
H2	0.4114	0.0804	0.3055	0.050*
C3	0.5393 (2)	0.2846 (6)	0.42271 (17)	0.0593 (8)

H3A	0.5269	0.1896	0.4545	0.071*
H3B	0.5683	0.1973	0.3977	0.071*
C4	0.6102 (3)	0.4646 (8)	0.4605 (2)	0.0811 (12)
H4A	0.5806	0.5522	0.4844	0.122*
H4B	0.6243	0.5547	0.4290	0.122*
H4C	0.6709	0.4014	0.4922	0.122*
C5	0.23970 (19)	0.0756 (5)	0.19795 (14)	0.0409 (6)
C6	0.1669 (2)	0.1548 (5)	0.13798 (15)	0.0488 (6)
H6	0.1396	0.2925	0.1373	0.059*
C7	0.1349 (2)	0.0315 (5)	0.07957 (16)	0.0544 (7)
H7	0.0868	0.0854	0.0395	0.065*
C8	0.1756 (2)	-0.1734 (5)	0.08164 (15)	0.0503 (7)
C9	0.2473 (2)	-0.2565 (5)	0.14005 (16)	0.0506 (7)
H9	0.2736	-0.3952	0.1405	0.061*
C10	0.2796 (2)	-0.1304 (5)	0.19819 (15)	0.0470 (6)
H10	0.3287	-0.1843	0.2379	0.056*
C11	0.1967 (2)	0.0520 (5)	0.36178 (14)	0.0468 (6)
C12	0.1754 (3)	0.2435 (6)	0.38661 (18)	0.0612 (8)
H12	0.2261	0.3392	0.4116	0.073*
C13	0.0772 (3)	0.2921 (7)	0.3739 (2)	0.0709 (10)
H13	0.0618	0.4222	0.3900	0.085*
C14	0.0026 (3)	0.1487 (8)	0.33776 (19)	0.0722 (10)
H14	-0.0631	0.1809	0.3298	0.087*
C15	0.0248 (3)	-0.0399 (8)	0.3137 (2)	0.0713 (10)
H15	-0.0259	-0.1368	0.2894	0.086*
C16	0.1217 (2)	-0.0899 (6)	0.32472 (17)	0.0577 (8)
H16	0.1362	-0.2181	0.3073	0.069*
O1	0.22999 (15)	0.3895 (3)	0.25913 (11)	0.0532 (5)
O2	0.45208 (18)	0.5313 (4)	0.31552 (12)	0.0643 (6)
O1W	0.4391 (2)	0.4022 (4)	0.18722 (14)	0.0729 (7)
H1W	0.4651	0.2794	0.1849	0.088*
H2W	0.4411	0.4158	0.2263	0.088*
S	0.42388 (5)	0.40496 (12)	0.36491 (4)	0.04610 (18)
Se	0.33094 (2)	-0.03515 (6)	0.380920 (17)	0.05674 (12)
Br	0.13073 (3)	-0.34569 (7)	0.001765 (19)	0.07905 (15)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0385 (13)	0.0460 (15)	0.0428 (15)	-0.0001 (11)	0.0165 (11)	0.0043 (11)
C2	0.0366 (13)	0.0488 (15)	0.0393 (14)	0.0033 (11)	0.0143 (11)	-0.0005 (11)
C3	0.0458 (16)	0.077 (2)	0.0477 (17)	-0.0005 (15)	0.0116 (13)	-0.0088 (16)
C4	0.054 (2)	0.114 (3)	0.067 (2)	-0.015 (2)	0.0156 (18)	-0.030 (2)
C5	0.0351 (12)	0.0465 (14)	0.0402 (14)	-0.0005 (11)	0.0144 (11)	0.0018 (11)
C6	0.0438 (14)	0.0497 (16)	0.0474 (16)	0.0058 (12)	0.0127 (12)	0.0029 (12)
C7	0.0494 (16)	0.0604 (18)	0.0426 (16)	0.0017 (14)	0.0073 (13)	0.0028 (13)
C8	0.0491 (15)	0.0589 (18)	0.0440 (15)	-0.0071 (13)	0.0199 (13)	-0.0048 (13)
C9	0.0498 (15)	0.0490 (16)	0.0527 (17)	0.0008 (13)	0.0203 (13)	-0.0028 (13)

C10	0.0410 (14)	0.0505 (16)	0.0442 (15)	0.0017 (12)	0.0116 (12)	0.0036 (12)
C11	0.0484 (15)	0.0545 (16)	0.0403 (15)	-0.0053 (12)	0.0207 (12)	0.0044 (12)
C12	0.0638 (19)	0.062 (2)	0.0597 (19)	-0.0092 (16)	0.0271 (16)	-0.0088 (16)
C13	0.078 (2)	0.077 (2)	0.070 (2)	0.0102 (19)	0.043 (2)	-0.0010 (19)
C14	0.0531 (19)	0.109 (3)	0.061 (2)	0.003 (2)	0.0298 (17)	0.014 (2)
C15	0.0543 (19)	0.099 (3)	0.062 (2)	-0.0205 (19)	0.0248 (17)	-0.007 (2)
C16	0.0591 (18)	0.0643 (19)	0.0530 (18)	-0.0161 (15)	0.0260 (15)	-0.0090 (15)
O1	0.0507 (11)	0.0509 (11)	0.0526 (12)	0.0099 (9)	0.0149 (9)	-0.0012 (9)
O2	0.0661 (14)	0.0659 (14)	0.0611 (14)	-0.0174 (11)	0.0258 (12)	0.0026 (11)
O1W	0.0846 (17)	0.0675 (15)	0.0766 (17)	0.0192 (13)	0.0428 (14)	0.0114 (13)
S	0.0449 (4)	0.0504 (4)	0.0427 (4)	-0.0023 (3)	0.0172 (3)	-0.0049 (3)
Se	0.05056 (19)	0.0595 (2)	0.0579 (2)	0.00582 (13)	0.01947 (15)	0.01869 (14)
Br	0.0929 (3)	0.0816 (3)	0.0526 (2)	-0.0047 (2)	0.01875 (19)	-0.02037 (18)

Geometric parameters (Å, °)

C1—O1	1.215 (3)	C8—Br	1.897 (3)
C1—C5	1.483 (4)	C9—C10	1.383 (4)
C1—C2	1.520 (4)	C9—H9	0.9300
C2—S	1.817 (3)	C10—H10	0.9300
C2—Se	1.969 (3)	C11—C12	1.375 (5)
C2—H2	0.9800	C11—C16	1.377 (4)
C3—C4	1.509 (5)	C11—Se	1.920 (3)
C3—S	1.809 (3)	C12—C13	1.388 (5)
C3—H3A	0.9700	C12—H12	0.9300
C3—H3B	0.9700	C13—C14	1.375 (6)
C4—H4A	0.9600	C13—H13	0.9300
C4—H4B	0.9600	C14—C15	1.357 (6)
C4—H4C	0.9600	C14—H14	0.9300
C5—C10	1.388 (4)	C15—C16	1.380 (5)
C5—C6	1.392 (4)	C15—H15	0.9300
C6—C7	1.378 (4)	C16—H16	0.9300
C6—H6	0.9300	O2—S	1.503 (2)
C7—C8	1.380 (4)	O1W—H1W	0.8525
C7—H7	0.9300	O1W—H2W	0.8362
C8—C9	1.374 (4)		
O1—C1—C5	122.1 (2)	C9—C8—Br	119.1 (2)
O1—C1—C2	119.1 (3)	C7—C8—Br	119.1 (2)
C5—C1—C2	118.8 (2)	C8—C9—C10	118.8 (3)
C1—C2—S	108.62 (19)	C8—C9—H9	120.6
C1—C2—Se	111.50 (17)	C10—C9—H9	120.6
S—C2—Se	109.77 (14)	C9—C10—C5	120.8 (3)
C1—C2—H2	109.0	C9—C10—H10	119.6
S—C2—H2	109.0	C5—C10—H10	119.6
Se—C2—H2	109.0	C12—C11—C16	120.4 (3)
C4—C3—S	109.2 (3)	C12—C11—Se	121.9 (2)
C4—C3—H3A	109.8	C16—C11—Se	117.6 (2)

S—C3—H3A	109.8	C11—C12—C13	119.2 (3)
C4—C3—H3B	109.8	C11—C12—H12	120.4
S—C3—H3B	109.8	C13—C12—H12	120.4
H3A—C3—H3B	108.3	C14—C13—C12	120.2 (4)
C3—C4—H4A	109.5	C14—C13—H13	119.9
C3—C4—H4B	109.5	C12—C13—H13	119.9
H4A—C4—H4B	109.5	C15—C14—C13	119.9 (3)
C3—C4—H4C	109.5	C15—C14—H14	120.0
H4A—C4—H4C	109.5	C13—C14—H14	120.0
H4B—C4—H4C	109.5	C14—C15—C16	120.8 (3)
C10—C5—C6	118.9 (3)	C14—C15—H15	119.6
C10—C5—C1	123.3 (2)	C16—C15—H15	119.6
C6—C5—C1	117.7 (2)	C11—C16—C15	119.4 (3)
C7—C6—C5	120.8 (3)	C11—C16—H16	120.3
C7—C6—H6	119.6	C15—C16—H16	120.3
C5—C6—H6	119.6	H1W—O1W—H2W	108.1
C6—C7—C8	118.8 (3)	O2—S—C3	104.37 (15)
C6—C7—H7	120.6	O2—S—C2	105.07 (13)
C8—C7—H7	120.6	C3—S—C2	98.16 (14)
C9—C8—C7	121.8 (3)	C11—Se—C2	101.82 (11)
O1—C1—C2—S	-28.1 (3)	C16—C11—C12—C13	0.1 (5)
C5—C1—C2—S	151.4 (2)	Se—C11—C12—C13	-176.7 (3)
O1—C1—C2—Se	93.0 (3)	C11—C12—C13—C14	0.9 (5)
C5—C1—C2—Se	-87.5 (2)	C12—C13—C14—C15	-0.8 (6)
O1—C1—C5—C10	-171.6 (3)	C13—C14—C15—C16	-0.3 (6)
C2—C1—C5—C10	8.9 (4)	C12—C11—C16—C15	-1.1 (5)
O1—C1—C5—C6	10.6 (4)	Se—C11—C16—C15	175.8 (3)
C2—C1—C5—C6	-168.9 (2)	C14—C15—C16—C11	1.3 (6)
C10—C5—C6—C7	-0.2 (4)	C4—C3—S—O2	64.1 (3)
C1—C5—C6—C7	177.7 (3)	C4—C3—S—C2	172.0 (3)
C5—C6—C7—C8	0.7 (5)	C1—C2—S—O2	-61.3 (2)
C6—C7—C8—C9	-0.5 (5)	Se—C2—S—O2	176.53 (14)
C6—C7—C8—Br	179.2 (2)	C1—C2—S—C3	-168.7 (2)
C7—C8—C9—C10	-0.2 (5)	Se—C2—S—C3	69.18 (16)
Br—C8—C9—C10	-179.9 (2)	C12—C11—Se—C2	-76.2 (3)
C8—C9—C10—C5	0.7 (4)	C16—C11—Se—C2	106.9 (2)
C6—C5—C10—C9	-0.5 (4)	C1—C2—Se—C11	-27.7 (2)
C1—C5—C10—C9	-178.3 (3)	S—C2—Se—C11	92.67 (15)

Hydrogen-bond geometry (Å, °)

Cg1 and Cg2 are the centroids of the C5—C10 and C11—C16 rings, respectively.

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O1w—H1w \cdots O2 ⁱ	0.85	1.95	2.788 (4)	169
O1w—H2w \cdots O2	0.84	1.99	2.810 (4)	165
C2—H2 \cdots O1w ⁱ	0.98	2.40	3.334 (4)	159
C3—H3b \cdots O1w ⁱ	0.97	2.54	3.434 (4)	153

C9—H9···O1w ⁱⁱ	0.93	2.55	3.320 (4)	141
C10—H10···O2 ⁱⁱ	0.93	2.58	3.456 (4)	157
C14—H14···Cg1 ⁱⁱⁱ	0.93	2.96	3.793 (5)	149
C8—Br···Cg2 ^{iv}	1.90 (1)	3.49 (1)	5.349 (3)	165 (1)

Symmetry codes: (i) $-x+1, y-1/2, -z+1/2$; (ii) $x, y-1, z$; (iii) $-x, y+1/2, -z+1/2$; (iv) $x, -y-1/2, z-1/2$.