

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

# 7-Diethylamino-2-oxo-2H-chromene-3-carbaldehyde

Hong-Da Li and Bing-Zhu Yin\*

Key Laboratory of Natural Resources of Changbai Mountain &amp; Functional Molecules (Yanbian University), Ministry of Education, Yanji 133002, People's Republic of China

Correspondence e-mail: zqcong@ybu.edu.cn


Received 8 June 2011; accepted 14 June 2011

 Key indicators: single-crystal X-ray study;  $T = 295$  K; mean  $\sigma(\text{C}-\text{C}) = 0.003$  Å; disorder in main residue;  $R$  factor = 0.048;  $wR$  factor = 0.144; data-to-parameter ratio = 15.5.

In the title compound,  $\text{C}_{14}\text{H}_{15}\text{NO}_3$ , all non-H atoms except for those of the methyl and the disordered ethyl group are approximately co-planar, the largest deviation from the mean plane being 0.0223 (13) Å at the N atom. In the crystal, the packing of molecules through weak intermolecular  $\text{C}-\text{H}\cdots\text{O}$  hydrogen-bonding interactions leads to the formation of layers parallel to  $bc$  plane. Within these layers, there exist slipped  $\pi-\pi$  stacking interactions between symmetry-related fused rings [centroid-centroid distances = 3.527 (3) and 3.554 (3), slippage = 0.988 and 1.011 Å, respectively]. One ethyl group is disordered over two sets of sites with site-occupation factors of 0.54 and 0.46.

## Related literature

For background to the title compound, an organic intermediate and a fluorescent probe for cyanide and amino acids, see: Kim *et al.* (2010). For electronic and photonic applications of coumarins, see: Murray *et al.* (1982). For the synthesis, see: Wu *et al.* (2007).



## Experimental

### Crystal data

$\text{C}_{14}\text{H}_{15}\text{NO}_3$	$V = 2517$ (3) Å <sup>3</sup>
$M_r = 245.27$	$Z = 8$
Monoclinic, $C2/c$	Mo $K\alpha$ radiation
$a = 25.488$ (17) Å	$\mu = 0.09$ mm <sup>-1</sup>
$b = 7.844$ (6) Å	$T = 295$ K
$c = 12.599$ (12) Å	$0.41 \times 0.39 \times 0.21$ mm
$\beta = 92.39$ (3)°	

### Data collection

Rigaku R-Axis RAPID diffractometer	11514 measured reflections
Absorption correction: multi-scan (ABSCOR; Higashi, 1995)	2850 independent reflections
$T_{\min} = 0.963$ , $T_{\max} = 0.981$	1724 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.033$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.048$	1 restraint
$wR(F^2) = 0.144$	H-atom parameters constrained
$S = 1.06$	$\Delta\rho_{\text{max}} = 0.14$ e Å <sup>-3</sup>
2850 reflections	$\Delta\rho_{\text{min}} = -0.15$ e Å <sup>-3</sup>
184 parameters	

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C8}-\text{H8}\cdots\text{O3}^{\text{i}}$	0.93	2.58	3.367 (4)	143
$\text{C9}-\text{H9}\cdots\text{O1}^{\text{ii}}$	0.93	2.55	3.432 (3)	158
$\text{C13}-\text{H13B}\cdots\text{O2}^{\text{iii}}$	0.97	2.53	3.388 (3)	147

 Symmetry codes: (i)  $-x + \frac{1}{2}, y + \frac{1}{2}, -z + \frac{5}{2}$ ; (ii)  $x, -y, z + \frac{1}{2}$ ; (iii)  $-x + \frac{1}{2}, y + \frac{1}{2}, -z + \frac{3}{2}$ .

Data collection: *RAPID-AUTO* (Rigaku, 1998); cell refinement: *RAPID-AUTO*; data reduction: *CrystalStructure* (Rigaku, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEPIII* (Burnett & Johnson, 1996) and *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97*.

The authors acknowledge financial support from the National Natural Science Foundation of China (grant No. 21062022) and the Open Project of the State Key Laboratory of Supramolecular Structure and Materials, Jilin University.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: DN2698).

## References

- Burnett, M. N. & Johnson, C. K. (1996). *ORTEPIII*. Report ORNL-6895. Oak Ridge National Laboratory, Tennessee, USA.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
- Kim, G. J. & Kim, H. J. (2010). *Tetrahedron Lett.* **51**, 2914–2916.
- Murray, R. D., Mendez, J. & Brown, S. A. (1982). *The Natural Coumarins: Occurrence, Chemistry and Biochemistry*, p. 227. New York: John Wiley and Sons.
- Rigaku (1998). *RAPID-AUTO*. Rigaku Corporation, Tokyo, Japan.
- Rigaku (2002). *CrystalStructure*. Rigaku/MSI Inc., The Woodlands, Texas, USA.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Wu, J. S., Liu, W. M., Zhuang, X. Q., Wang, F., Wang, P. F., Tao, S. L., Zhang, X. H., Wu, S. K. & Lee, S. T. (2007). *Org. Lett.* **9**, 33–36.

## supporting information

*Acta Cryst.* (2011). E67, o1713 [doi:10.1107/S160053681102294X]

**7-Diethylamino-2-oxo-2H-chromene-3-carbaldehyde****Hong-Da Li and Bing-Zhu Yin****S1. Comment**

Coumarins are an important class of naturally occurring and synthetic compounds, which have been extensively investigated for electronic and photonic applications (Murray *et al.*, 1982). Herein, we reported the crystal structure of the title compound, an important organic intermediate and a fluorescent probe for cyanide and amino acids (Kim *et al.*, 2010).

The molecule of title compound formed by two fused rings is mainly planar with the exception of the methyl and disordered ethyl group (Fig. 1). The weak intermolecular C—H $\cdots$ O hydrogen bonds (Table 1) link the molecules into layers parallel to the (100) plane. Furthermore, slipped  $\pi$ - $\pi$  stacking occurs between symmetry related fused rings within the layers (Table 2)


**S2. Experimental**

The title compound was prepared according to the literature (Wu *et al.*, 2007). Single crystals suitable for X-ray diffraction were prepared by recrystallization from mixture of dichloromethane and petroleum (60–90 °C).

**S3. Refinement**

Carbon-bound H-atoms were placed in calculated positions (C—H 0.93 to 0.97 Å) and were included in the refinement in the riding model with  $U_{\text{iso}}(\text{H}) = 1.2$  or  $1.5 U_{\text{eq}}(\text{C}_{\text{methyl}})$ .

One of the ethyl group is disordered over two positions with a site occupancy in the ratio 0.54/0.46. The refinement of the disordered moieties was carried out using the PART instruction and restraining them to have identical geometry with the SAME instruction available in SHELXL-97 (Sheldrick, 2008)

**Figure 1**

The crystal structure of the title compound, with the atom numbering scheme. Displacement ellipsoids of non-H atoms are drawn at the 30% probability level. H atoms are shown as small spheres of arbitrary radii. Only the major component of the disordered ethyl is represented for clarity.

### 7-Diethylamino-2-oxo-2H-chromene-3-carbaldehyde

#### Crystal data

$C_{14}H_{15}NO_3$

$M_r = 245.27$

Monoclinic,  $C2/c$

Hall symbol:  $-C 2yc$

$a = 25.488 (17) \text{ \AA}$

$b = 7.844 (6) \text{ \AA}$

$c = 12.599 (12) \text{ \AA}$

$\beta = 92.39 (3)^\circ$

$V = 2517 (3) \text{ \AA}^3$

$Z = 8$

$F(000) = 1040$

$D_x = 1.295 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 6542 reflections

$\theta = 3.2\text{--}27.5^\circ$

$\mu = 0.09 \text{ mm}^{-1}$

$T = 295 \text{ K}$

Block, brown

$0.41 \times 0.39 \times 0.21 \text{ mm}$

#### Data collection

Rigaku R-AXIS RAPID

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

$\omega$  scans

Absorption correction: multi-scan

(*ABSCOR*; Higashi, 1995)

$T_{\min} = 0.963$ ,  $T_{\max} = 0.981$

11514 measured reflections

2850 independent reflections

1724 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.033$

$\theta_{\max} = 27.5^\circ$ ,  $\theta_{\min} = 3.2^\circ$

$h = -32 \rightarrow 32$

$k = -9 \rightarrow 10$

$l = -16 \rightarrow 16$

*Refinement*

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.048$   
 $wR(F^2) = 0.144$   
 $S = 1.06$   
 2850 reflections  
 184 parameters  
 1 restraint  
 Primary atom site location: structure-invariant  
 direct methods

Secondary atom site location: difference Fourier  
 map  
 Hydrogen site location: inferred from  
 neighbouring sites  
 H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0674P)^2 + 0.3544P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} < 0.001$   
 $\Delta\rho_{\max} = 0.14 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.15 \text{ e } \text{\AA}^{-3}$

*Special details*

**Experimental.** (See detailed section in the paper)

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
O1	0.24299 (4)	0.00447 (14)	0.88750 (8)	0.0556 (3)	
O2	0.17217 (5)	-0.12688 (18)	0.82505 (10)	0.0787 (4)	
O3	0.12022 (6)	-0.24463 (18)	1.11710 (12)	0.0874 (5)	
N1	0.40226 (6)	0.2780 (2)	0.99586 (12)	0.0747 (5)	
C1	0.19772 (6)	-0.0872 (2)	0.90386 (13)	0.0562 (4)	
C2	0.18575 (6)	-0.12332 (19)	1.01323 (12)	0.0533 (4)	
C3	0.21890 (6)	-0.0707 (2)	1.09396 (12)	0.0546 (4)	
H3	0.2105	-0.0940	1.1636	0.066*	
C4	0.26540 (6)	0.01794 (19)	1.07523 (11)	0.0503 (4)	
C5	0.27674 (6)	0.05510 (19)	0.96975 (11)	0.0478 (4)	
C6	0.32073 (6)	0.1413 (2)	0.94200 (12)	0.0533 (4)	
H6	0.3261	0.1652	0.8710	0.064*	
C7	0.35790 (6)	0.1938 (2)	1.02108 (13)	0.0578 (4)	
C8	0.34695 (7)	0.1572 (2)	1.12853 (13)	0.0636 (5)	
H8	0.3707	0.1906	1.1826	0.076*	
C9	0.30257 (7)	0.0747 (2)	1.15318 (12)	0.0613 (4)	
H9	0.2963	0.0546	1.2243	0.074*	
C10	0.13764 (7)	-0.2172 (2)	1.03121 (16)	0.0679 (5)	
H10	0.1189	-0.2592	0.9719	0.081*	
C11A	0.4515 (3)	0.2676 (10)	1.0761 (7)	0.086 (2)	0.46
H11A	0.4478	0.1756	1.1267	0.103*	0.46
H11B	0.4836	0.2509	1.0387	0.103*	0.46
C12A	0.4513 (2)	0.4378 (8)	1.1303 (5)	0.1030 (17)	0.46
H12A	0.4539	0.5266	1.0784	0.154*	0.46

H12B	0.4807	0.4448	1.1805	0.154*	0.46
H12C	0.4193	0.4510	1.1669	0.154*	0.46
C11B	0.4390 (2)	0.3468 (8)	1.0753 (5)	0.0750 (15)	0.54
H11C	0.4204	0.3835	1.1370	0.090*	0.54
H11D	0.4570	0.4445	1.0469	0.090*	0.54
C12B	0.4784 (2)	0.2088 (8)	1.1068 (5)	0.125 (2)	0.54
H12D	0.4602	0.1130	1.1354	0.188*	0.54
H12E	0.5031	0.2527	1.1595	0.188*	0.54
H12F	0.4966	0.1734	1.0455	0.188*	0.54
C13	0.41299 (8)	0.3169 (3)	0.88577 (15)	0.0742 (5)	
H13A	0.4374	0.4116	0.8847	0.089*	
H13B	0.3806	0.3533	0.8495	0.089*	
C14	0.43552 (9)	0.1696 (3)	0.8251 (2)	0.1025 (8)	
H14A	0.4692	0.1394	0.8563	0.154*	
H14B	0.4393	0.2024	0.7524	0.154*	
H14C	0.4123	0.0735	0.8280	0.154*	

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O1	0.0602 (6)	0.0714 (7)	0.0356 (6)	−0.0004 (6)	0.0062 (4)	0.0006 (5)
O2	0.0830 (8)	0.1019 (10)	0.0503 (7)	−0.0158 (7)	−0.0056 (6)	0.0026 (7)
O3	0.0943 (10)	0.0941 (10)	0.0756 (9)	−0.0150 (8)	0.0259 (8)	0.0101 (7)
N1	0.0630 (8)	0.1053 (12)	0.0559 (9)	−0.0141 (8)	0.0021 (7)	0.0042 (8)
C1	0.0627 (9)	0.0599 (10)	0.0459 (9)	0.0078 (8)	0.0030 (7)	0.0007 (7)
C2	0.0613 (9)	0.0508 (9)	0.0486 (9)	0.0078 (8)	0.0114 (7)	0.0042 (7)
C3	0.0697 (10)	0.0563 (9)	0.0387 (8)	0.0110 (8)	0.0125 (7)	0.0036 (7)
C4	0.0612 (9)	0.0544 (9)	0.0357 (8)	0.0106 (7)	0.0091 (6)	0.0020 (6)
C5	0.0565 (8)	0.0530 (9)	0.0342 (7)	0.0129 (7)	0.0044 (6)	−0.0019 (6)
C6	0.0602 (9)	0.0646 (10)	0.0358 (8)	0.0082 (8)	0.0089 (6)	0.0018 (7)
C7	0.0604 (9)	0.0651 (10)	0.0479 (9)	0.0078 (8)	0.0044 (7)	0.0016 (7)
C8	0.0707 (11)	0.0787 (12)	0.0408 (9)	0.0001 (9)	−0.0031 (7)	0.0005 (8)
C9	0.0784 (11)	0.0727 (11)	0.0329 (8)	0.0061 (9)	0.0046 (7)	0.0034 (7)
C10	0.0754 (11)	0.0657 (11)	0.0631 (12)	0.0018 (9)	0.0103 (9)	0.0037 (9)
C11A	0.069 (4)	0.099 (6)	0.091 (4)	0.013 (4)	0.007 (3)	0.022 (4)
C12A	0.093 (4)	0.121 (5)	0.094 (4)	−0.022 (3)	−0.009 (3)	−0.008 (4)
C11B	0.065 (3)	0.086 (4)	0.073 (3)	−0.015 (3)	−0.013 (2)	0.005 (3)
C12B	0.083 (3)	0.152 (5)	0.137 (5)	0.036 (3)	−0.040 (3)	0.021 (4)
C13	0.0713 (11)	0.0878 (13)	0.0643 (11)	−0.0101 (10)	0.0123 (9)	0.0046 (10)
C14	0.0940 (15)	0.1172 (19)	0.0991 (18)	0.0080 (14)	0.0368 (13)	−0.0051 (15)

*Geometric parameters (Å, °)*

O1—C5	1.3774 (19)	C9—H9	0.9300
O1—C1	1.382 (2)	C10—H10	0.9300
O2—C1	1.205 (2)	C11A—C12A	1.500 (8)
O3—C10	1.206 (2)	C11A—H11A	0.9700
N1—C7	1.358 (2)	C11A—H11B	0.9700

N1—C11B	1.446 (6)	C12A—H12A	0.9600
N1—C13	1.457 (3)	C12A—H12B	0.9600
N1—C11A	1.581 (8)	C12A—H12C	0.9600
C1—C2	1.452 (3)	C11B—C12B	1.517 (7)
C2—C3	1.359 (2)	C11B—H11C	0.9700
C2—C10	1.456 (3)	C11B—H11D	0.9700
C3—C4	1.402 (2)	C12B—H12D	0.9600
C3—H3	0.9300	C12B—H12E	0.9600
C4—C5	1.402 (2)	C12B—H12F	0.9600
C4—C9	1.408 (2)	C13—C14	1.512 (3)
C5—C6	1.367 (2)	C13—H13A	0.9700
C6—C7	1.408 (2)	C13—H13B	0.9700
C6—H6	0.9300	C14—H14A	0.9600
C7—C8	1.423 (2)	C14—H14B	0.9600
C8—C9	1.351 (2)	C14—H14C	0.9600
C8—H8	0.9300		
C5—O1—C1	122.47 (12)	O3—C10—C2	125.00 (19)
C7—N1—C11B	122.7 (3)	O3—C10—H10	117.5
C7—N1—C13	121.03 (15)	C2—C10—H10	117.5
C11B—N1—C13	116.1 (3)	C12A—C11A—N1	103.2 (5)
C7—N1—C11A	118.2 (3)	C12A—C11A—H11A	111.1
C13—N1—C11A	116.3 (3)	N1—C11A—H11A	111.1
O2—C1—O1	115.93 (15)	C12A—C11A—H11B	111.1
O2—C1—C2	127.14 (17)	N1—C11A—H11B	111.1
O1—C1—C2	116.93 (14)	H11A—C11A—H11B	109.1
C3—C2—C1	120.15 (16)	N1—C11B—C12B	108.5 (5)
C3—C2—C10	122.59 (16)	N1—C11B—H11C	110.0
C1—C2—C10	117.26 (16)	C12B—C11B—H11C	110.0
C2—C3—C4	121.88 (14)	N1—C11B—H11D	110.0
C2—C3—H3	119.1	C12B—C11B—H11D	110.0
C4—C3—H3	119.1	H11C—C11B—H11D	108.4
C3—C4—C5	118.13 (14)	C11B—C12B—H12D	109.5
C3—C4—C9	126.03 (14)	C11B—C12B—H12E	109.5
C5—C4—C9	115.84 (15)	H12D—C12B—H12E	109.5
C6—C5—O1	116.32 (13)	C11B—C12B—H12F	109.5
C6—C5—C4	123.28 (14)	H12D—C12B—H12F	109.5
O1—C5—C4	120.40 (15)	H12E—C12B—H12F	109.5
C5—C6—C7	119.92 (15)	N1—C13—C14	114.33 (19)
C5—C6—H6	120.0	N1—C13—H13A	108.7
C7—C6—H6	120.0	C14—C13—H13A	108.7
N1—C7—C6	121.26 (15)	N1—C13—H13B	108.7
N1—C7—C8	121.26 (16)	C14—C13—H13B	108.7
C6—C7—C8	117.48 (16)	H13A—C13—H13B	107.6
C9—C8—C7	120.99 (16)	C13—C14—H14A	109.5
C9—C8—H8	119.5	C13—C14—H14B	109.5
C7—C8—H8	119.5	H14A—C14—H14B	109.5
C8—C9—C4	122.47 (15)	C13—C14—H14C	109.5

C8—C9—H9	118.8	H14A—C14—H14C	109.5
C4—C9—H9	118.8	H14B—C14—H14C	109.5

*Hydrogen-bond geometry (Å, °)*

<i>D—H···A</i>	<i>D—H</i>	<i>H···A</i>	<i>D···A</i>	<i>D—H···A</i>
C8—H8···O3 <sup>i</sup>	0.93	2.58	3.367 (4)	143
C9—H9···O1 <sup>ii</sup>	0.93	2.55	3.432 (3)	158
C13—H13B···O2 <sup>iii</sup>	0.97	2.53	3.388 (3)	147

Symmetry codes: (i)  $-x+1/2, y+1/2, -z+5/2$ ; (ii)  $x, -y, z+1/2$ ; (iii)  $-x+1/2, y+1/2, -z+3/2$ .