

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Sodium piperidine-1-carbodithioate dihydrate

Ana C. Mafud* and Maria Teresa P. Gambardella

Instituto de Química de São Carlos, Universidade de São Paulo, Av. Trabalhador São-carlense 400, Caixa Postal 780, 13560-970, São Carlos, SP, Brazil
Correspondence e-mail: mafud@iqsc.usp.br


Received 8 June 2011; accepted 10 June 2011

Key indicators: single-crystal X-ray study; $T = 290$ K; mean $\sigma(\text{S}-\text{Na}) = 0.003$ Å; disorder in main residue; R factor = 0.054; wR factor = 0.152; data-to-parameter ratio = 10.6.

The asymmetric unit of the title compound, $\text{Na}^+ \cdot \text{C}_6\text{H}_{10}\text{NS}_2 \cdot 2\text{H}_2\text{O}$, is composed of a sodium cation, a piperidinedithiocarbamate anion which exhibits positional disorder, and two lattice water molecules. The atoms of the piperidine ring are divided over two sites with occupancy factors of 0.554 (6) and 0.446 (6). In the crystal, the sodium cation (coordination number of 6) and the piperidinedithiocarbamate anion are linked, forming an infinite two-dimensional network extending parallel to (001). $\text{O}-\text{H} \cdots \text{S}$ hydrogen bonds, involving the lattice water molecules, also aid in stabilizing the crystal structure.

Related literature

For the crystal structures of similar compounds, see: Oskarsson *et al.* (1979); Albertsson *et al.* (1980); Ymén (1982); Mafud & Gambardella (2011). For puckering parameters, see: Cremer & Pople (1975).



Experimental

Crystal data

$\text{Na}^+ \cdot \text{C}_6\text{H}_{10}\text{NS}_2 \cdot 2\text{H}_2\text{O}$
 $M_r = 219.29$
Monoclinic, $P2_1/a$
 $a = 12.241$ (5) Å
 $b = 5.909$ (5) Å

$c = 14.690$ (5) Å
 $\beta = 95.519$ (5)°
 $V = 1057.6$ (11) Å³
 $Z = 4$
Mo $K\alpha$ radiation

$\mu = 0.51$ mm⁻¹
 $T = 290$ K

0.02 × 0.02 × 0.02 mm

Data collection

Nonius KappaCCD diffractometer
7553 measured reflections
1863 independent reflections

1482 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.145$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.054$
 $wR(F^2) = 0.152$
 $S = 1.02$
1863 reflections
176 parameters
6 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.33$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.36$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
$\text{O1}-\text{H11O} \cdots \text{S2}^{\text{i}}$	0.84 (2)	2.39 (2)	3.214 (2)	167 (3)
$\text{O1}-\text{H12O} \cdots \text{S1}^{\text{ii}}$	0.85 (2)	2.49 (2)	3.322 (3)	167 (3)
$\text{O2}-\text{H21O} \cdots \text{S2}^{\text{ii}}$	0.83 (2)	2.48 (2)	3.283 (2)	163 (3)
$\text{O2}-\text{H22O} \cdots \text{S1}^{\text{iii}}$	0.86 (2)	2.46 (2)	3.313 (2)	174 (3)

Symmetry codes: (i) $x - \frac{1}{2}, -y + \frac{1}{2}, z$; (ii) $x - \frac{1}{2}, -y - \frac{1}{2}, z$; (iii) $x, y - 1, z$.

Data collection: *COLLECT* (Nonius, 1998); cell refinement: *SCALEPACK* (Otwinowski & Minor, 1997); data reduction: *DENZO* (Otwinowski & Minor, 1997) and *SCALEPACK*; program(s) used to solve structure: *SIR92* (Altomare *et al.*, 1994); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *Mercury* (Macrae *et al.*, 2006); software used to prepare material for publication: *WinGX* (Farrugia, 1999) and *pubCIF* (Westrip, 2010).

We are grateful to the Instituto de Química de São Carlos and the Universidade de São Paulo for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SU2283).

References

- Albertsson, J., Oskarsson, Å., Ståhl, K., Svensson, C. & Ymén, I. (1980). *Acta Cryst.* **B36**, 3072–3078.
Altomare, A., Casciarano, G., Giacovazzo, C., Guagliardi, A., Burla, M. C., Polidori, G. & Camalli, M. (1994). *J. Appl. Cryst.* **27**, 435.
Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
Mafud, A. C. & Gambardella, M. T. P. (2011). *Acta Cryst.* **E67**, o879.
Nonius (1998). *COLLECT*. Nonius BV, Delft, The Netherlands.
Oskarsson, A., Stahl, K., Svensson, C. & Ymén, I. (1979). *Eur. Cryst. Meet.* **5**, 67.
Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.
Ymén, I. (1982). *Acta Cryst.* **B38**, 2671–2674.

supporting information

Acta Cryst. (2011). E67, m942 [doi:10.1107/S1600536811022604]

Sodium piperidine-1-carbodithioate dihydrate

Ana C. Mafud and Maria Teresa P. Gambardella

S1. Comment

The title compound is composed of a piperidinedithiocarbamate anion, a sodium cation, and two lattice water molecules (Fig. 1). The crystal structures of similar compounds, for example sodium 1-pyrrolidinecarbodithioate dihydrate, has been reported on previously (Oskarsson *et al.*, 1979; Albertsson *et al.*, 1980; Ymén, 1982). The crystal structure of ammonium piperidine-1-carbodithioate dihydrate has been described by our group recently (Mafud & Gambardella, 2011).

The atoms of the piperidine ring are disordered, occupying two positions (A = C2',C3',C4',C5',N1' and B = C2,C3,C4,C5,N1) with occupancies of 0.554 (6)/0.446 (6). Both of these six-membered rings have a chair conformation, with puckering parameters, $Q = 0.552$ (13) Å, $\theta = 180.0$ (13) °, $\varphi_2 = 128$ (25) °, for ring A, and $Q = 0.577$ (15) Å, $\theta = 0.0$ (15) °, $\varphi_2 = 313$ (27) °, for ring B (Cremer & Pople, 1975).

The sodium atoms are coordinated to two sulfur atoms [Na1...S1 3.0649 (15) Å and Na1...S1ⁱ 2.9644 (15) Å; symmetry code: (i) $-x+3/2, y-1/2, -z+1$] and four oxygens [Na1...O2 2.360 (3) Å, Na1...O1 2.385 (2) Å, Na1...O1ⁱⁱ 2.416 (3) Å, Na1...O2ⁱⁱⁱ 2.515 (2) Å; symmetry codes: (ii) $-x+1, -y, -z+1$; (iii) $-x+1, -y-1, -z+1$], with a bi-pyramidal reversed geometry. This configuration generates close packed layers which remain cohesive in crystal stacking by van der Waals interactions. The distances of these contacts are slightly less than the sum of the van der Waals radii.

In the crystal O-H...S hydrogen bonds, involving the lattice water molecules, aid in stabilizing the crystal structure (Table 1). The crystal packing gives rise to a supramolecular structure, whose infinite two-dimensional network, with base vectors: #1 = [0 1 0], #2 = [1 0 0], grows parallel to (001), as shown in Fig. 2.

S2. Experimental

The title compound was prepared by slow addition of 0.1 mol of CS₂ to a cold solution containing 0.2 mol of piperidine and a stoichiometric amount of sodium hydroxide in ethanol/water 1:1 (v/v). The reaction mixture was placed in the freezer for 12 h and then filtered through a Büchner funnel, washed with cold ether and the product recrystallized in an ethanol water mixture 1:1 (v/v). Colourless single crystals, suitable for X-ray diffraction analysis, were obtained. On heating they sublimed and decomposed.

S3. Refinement

The H-atom positions of the water molecules were located in a difference Fourier map, they were refined with distance restraints, O-H = 0.84 (2) Å, with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{parent O-atom})$. The C-bound H-atoms of the anion were included in calculated positions and treated as riding atoms: C—H = 0.97 Å, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{parent C-atom})$.




Figure 1

Perspective view of asymmetric unit of the molecular structure of the title compound, with the numbering scheme and displacement ellipsoids drawn at the 50% probability level. The two components (A = C2',C3',C4',C5',N1' and B = C2,C3,C4,C5,N1) of the disordered piperidine ring are shown.




Figure 2

A view along the *b*-axis of the crystal packing of the title compound. Only the minor (B) component of the disordered piperidine ring is shown. The O-H...S hydrogen bonds are shown as dashed cyan lines and the C-bound H atoms have been omitted for clarity.

Sodium piperidine-1-carbodithioate dihydrate

Crystal data

$\text{Na}^+ \cdot \text{C}_6\text{H}_{10}\text{NS}_2 \cdot 2\text{H}_2\text{O}$

$M_r = 219.29$

Monoclinic, $P2_1/a$

Hall symbol: $-P\ 2yab$

$a = 12.241\ (5)\ \text{\AA}$

$b = 5.909\ (5)\ \text{\AA}$

$c = 14.690\ (5)\ \text{\AA}$

$\beta = 95.519\ (5)^\circ$

$V = 1057.6\ (11)\ \text{\AA}^3$

$Z = 4$

$F(000) = 464$

$D_x = 1.377\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 24030 reflections

$\theta = 2.9\text{--}26.7^\circ$

$\mu = 0.51\ \text{mm}^{-1}$

$T = 290\ \text{K}$

Prism, colourless

$0.02 \times 0.02 \times 0.02\ \text{mm}$

Data collection

Nonius KappaCCD

diffractometer

Radiation source: Enraf Nonius FR590

Graphite monochromator

Detector resolution: $9\ \text{pixels mm}^{-1}$

CCD rotation images, thick slices scans

7553 measured reflections

1863 independent reflections

1482 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.145$

$\theta_{\text{max}} = 25.1^\circ$, $\theta_{\text{min}} = 3.3^\circ$

$h = -14 \rightarrow 14$

$k = -7 \rightarrow 6$

$l = -17 \rightarrow 16$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.054$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.152$	$w = 1/[\sigma^2(F_o^2) + (0.0964P)^2]$
$S = 1.02$	where $P = (F_o^2 + 2F_c^2)/3$
1863 reflections	$(\Delta/\sigma)_{\max} < 0.001$
176 parameters	$\Delta\rho_{\max} = 0.33 \text{ e } \text{\AA}^{-3}$
6 restraints	$\Delta\rho_{\min} = -0.36 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
S1	0.74854 (5)	0.03097 (13)	0.39216 (4)	0.0481 (3)	
S2	0.86593 (6)	0.24242 (13)	0.24400 (5)	0.0577 (3)	
Na1	0.57813 (8)	-0.24210 (16)	0.48568 (8)	0.0509 (4)	
O1	0.41274 (14)	-0.0826 (4)	0.41396 (13)	0.0542 (5)	
H11O	0.403 (3)	-0.012 (5)	0.3644 (17)	0.081*	
H12O	0.369 (2)	-0.194 (4)	0.418 (2)	0.081*	
O2	0.55132 (14)	-0.5918 (4)	0.40928 (13)	0.0533 (5)	
H21O	0.509 (2)	-0.604 (6)	0.3618 (16)	0.08*	
H22O	0.604 (2)	-0.687 (5)	0.409 (2)	0.08*	
C1	0.7694 (2)	0.0599 (5)	0.27824 (18)	0.0551 (7)	
N1	0.6797 (6)	0.0084 (12)	0.2149 (4)	0.0541 (16)	0.446 (6)
C2	0.6746 (8)	0.0673 (16)	0.1177 (5)	0.081 (3)	0.446 (6)
H2A	0.73	0.1807	0.1092	0.097*	0.446 (6)
H2B	0.6034	0.1336	0.0991	0.097*	0.446 (6)
C3	0.6917 (19)	-0.128 (5)	0.0585 (18)	0.095 (8)	0.446 (6)
H3A	0.7657	-0.1862	0.0716	0.114*	0.446 (6)
H3B	0.6818	-0.0832	-0.0053	0.114*	0.446 (6)
C4	0.6069 (10)	-0.314 (3)	0.0773 (10)	0.079 (4)	0.446 (6)
H4A	0.5339	-0.2616	0.0555	0.095*	0.446 (6)
H4B	0.6218	-0.449	0.043	0.095*	0.446 (6)
C5	0.6089 (7)	-0.3753 (16)	0.1787 (6)	0.076 (2)	0.446 (6)
H5A	0.5505	-0.4818	0.1877	0.091*	0.446 (6)
H5B	0.6784	-0.4451	0.1998	0.091*	0.446 (6)
C6	0.5931 (7)	-0.1592 (18)	0.2320 (6)	0.067 (2)	0.446 (6)

H6A	0.5213	-0.0958	0.2135	0.08*	0.446 (6)
H6B	0.5969	-0.1932	0.2969	0.08*	0.446 (6)
N1'	0.7338 (4)	-0.1085 (10)	0.2187 (3)	0.0516 (13)	0.554 (6)
C2'	0.7582 (5)	-0.1213 (13)	0.1240 (4)	0.0677 (19)	0.554 (6)
H2'1	0.7906	-0.267	0.1125	0.081*	0.554 (6)
H2'2	0.8103	-0.0042	0.1119	0.081*	0.554 (6)
C3'	0.6545 (14)	-0.091 (5)	0.0626 (11)	0.095 (6)	0.554 (6)
H3'1	0.6717	-0.1086	-0.0002	0.115*	0.554 (6)
H3'2	0.6291	0.0629	0.0693	0.115*	0.554 (6)
C4'	0.5616 (9)	-0.249 (3)	0.0778 (10)	0.111 (5)	0.554 (6)
H4'1	0.5798	-0.4026	0.0616	0.133*	0.554 (6)
H4'2	0.4954	-0.2037	0.0407	0.133*	0.554 (6)
C5'	0.5447 (6)	-0.2350 (14)	0.1783 (6)	0.082 (2)	0.554 (6)
H5'1	0.5157	-0.0869	0.1912	0.098*	0.554 (6)
H5'2	0.4909	-0.3471	0.1922	0.098*	0.554 (6)
C6'	0.6494 (6)	-0.2736 (13)	0.2392 (4)	0.0629 (18)	0.554 (6)
H6'1	0.6349	-0.2592	0.3027	0.075*	0.554 (6)
H6'2	0.6759	-0.4258	0.2299	0.075*	0.554 (6)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0420 (4)	0.0551 (5)	0.0476 (4)	0.0008 (3)	0.0057 (3)	-0.0027 (3)
S2	0.0586 (5)	0.0585 (6)	0.0565 (5)	-0.0121 (3)	0.0090 (3)	0.0002 (3)
Na1	0.0465 (6)	0.0453 (8)	0.0609 (7)	0.0002 (4)	0.0053 (5)	-0.0005 (5)
O1	0.0534 (11)	0.0470 (13)	0.0608 (12)	-0.0039 (9)	-0.0015 (9)	0.0059 (10)
O2	0.0502 (10)	0.0497 (13)	0.0597 (12)	0.0060 (9)	0.0033 (8)	-0.0031 (10)
C1	0.0574 (15)	0.056 (2)	0.0512 (15)	-0.0111 (13)	0.0027 (12)	-0.0035 (14)
N1	0.062 (4)	0.052 (4)	0.047 (3)	-0.008 (3)	-0.001 (3)	0.000 (2)
C2	0.108 (6)	0.080 (7)	0.051 (4)	-0.026 (5)	-0.008 (4)	0.007 (4)
C3	0.104 (14)	0.107 (13)	0.079 (10)	-0.051 (11)	0.044 (10)	-0.030 (8)
C4	0.077 (8)	0.090 (9)	0.070 (6)	-0.033 (7)	0.003 (6)	-0.028 (7)
C5	0.070 (5)	0.070 (6)	0.088 (6)	-0.017 (5)	0.014 (4)	0.000 (5)
C6	0.053 (4)	0.086 (7)	0.061 (4)	-0.022 (5)	0.010 (4)	-0.010 (5)
N1'	0.052 (3)	0.056 (3)	0.047 (2)	-0.011 (2)	0.0082 (18)	-0.007 (2)
C2'	0.071 (4)	0.084 (5)	0.050 (3)	-0.020 (3)	0.016 (3)	-0.018 (3)
C3'	0.105 (12)	0.130 (13)	0.049 (6)	-0.041 (9)	-0.004 (7)	0.016 (7)
C4'	0.082 (8)	0.148 (14)	0.096 (7)	-0.042 (7)	-0.030 (6)	0.037 (7)
C5'	0.061 (4)	0.077 (6)	0.107 (6)	-0.020 (4)	0.007 (4)	-0.002 (4)
C6'	0.066 (4)	0.060 (5)	0.061 (4)	-0.027 (3)	0.000 (3)	-0.009 (3)

Geometric parameters (Å, °)

S1—C1	1.726 (3)	C3—H3A	0.97
S1—Na1 ⁱ	2.9644 (15)	C3—H3B	0.97
S1—Na1	3.0649 (15)	C4—C5	1.530 (18)
S2—C1	1.710 (3)	C4—H4A	0.97
Na1—O2	2.360 (3)	C4—H4B	0.97

Na1—O1	2.385 (2)	C5—C6	1.521 (14)
Na1—O1 ⁱⁱ	2.416 (3)	C5—H5A	0.97
Na1—O2 ⁱⁱⁱ	2.515 (2)	C5—H5B	0.97
Na1—S1 ^{iv}	2.9644 (15)	C6—H6A	0.97
Na1—Na1 ⁱⁱ	3.490 (3)	C6—H6B	0.97
Na1—Na1 ⁱⁱⁱ	3.644 (3)	N1'—C2'	1.452 (6)
Na1—H12O	2.67 (3)	N1'—C6'	1.473 (7)
O1—Na1 ⁱⁱ	2.416 (3)	C2'—C3'	1.495 (19)
O1—H11O	0.839 (17)	C2'—H2'1	0.97
O1—H12O	0.852 (17)	C2'—H2'2	0.97
O2—Na1 ⁱⁱⁱ	2.515 (2)	C3'—C4'	1.50 (3)
O2—H21O	0.834 (17)	C3'—H3'1	0.97
O2—H22O	0.856 (17)	C3'—H3'2	0.97
C1—N1'	1.368 (5)	C4'—C5'	1.512 (17)
C1—N1	1.403 (6)	C4'—H4'1	0.97
N1—C2	1.465 (9)	C4'—H4'2	0.97
N1—C6	1.489 (10)	C5'—C6'	1.507 (11)
C2—C3	1.47 (3)	C5'—H5'1	0.97
C2—H2A	0.97	C5'—H5'2	0.97
C2—H2B	0.97	C6'—H6'1	0.97
C3—C4	1.55 (3)	C6'—H6'2	0.97
C1—S1—Na1 ⁱ	112.20 (10)	C3—C2—H2A	108.9
C1—S1—Na1	131.13 (10)	N1—C2—H2B	108.9
Na1 ⁱ —S1—Na1	116.43 (4)	C3—C2—H2B	108.9
O2—Na1—O1	93.59 (8)	H2A—C2—H2B	107.8
O2—Na1—O1 ⁱⁱ	169.15 (8)	C2—C3—C4	108.3 (14)
O1—Na1—O1 ⁱⁱ	86.75 (8)	C2—C3—H3A	110
O2—Na1—O2 ⁱⁱⁱ	83.29 (8)	C4—C3—H3A	110
O1—Na1—O2 ⁱⁱⁱ	82.36 (7)	C2—C3—H3B	110
O1 ⁱⁱ —Na1—O2 ⁱⁱⁱ	86.01 (8)	C4—C3—H3B	110
O2—Na1—S1 ^{iv}	87.16 (7)	H3A—C3—H3B	108.4
O1—Na1—S1 ^{iv}	166.91 (6)	C5—C4—C3	113.1 (13)
O1 ⁱⁱ —Na1—S1 ^{iv}	90.09 (7)	C5—C4—H4A	109
O2 ⁱⁱⁱ —Na1—S1 ^{iv}	84.76 (6)	C3—C4—H4A	109
O2—Na1—S1	108.51 (7)	C5—C4—H4B	109
O1—Na1—S1	100.34 (7)	C3—C4—H4B	109
O1 ⁱⁱ —Na1—S1	82.06 (7)	H4A—C4—H4B	107.8
O2 ⁱⁱⁱ —Na1—S1	167.57 (6)	C6—C5—C4	108.2 (9)
S1 ^{iv} —Na1—S1	91.78 (4)	C6—C5—H5A	110.1
O2—Na1—Na1 ⁱⁱ	136.28 (8)	C4—C5—H5A	110.1
O1—Na1—Na1 ⁱⁱ	43.73 (6)	C6—C5—H5B	110.1
O1 ⁱⁱ —Na1—Na1 ⁱⁱ	43.02 (5)	C4—C5—H5B	110.1
O2 ⁱⁱⁱ —Na1—Na1 ⁱⁱ	82.00 (7)	H5A—C5—H5B	108.4
S1 ^{iv} —Na1—Na1 ⁱⁱ	131.81 (6)	N1—C6—C5	110.1 (6)
S1—Na1—Na1 ⁱⁱ	91.55 (6)	N1—C6—H6A	109.6
O2—Na1—Na1 ⁱⁱⁱ	43.27 (6)	C5—C6—H6A	109.6
O1—Na1—Na1 ⁱⁱⁱ	87.06 (7)	N1—C6—H6B	109.6

O1 ⁱⁱ —Na1—Na1 ⁱⁱⁱ	126.00 (8)	C5—C6—H6B	109.6
O2 ⁱⁱⁱ —Na1—Na1 ⁱⁱⁱ	40.02 (5)	H6A—C6—H6B	108.2
S1 ^{iv} —Na1—Na1 ⁱⁱⁱ	84.54 (5)	C1—N1'—C2'	124.6 (4)
S1—Na1—Na1 ⁱⁱⁱ	151.58 (5)	C1—N1'—C6'	122.6 (4)
Na1 ⁱⁱ —Na1—Na1 ⁱⁱⁱ	111.83 (7)	C2'—N1'—C6'	111.9 (5)
O2—Na1—H12O	80.3 (6)	N1'—C2'—C3'	109.4 (7)
O1—Na1—H12O	18.4 (4)	N1'—C2'—H2'1	109.8
O1 ⁱⁱ —Na1—H12O	97.4 (7)	C3'—C2'—H2'1	109.8
O2 ⁱⁱⁱ —Na1—H12O	68.4 (5)	N1'—C2'—H2'2	109.8
S1 ^{iv} —Na1—H12O	151.4 (5)	C3'—C2'—H2'2	109.8
S1—Na1—H12O	116.5 (5)	H2'1—C2'—H2'2	108.2
Na1 ⁱⁱ —Na1—H12O	56.0 (6)	C2'—C3'—C4'	116.5 (14)
Na1 ⁱⁱⁱ —Na1—H12O	68.7 (4)	C2'—C3'—H3'1	108.2
Na1—O1—Na1 ⁱⁱ	93.25 (8)	C4'—C3'—H3'1	108.2
Na1—O1—H11O	129 (2)	C2'—C3'—H3'2	108.2
Na1 ⁱⁱ —O1—H11O	97 (3)	C4'—C3'—H3'2	108.2
Na1—O1—H12O	100 (2)	H3'1—C3'—H3'2	107.3
Na1 ⁱⁱ —O1—H12O	124 (2)	C3'—C4'—C5'	106.7 (11)
H11O—O1—H12O	114 (3)	C3'—C4'—H4'1	110.4
Na1—O2—Na1 ⁱⁱⁱ	96.71 (8)	C5'—C4'—H4'1	110.4
Na1—O2—H21O	121 (3)	C3'—C4'—H4'2	110.4
Na1 ⁱⁱⁱ —O2—H21O	96 (2)	C5'—C4'—H4'2	110.4
Na1—O2—H22O	121 (2)	H4'1—C4'—H4'2	108.6
Na1 ⁱⁱⁱ —O2—H22O	106 (2)	C6'—C5'—C4'	112.5 (7)
H21O—O2—H22O	111 (2)	C6'—C5'—H5'1	109.1
N1'—C1—N1	40.3 (3)	C4'—C5'—H5'1	109.1
N1'—C1—S2	117.2 (3)	C6'—C5'—H5'2	109.1
N1—C1—S2	117.4 (3)	C4'—C5'—H5'2	109.1
N1'—C1—S1	118.6 (3)	H5'1—C5'—H5'2	107.8
N1—C1—S1	116.2 (3)	N1'—C6'—C5'	110.6 (5)
S2—C1—S1	121.25 (16)	N1'—C6'—H6'1	109.5
C1—N1—C2	123.5 (5)	C5'—C6'—H6'1	109.5
C1—N1—C6	123.9 (6)	N1'—C6'—H6'2	109.5
C2—N1—C6	111.0 (6)	C5'—C6'—H6'2	109.5
N1—C2—C3	113.2 (13)	H6'1—C6'—H6'2	108.1
N1—C2—H2A	108.9		
C1—S1—Na1—O2	38.84 (16)	Na1 ⁱ —S1—C1—S2	-10.8 (2)
Na1 ⁱ —S1—Na1—O2	-135.11 (7)	Na1—S1—C1—S2	175.08 (10)
C1—S1—Na1—O1	-58.52 (16)	N1'—C1—N1—C2	-89.0 (9)
Na1 ⁱ —S1—Na1—O1	127.53 (7)	S2—C1—N1—C2	11.7 (9)
C1—S1—Na1—O1 ⁱⁱ	-143.72 (15)	S1—C1—N1—C2	166.8 (6)
Na1 ⁱ —S1—Na1—O1 ⁱⁱ	42.34 (6)	N1'—C1—N1—C6	75.0 (8)
C1—S1—Na1—O2 ⁱⁱⁱ	-160.0 (3)	S2—C1—N1—C6	175.6 (6)
Na1 ⁱ —S1—Na1—O2 ⁱⁱⁱ	26.0 (3)	S1—C1—N1—C6	-29.3 (9)
C1—S1—Na1—S1 ^{iv}	126.44 (14)	C1—N1—C2—C3	104.9 (12)
Na1 ⁱ —S1—Na1—S1 ^{iv}	-47.50 (6)	C6—N1—C2—C3	-60.8 (14)
C1—S1—Na1—Na1 ⁱⁱ	-101.64 (15)	N1—C2—C3—C4	55 (2)

Na1 ⁱ —S1—Na1—Na1 ⁱⁱ	84.41 (5)	C2—C3—C4—C5	-53 (2)
C1—S1—Na1—Na1 ⁱⁱⁱ	44.63 (19)	C3—C4—C5—C6	54.8 (15)
Na1 ⁱ —S1—Na1—Na1 ⁱⁱⁱ	-129.32 (11)	C1—N1—C6—C5	-105.1 (9)
O2—Na1—O1—Na1 ⁱⁱ	169.13 (8)	C2—N1—C6—C5	60.6 (11)
O1 ⁱⁱ —Na1—O1—Na1 ⁱⁱ	0	C4—C5—C6—N1	-57.2 (11)
O2 ⁱⁱⁱ —Na1—O1—Na1 ⁱⁱ	86.40 (8)	N1—C1—N1'—C2'	90.7 (8)
S1 ^{iv} —Na1—O1—Na1 ⁱⁱ	76.2 (3)	S2—C1—N1'—C2'	-10.6 (7)
S1—Na1—O1—Na1 ⁱⁱ	-81.31 (7)	S1—C1—N1'—C2'	-171.6 (4)
Na1 ⁱⁱⁱ —Na1—O1—Na1 ⁱⁱ	126.33 (8)	N1—C1—N1'—C6'	-77.5 (6)
O1—Na1—O2—Na1 ⁱⁱⁱ	-81.86 (8)	S2—C1—N1'—C6'	-178.7 (4)
O1 ⁱⁱ —Na1—O2—Na1 ⁱⁱⁱ	9.6 (5)	S1—C1—N1'—C6'	20.2 (7)
O2 ⁱⁱⁱ —Na1—O2—Na1 ⁱⁱⁱ	0	C1—N1'—C2'—C3'	-113.5 (12)
S1 ^{iv} —Na1—O2—Na1 ⁱⁱⁱ	85.05 (6)	C6'—N1'—C2'—C3'	55.7 (13)
S1—Na1—O2—Na1 ⁱⁱⁱ	175.98 (6)	N1'—C2'—C3'—C4'	-55 (2)
Na1 ⁱⁱ —Na1—O2—Na1 ⁱⁱⁱ	-70.99 (11)	C2'—C3'—C4'—C5'	52.4 (19)
Na1 ⁱ —S1—C1—N1'	149.5 (3)	C3'—C4'—C5'—C6'	-52.4 (15)
Na1—S1—C1—N1'	-24.6 (4)	C1—N1'—C6'—C5'	110.7 (7)
Na1 ⁱ —S1—C1—N1	-164.9 (4)	C2'—N1'—C6'—C5'	-58.7 (9)
Na1—S1—C1—N1	21.0 (5)	C4'—C5'—C6'—N1'	57.7 (11)

Symmetry codes: (i) $-x+3/2, y+1/2, -z+1$; (ii) $-x+1, -y, -z+1$; (iii) $-x+1, -y-1, -z+1$; (iv) $-x+3/2, y-1/2, -z+1$.

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
O1—H11O...S2 ^v	0.84 (2)	2.39 (2)	3.214 (2)	167 (3)
O1—H12O...S1 ^{vi}	0.85 (2)	2.49 (2)	3.322 (3)	167 (3)
O2—H21O...S2 ^{vi}	0.83 (2)	2.48 (2)	3.283 (2)	163 (3)
O2—H22O...S1 ^{vii}	0.86 (2)	2.46 (2)	3.313 (2)	174 (3)

Symmetry codes: (v) $x-1/2, -y+1/2, z$; (vi) $x-1/2, -y-1/2, z$; (vii) $x, y-1, z$.