

(*E,E*)-4-[4-(4-Chloroanilino)-1-hydroxybut-2-enylidene]-3-methyl-5-oxo-4,5-dihydro-1*H*-pyrazol-1-yl}-benzenesulfonamide

Abdullah M. Asiri,^a Hassan M. Faidallah^a and Seik Weng Ng^{b*}

^aChemistry Department, Faculty of Science, King Abdul Aziz University, Jeddah 21589, Saudi Arabia, and ^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: seikweng@um.edu.my


Received 26 May 2011; accepted 30 May 2011

Key indicators: single-crystal X-ray study; $T = 100\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$; R factor = 0.054; wR factor = 0.125; data-to-parameter ratio = 12.5.

The molecule of the title compound, $\text{C}_{20}\text{H}_{19}\text{ClN}_4\text{O}_4\text{S}$, features a central pyrazole ring that possesses a benzene substituent, as well as a conjugated $=\text{C}-\text{C}=\text{C}-\text{C}_{\text{methyl}}$ substituent. The benzene ring is slightly twisted [dihedral angle = $7.7(2)^\circ$] with respect to the five-membered ring; the mean plane of the zigzag $=\text{C}-\text{C}=\text{C}-\text{C}$ fragment [torsion angle = $178.0(4)^\circ$] is also slightly twisted [dihedral angle = $10.6(4)^\circ$]. The amine and hydroxy groups form intramolecular hydrogen bonds. The amide group uses one of its H atoms to form a hydrogen bond to the sulfamyl O atom of an inversion-related molecule. Adjacent dimers are further linked by an $\text{N}-\text{H}_{\text{amido}}\cdots\text{N}_{\text{pyrazole}}$ hydrogen bond to generate a linear chain. The crystal studied is a nonmerohedral twin with a minor twin component of 25.6 (2)%.

Related literature

For the synthesis of 4-acetoacetyl-3-methyl-5-onyl-1-phenyl-pyrazole, see: Gelin *et al.* (1983).



Experimental

Crystal data

$\text{C}_{20}\text{H}_{19}\text{ClN}_4\text{O}_4\text{S}$	$V = 1932.6(4)\text{ \AA}^3$
$M_r = 446.90$	$Z = 4$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 14.7513(17)\text{ \AA}$	$\mu = 0.34\text{ mm}^{-1}$
$b = 17.545(2)\text{ \AA}$	$T = 100\text{ K}$
$c = 7.6203(9)\text{ \AA}$	$0.20 \times 0.02 \times 0.02\text{ mm}$
$\beta = 101.496(2)^\circ$	

Data collection

Bruker SMART APEX	32586 measured reflections
diffractometer	3426 independent reflections
Absorption correction: multi-scan (TWINABS; Bruker, 2009)	2605 reflections with $I > 2\sigma(I)$
$T_{\min} = 0.935$, $T_{\max} = 0.993$	$R_{\text{int}} = 0.099$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.054$	275 parameters
$wR(F^2) = 0.125$	H-atom parameters constrained
$S = 1.04$	$\Delta\rho_{\max} = 0.39\text{ e \AA}^{-3}$
3426 reflections	$\Delta\rho_{\min} = -0.47\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1—H1O \cdots O2	0.84	1.79	2.498 (4)	141
N1—H1 \cdots O1	0.88	2.00	2.659 (4)	131
N1—H1 \cdots O3 ⁱ	0.88	2.34	3.093 (4)	143
N4—H41 \cdots N2 ⁱⁱ	0.88	2.16	3.003 (4)	161
N4—H42 \cdots O4 ⁱⁱⁱ	0.88	2.09	2.917 (4)	156

Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$; (ii) $x, y, z - 1$; (iii) $-x, -y + 1, -z$.

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2010).

We thank King Abdul Aziz University and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU5227).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Bruker (2009). *APEX2, SAINT* and *TWINABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Gelin, S., Chantegrel, B. & Nadi, A. I. (1983). *J. Org. Chem.* **48**, 4078–4082.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2011). E67, o1590 [doi:10.1107/S1600536811020617]

(E,E)-4-{4-[3-(4-Chloroanilino)-1-hydroxybut-2-enylidene]-3-methyl-5-oxo-4,5-dihydro-1H-pyrazol-1-yl}benzenesulfonamide

Abdullah M. Asiri, Hassan M. Faidallah and Seik Weng Ng

S1. Comment

The compound, 4-acetoacetyl-3-methyl-5-onyl-1-phenylpyrazole rearranges under the influence of acetic acid to form functionalized 4-oxopyrano[2,3-*c*]pyrazoles (Gelin *et al.*, 1983). The addition of a sulfamido unit to the phenyl ring is expected to improve its biological activity; in the present study, the *p*-sulfamyl analog is reacted with chloroaniline to yield a new *p*-sulfamylphenylpyrazole derivative (Scheme I). The C₂₀H₁₉ClN₄O₄S (Fig. 1) features a central pyrazole ring that possesses a benzene substituent as well as a conjugated =C—C=C—C_{methyl} substituent. The benzene ring is slightly twisted with respect to the five-membered ring; the mean plane of the zigzag =C—C=C—C fragment is also slightly twisted.

The amino and hydroxy groups are intramolecular hydrogen-bond donors. The amido group uses one of its H atoms to form an hydrogen bond to the sulfamyl O atom of an inversion-related molecule. Adjacent dimers are further linked by an N—H_{amido}···N_{pyrazole} hydrogen bond to generate a linear chain motif (Table 2, Fig. 2).

S2. Experimental


4-Acetoacetyl-3-methyl-5-onyl-1-*p*-sulfamylphenylpyrazole was synthesized by using a literature procedure (Gelin *et al.*, 1983). The compound (1.70 g, 0.005 mol) and 4-chloroaniline (0.63 g, 0.005 mol) were heated in ethanol (25 ml) for 2 h. The solid that separated from solution was collected and recrystallized from ethanol to yield yellow prismatic crystals.

S3. Refinement


The crystal is a non-merohedral twin with a minor twin domain of 25.6 (2)%. Owing to twinning, the amino, amido and hydroxy H-atoms were generated geometrically.

Carbon-bound H-atoms were placed in calculated positions (C—H 0.95 to 0.98, N—H 0.86, O—H 0.84 Å) and were included in the refinement in the riding model approximation, with *U*(H) set to 1.2 to 1.5*U*_{eq}(C,N,O). An *sp*²-type of hybridization was assumed for the amino and hydroxy H atoms, and an *sp*³-type of hybridization for the amido H atoms.

Omitted because of bad disagreement was (4 4 1).

**Figure 1**

Thermal ellipsoid plot (Barbour, 2001) of $C_{20}H_{19}ClN_4O_4S$ at the 70% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.

**Figure 2**

Hydrogen-bonded chain motif.

(E,E)-4-[3-(4-Chloroanilino)-1-hydroxybut-2-enylidene]- 3-methyl-5-oxo-4,5-dihydro-1*H*-pyrazol-1-yl}benzenesulfonamide

Crystal data

$C_{20}H_{19}ClN_4O_4S$

$M_r = 446.90$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 14.7513 (17) \text{ \AA}$

$b = 17.545 (2) \text{ \AA}$

$c = 7.6203 (9) \text{ \AA}$

$\beta = 101.496 (2)^\circ$

$V = 1932.6 (4) \text{ \AA}^3$

$Z = 4$

$F(000) = 928$

$D_x = 1.536 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 656 reflections

$\theta = 2.3\text{--}20.1^\circ$

$\mu = 0.34 \text{ mm}^{-1}$

$T = 100 \text{ K}$

Prism, yellow

$0.20 \times 0.02 \times 0.02 \text{ mm}$

Data collection

Bruker SMART APEX
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 ω scans
Absorption correction: multi-scan
(TWINABS; Bruker, 2009)
 $T_{\min} = 0.935$, $T_{\max} = 0.993$

32586 measured reflections
3426 independent reflections
2605 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.099$
 $\theta_{\max} = 25.1^\circ$, $\theta_{\min} = 1.4^\circ$
 $h = -17 \rightarrow 17$
 $k = 0 \rightarrow 20$
 $l = 0 \rightarrow 9$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.054$
 $wR(F^2) = 0.125$
 $S = 1.04$
3426 reflections
275 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0457P)^2 + 2.8595P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.39 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.47 \text{ e } \text{\AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	1.06569 (7)	0.72948 (6)	1.73048 (15)	0.0256 (3)
S1	0.14888 (7)	0.47564 (6)	0.00333 (13)	0.0137 (2)
O1	0.58091 (18)	0.59404 (16)	1.0470 (4)	0.0200 (7)
H1O	0.5745	0.5767	0.9424	0.030*
O2	0.48625 (17)	0.55620 (16)	0.7504 (4)	0.0187 (6)
O3	0.22069 (18)	0.43897 (16)	-0.0680 (4)	0.0194 (7)
O4	0.07052 (18)	0.43097 (15)	0.0288 (4)	0.0169 (6)
N1	0.6886 (2)	0.64430 (18)	1.3466 (4)	0.0147 (7)
H1	0.6866	0.6177	1.2480	0.018*
N2	0.2764 (2)	0.63178 (18)	0.8250 (4)	0.0129 (7)
N3	0.3294 (2)	0.59328 (17)	0.7196 (4)	0.0115 (7)
N4	0.1079 (2)	0.54401 (18)	-0.1294 (4)	0.0158 (7)
H41	0.1513	0.5783	-0.1304	0.024*
H42	0.0610	0.5652	-0.0922	0.024*
C1	0.7783 (3)	0.6666 (2)	1.4438 (5)	0.0181 (9)
C2	0.8495 (3)	0.6126 (2)	1.4652 (5)	0.0195 (9)
H2	0.8370	0.5625	1.4200	0.023*
C3	0.9374 (3)	0.6319 (2)	1.5512 (5)	0.0200 (9)
H3	0.9860	0.5955	1.5648	0.024*
C4	0.9541 (3)	0.7048 (2)	1.6175 (5)	0.0187 (9)
C5	0.8863 (3)	0.7590 (2)	1.5964 (5)	0.0214 (10)
H5	0.8998	0.8088	1.6429	0.026*
C6	0.7972 (3)	0.7406 (2)	1.5065 (6)	0.0205 (9)
H6	0.7499	0.7782	1.4880	0.025*
C7	0.6074 (3)	0.6606 (2)	1.3933 (5)	0.0140 (9)

C8	0.6086 (3)	0.6938 (2)	1.5746 (5)	0.0218 (10)
H8A	0.6646	0.6771	1.6577	0.033*
H8B	0.5539	0.6766	1.6182	0.033*
H8C	0.6081	0.7496	1.5666	0.033*
C9	0.5228 (3)	0.6492 (2)	1.2803 (5)	0.0139 (8)
H9	0.4692	0.6617	1.3257	0.017*
C10	0.5088 (3)	0.6208 (2)	1.1047 (5)	0.0149 (8)
C11	0.2967 (3)	0.6930 (2)	1.1153 (5)	0.0172 (9)
H11A	0.2318	0.7065	1.0700	0.026*
H11B	0.3335	0.7397	1.1420	0.026*
H11C	0.3018	0.6626	1.2246	0.026*
C12	0.3320 (3)	0.6480 (2)	0.9771 (5)	0.0118 (8)
C13	0.4237 (3)	0.6199 (2)	0.9796 (5)	0.0141 (8)
C14	0.4195 (3)	0.5859 (2)	0.8100 (5)	0.0133 (8)
C15	0.2875 (2)	0.5682 (2)	0.5466 (5)	0.0113 (8)
C16	0.3375 (3)	0.5229 (2)	0.4485 (5)	0.0148 (8)
H16	0.4003	0.5103	0.4957	0.018*
C17	0.2940 (3)	0.4966 (2)	0.2817 (5)	0.0149 (8)
H17	0.3275	0.4660	0.2137	0.018*
C18	0.2025 (3)	0.5143 (2)	0.2129 (5)	0.0136 (8)
C19	0.1535 (3)	0.5618 (2)	0.3080 (5)	0.0154 (9)
H19	0.0913	0.5754	0.2589	0.018*
C20	0.1966 (3)	0.5890 (2)	0.4746 (5)	0.0134 (8)
H20	0.1641	0.6218	0.5397	0.016*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl1	0.0176 (5)	0.0285 (6)	0.0281 (6)	-0.0015 (4)	-0.0013 (4)	-0.0037 (5)
S1	0.0109 (5)	0.0185 (5)	0.0116 (5)	0.0000 (4)	0.0020 (4)	-0.0030 (4)
O1	0.0134 (14)	0.0299 (17)	0.0165 (15)	0.0024 (12)	0.0022 (12)	-0.0079 (13)
O2	0.0117 (14)	0.0284 (16)	0.0166 (15)	0.0036 (12)	0.0045 (12)	-0.0071 (12)
O3	0.0152 (14)	0.0258 (16)	0.0168 (15)	0.0035 (12)	0.0022 (12)	-0.0089 (12)
O4	0.0136 (14)	0.0178 (15)	0.0196 (15)	-0.0041 (12)	0.0036 (12)	-0.0012 (12)
N1	0.0106 (16)	0.0211 (18)	0.0110 (17)	0.0011 (14)	-0.0017 (13)	0.0021 (14)
N2	0.0121 (17)	0.0146 (18)	0.0129 (17)	0.0005 (14)	0.0045 (14)	-0.0004 (13)
N3	0.0080 (16)	0.0151 (17)	0.0119 (17)	0.0014 (13)	0.0032 (13)	-0.0041 (13)
N4	0.0108 (16)	0.0227 (19)	0.0133 (17)	-0.0015 (14)	0.0012 (14)	0.0002 (14)
C1	0.018 (2)	0.022 (2)	0.013 (2)	-0.0011 (18)	0.0024 (17)	0.0011 (17)
C2	0.027 (2)	0.016 (2)	0.014 (2)	0.0011 (18)	0.0015 (18)	-0.0009 (17)
C3	0.018 (2)	0.022 (2)	0.020 (2)	0.0026 (18)	0.0022 (19)	0.0029 (18)
C4	0.019 (2)	0.023 (2)	0.014 (2)	0.0006 (18)	0.0035 (17)	0.0010 (17)
C5	0.024 (2)	0.020 (2)	0.020 (2)	-0.0020 (19)	0.0022 (18)	-0.0028 (18)
C6	0.020 (2)	0.018 (2)	0.023 (2)	0.0012 (19)	0.0032 (18)	0.0014 (18)
C7	0.017 (2)	0.010 (2)	0.015 (2)	0.0028 (16)	0.0034 (17)	0.0035 (16)
C8	0.022 (2)	0.026 (2)	0.017 (2)	-0.0044 (19)	0.0022 (18)	-0.0032 (18)
C9	0.0119 (19)	0.018 (2)	0.013 (2)	0.0014 (16)	0.0033 (16)	0.0000 (16)
C10	0.014 (2)	0.014 (2)	0.017 (2)	0.0003 (17)	0.0055 (17)	-0.0001 (17)

C11	0.017 (2)	0.022 (2)	0.014 (2)	0.0027 (18)	0.0042 (17)	-0.0014 (17)
C12	0.013 (2)	0.0102 (19)	0.013 (2)	-0.0018 (16)	0.0026 (16)	0.0002 (15)
C13	0.0119 (19)	0.016 (2)	0.015 (2)	-0.0017 (16)	0.0040 (17)	-0.0009 (16)
C14	0.0120 (19)	0.014 (2)	0.013 (2)	-0.0008 (16)	0.0006 (16)	-0.0002 (16)
C15	0.0139 (19)	0.011 (2)	0.0101 (19)	-0.0040 (15)	0.0037 (16)	0.0020 (15)
C16	0.0108 (19)	0.019 (2)	0.014 (2)	-0.0025 (17)	0.0016 (16)	-0.0009 (17)
C17	0.0153 (19)	0.018 (2)	0.013 (2)	0.0041 (17)	0.0055 (17)	0.0008 (16)
C18	0.015 (2)	0.015 (2)	0.0098 (19)	-0.0024 (17)	0.0006 (16)	-0.0030 (16)
C19	0.0121 (19)	0.018 (2)	0.016 (2)	0.0027 (17)	0.0023 (16)	0.0000 (17)
C20	0.013 (2)	0.015 (2)	0.014 (2)	0.0018 (16)	0.0064 (17)	-0.0009 (16)

Geometric parameters (\AA , $^\circ$)

C11—C4	1.753 (4)	C6—H6	0.9500
S1—O3	1.436 (3)	C7—C9	1.382 (5)
S1—O4	1.441 (3)	C7—C8	1.497 (5)
S1—N4	1.607 (3)	C8—H8A	0.9800
S1—C18	1.770 (4)	C8—H8B	0.9800
O1—C10	1.316 (5)	C8—H8C	0.9800
O1—H1O	0.8400	C9—C10	1.404 (5)
O2—C14	1.275 (5)	C9—H9	0.9500
N1—C7	1.346 (5)	C10—C13	1.418 (5)
N1—C1	1.435 (5)	C11—C12	1.491 (5)
N1—H1	0.8800	C11—H11A	0.9800
N2—C12	1.310 (5)	C11—H11B	0.9800
N2—N3	1.401 (4)	C11—H11C	0.9800
N3—C14	1.376 (5)	C12—C13	1.435 (5)
N3—C15	1.410 (5)	C13—C14	1.414 (5)
N4—H41	0.8800	C15—C20	1.392 (5)
N4—H42	0.8800	C15—C16	1.398 (5)
C1—C6	1.393 (6)	C16—C17	1.383 (5)
C1—C2	1.399 (6)	C16—H16	0.9500
C2—C3	1.374 (6)	C17—C18	1.382 (5)
C2—H2	0.9500	C17—H17	0.9500
C3—C4	1.379 (6)	C18—C19	1.397 (5)
C3—H3	0.9500	C19—C20	1.387 (5)
C4—C5	1.365 (6)	C19—H19	0.9500
C5—C6	1.393 (5)	C20—H20	0.9500
C5—H5	0.9500		
O3—S1—O4	118.65 (17)	H8A—C8—H8C	109.5
O3—S1—N4	108.04 (17)	H8B—C8—H8C	109.5
O4—S1—N4	106.22 (17)	C7—C9—C10	126.0 (4)
O3—S1—C18	106.40 (17)	C7—C9—H9	117.0
O4—S1—C18	108.26 (17)	C10—C9—H9	117.0
N4—S1—C18	109.02 (17)	O1—C10—C9	118.1 (3)
C10—O1—H1O	120.0	O1—C10—C13	116.0 (3)
C7—N1—C1	125.6 (3)	C9—C10—C13	126.0 (4)

C7—N1—H1	117.2	C12—C11—H11A	109.5
C1—N1—H1	117.2	C12—C11—H11B	109.5
C12—N2—N3	106.9 (3)	H11A—C11—H11B	109.5
C14—N3—N2	110.6 (3)	C12—C11—H11C	109.5
C14—N3—C15	129.7 (3)	H11A—C11—H11C	109.5
N2—N3—C15	119.8 (3)	H11B—C11—H11C	109.5
S1—N4—H41	109.5	N2—C12—C13	111.0 (3)
S1—N4—H42	109.5	N2—C12—C11	119.7 (3)
H41—N4—H42	109.5	C13—C12—C11	129.2 (3)
C6—C1—C2	119.7 (4)	C14—C13—C10	119.4 (3)
C6—C1—N1	122.2 (4)	C14—C13—C12	105.3 (3)
C2—C1—N1	118.0 (4)	C10—C13—C12	135.3 (4)
C3—C2—C1	120.3 (4)	O2—C14—N3	126.7 (3)
C3—C2—H2	119.9	O2—C14—C13	127.0 (3)
C1—C2—H2	119.9	N3—C14—C13	106.2 (3)
C2—C3—C4	119.1 (4)	C20—C15—C16	120.5 (3)
C2—C3—H3	120.5	C20—C15—N3	119.6 (3)
C4—C3—H3	120.5	C16—C15—N3	119.9 (3)
C5—C4—C3	121.9 (4)	C17—C16—C15	118.9 (4)
C5—C4—C11	118.7 (3)	C17—C16—H16	120.5
C3—C4—C11	119.4 (3)	C15—C16—H16	120.5
C4—C5—C6	119.6 (4)	C18—C17—C16	120.8 (4)
C4—C5—H5	120.2	C18—C17—H17	119.6
C6—C5—H5	120.2	C16—C17—H17	119.6
C1—C6—C5	119.3 (4)	C17—C18—C19	120.3 (3)
C1—C6—H6	120.3	C17—C18—S1	118.9 (3)
C5—C6—H6	120.3	C19—C18—S1	120.8 (3)
N1—C7—C9	123.0 (4)	C20—C19—C18	119.3 (4)
N1—C7—C8	118.6 (3)	C20—C19—H19	120.3
C9—C7—C8	118.3 (3)	C18—C19—H19	120.3
C7—C8—H8A	109.5	C19—C20—C15	120.0 (4)
C7—C8—H8B	109.5	C19—C20—H20	120.0
H8A—C8—H8B	109.5	C15—C20—H20	120.0
C7—C8—H8C	109.5		
C12—N2—N3—C14	0.0 (4)	C11—C12—C13—C10	2.9 (8)
C12—N2—N3—C15	179.8 (3)	N2—N3—C14—O2	-177.4 (4)
C7—N1—C1—C6	-46.9 (6)	C15—N3—C14—O2	2.8 (7)
C7—N1—C1—C2	137.2 (4)	N2—N3—C14—C13	0.8 (4)
C6—C1—C2—C3	1.5 (6)	C15—N3—C14—C13	-179.0 (4)
N1—C1—C2—C3	177.5 (4)	C10—C13—C14—O2	-2.1 (6)
C1—C2—C3—C4	0.6 (6)	C12—C13—C14—O2	177.0 (4)
C2—C3—C4—C5	-1.6 (6)	C10—C13—C14—N3	179.7 (3)
C2—C3—C4—C11	179.0 (3)	C12—C13—C14—N3	-1.2 (4)
C3—C4—C5—C6	0.4 (6)	C14—N3—C15—C20	-173.4 (4)
C11—C4—C5—C6	179.7 (3)	N2—N3—C15—C20	6.9 (5)
C2—C1—C6—C5	-2.7 (6)	C14—N3—C15—C16	6.5 (6)
N1—C1—C6—C5	-178.6 (4)	N2—N3—C15—C16	-173.2 (3)

C4—C5—C6—C1	1.8 (6)	C20—C15—C16—C17	-2.4 (6)
C1—N1—C7—C9	168.6 (4)	N3—C15—C16—C17	177.7 (3)
C1—N1—C7—C8	-10.3 (6)	C15—C16—C17—C18	-0.4 (6)
N1—C7—C9—C10	-0.9 (6)	C16—C17—C18—C19	2.7 (6)
C8—C7—C9—C10	178.0 (4)	C16—C17—C18—S1	-177.2 (3)
C7—C9—C10—O1	8.5 (6)	O3—S1—C18—C17	-7.9 (4)
C7—C9—C10—C13	-169.7 (4)	O4—S1—C18—C17	120.6 (3)
N3—N2—C12—C13	-0.8 (4)	N4—S1—C18—C17	-124.2 (3)
N3—N2—C12—C11	176.8 (3)	O3—S1—C18—C19	172.2 (3)
O1—C10—C13—C14	4.9 (6)	O4—S1—C18—C19	-59.2 (4)
C9—C10—C13—C14	-176.9 (4)	N4—S1—C18—C19	56.0 (4)
O1—C10—C13—C12	-173.8 (4)	C17—C18—C19—C20	-2.1 (6)
C9—C10—C13—C12	4.4 (8)	S1—C18—C19—C20	177.7 (3)
N2—C12—C13—C14	1.3 (4)	C18—C19—C20—C15	-0.7 (6)
C11—C12—C13—C14	-176.0 (4)	C16—C15—C20—C19	3.0 (6)
N2—C12—C13—C10	-179.8 (4)	N3—C15—C20—C19	-177.1 (3)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
O1—H1O···O2	0.84	1.79	2.498 (4)	141
N1—H1···O1	0.88	2.00	2.659 (4)	131
N1—H1···O3 ⁱ	0.88	2.34	3.093 (4)	143
N4—H41···N2 ⁱⁱ	0.88	2.16	3.003 (4)	161
N4—H42···O4 ⁱⁱⁱ	0.88	2.09	2.917 (4)	156

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $x, y, z-1$; (iii) $-x, -y+1, -z$.