

A second monoclinic polymorph of 2-[2-(4-methoxyphenyl)hydrazinylidene]-1,3-diphenylpropane-1,3-dione

Carlos Bustos,^a Luis Alvarez-Thon,^{b*} Daniela Barría,^a Juan-Guillermo Cárcamo^c and María Teresa Garland^d

^aInstituto de Ciencias Químicas, Universidad Austral de Chile, Avda. Los Robles s/n, Campus Isla Teja, Casilla 567, Valdivia, Chile, ^bDepartamento de Ciencias Físicas, Universidad Andres Bello, Avda. República 220, Santiago de Chile, Chile, ^cInstituto de Ciencias Moleculares y Microbiología, Universidad Austral de Chile, Avda. Los Robles s/n, Campus Isla Teja, Casilla 567, Valdivia, Chile, and ^dLaboratorio de Cristalografía, Departamento de Física, Facultad de Ciencias Físicas y Matemáticas, Universidad de Chile, Av. Blanco Encalada 2008, Santiago de Chile, Chile
Correspondence e-mail: lalvarez@unab.cl


Received 17 June 2011; accepted 21 June 2011

Key indicators: single-crystal X-ray study; $T = 150\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.039; wR factor = 0.105; data-to-parameter ratio = 14.5.

The title compound, $\text{C}_{22}\text{H}_{18}\text{N}_2\text{O}_3$ is the second monoclinic polymorph ($P2_1/c$) of the compound, the first being reported in space group $P2_1$ [Bertolasi *et al.* (1993). *J. Chem. Soc. Perkin Trans. 2*, pp. 2223–2228]. In the molecular structure of the title compound, the interplanar angle between the benzoyl units is $80.04(5)^\circ$, while the corresponding angles between the phenylhydrazinylidene and benzoyl groups are $36.11(5)$ and $55.77(2)^\circ$. A strong resonance-assisted intramolecular N—H···O hydrogen bond is found. In the crystal, the entire supramolecular structure is constructed by weak intermolecular C—H···O interactions and an inter-ring π – π interaction [centroid–centroid distance = $3.6088(8)\text{ \AA}$].

Related literature

For details of the synthesis, see: Yao (1964). For resonance-assisted hydrogen bonds and related structures, see: Bertolasi *et al.* (1993); Bustos, Alvarez-Thon, Barría *et al.* (2011); Bustos, Alvarez-Thon, Cárcamo, Garland & Sánchez (2011); Bustos, Alvarez-Thon, Cárcamo, Ibañez & Sánchez (2011); Gilli *et al.* (1993).



Experimental

Crystal data

$\text{C}_{22}\text{H}_{18}\text{N}_2\text{O}_3$	$V = 1774.3(3)\text{ \AA}^3$
$M_r = 358.38$	$Z = 4$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 12.3045(11)\text{ \AA}$	$\mu = 0.09\text{ mm}^{-1}$
$b = 11.055(1)\text{ \AA}$	$T = 150\text{ K}$
$c = 14.2435(13)\text{ \AA}$	$0.52 \times 0.26 \times 0.15\text{ mm}$
$\beta = 113.683(1)^\circ$	

Data collection

Bruker D8 Discover diffractometer with SMART CCD area detector 10646 measured reflections	3600 independent reflections 2894 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.036$
---	--

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.039$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.105$	$\Delta\rho_{\text{max}} = 0.26\text{ e \AA}^{-3}$
$S = 1.04$	$\Delta\rho_{\text{min}} = -0.18\text{ e \AA}^{-3}$
3600 reflections	
249 parameters	

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D\cdots H\cdots A$	$D\cdots H$	$H\cdots A$	$D\cdots A$	$D\cdots H\cdots A$
N2—H1···O2	0.884 (14)	1.977 (15)	2.6427 (14)	131.0 (14)
C15—H15···O3 ⁱ	0.95	2.64	3.5859 (16)	174
C21—H21···O1 ⁱⁱ	0.95	2.63	3.3441 (15)	132

Symmetry codes: (i) $-x + 1, -y + 1, -z + 2$; (ii) $-x + 1, -y, -z + 2$.

Data collection: SMART (Bruker, 2001); cell refinement: SAINT (Bruker, 2000); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: XP in SHELXTL-PC (Sheldrick, 2008); software used to prepare material for publication: PLATON (Spek, 2009) and Mercury (Macrae *et al.*, 2006).

The authors thank the Fondo Nacional de Desarrollo Científico y Tecnológico (FONDECYT; grant Nos. 11100446 and 1080269) and the Universidad Andrés Bello (grant No. DI-06-10-R) for financial assistance.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: ZS2122).

References

- Bertolasi, V., Ferretti, V., Gilli, P., Gilli, G., Issa, Y. M. & Sherif, O. E. (1993). *J. Chem. Soc. Perkin Trans. 2*, pp. 2223–2228.
- Bruker (2000). *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2001). *SMART*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bustos, C., Alvarez-Thon, L., Barría, D., Garland, M. T. & Sánchez, C. (2011). *Acta Cryst. E*67, o1587.
- Bustos, C., Alvarez-Thon, L., Cárcamo, J.-G., Garland, M. T. & Sánchez, C. (2011). *Acta Cryst. E*67, o1426.
- Bustos, C., Alvarez-Thon, L., Cárcamo, J.-G., Ibañez, A. & Sánchez, C. (2011). *Acta Cryst. E*67, o1450–o1451.
- Gilli, G., Bertolasi, V., Ferretti, V. & Gilli, P. (1993). *Acta Cryst. B*49, 564–576.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- Sheldrick, G. M. (2008). *Acta Cryst. A*64, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D*65, 148–155.
- Yao, H. C. (1964). *J. Org. Chem.* **29**, 2959–2962.

supporting information

Acta Cryst. (2011). E67, o1830–o1831 [doi:10.1107/S1600536811024445]

A second monoclinic polymorph of 2-[2-(4-methoxyphenyl)hydrazinylidene]-1,3-diphenylpropane-1,3-dione

Carlos Bustos, Luis Alvarez-Thon, Daniela Barría, Juan-Guillermo Cárcamo and María Teresa Garland

S1. Comment

Extensive experimental and theoretical efforts have been devoted to understand the reasons for the formation of the very strong intramolecular O—H···O hydrogen bond in β -diketoenolic systems (Bertolasi *et al.*, 1993). A model called RAHB (Resonance Assisted Hydrogen Bond) was suggested, which is essentially a synergistic mutual reinforcement of the H-bond with the π -delocalization within this heterodienic system (Gilli *et al.*, 1993). Besides, this concept has been applied to other heterodienic systems such as enaminones and ketohydrazones. On the other hand, in previous works we have reported the crystal structures of three β -diketohydrzones of the type 2-(*R*-phenyl)hydrazinylidene)-1,3-diphenylpropane-1,3-dione (*R* = 4-Br, 4-NO₂, 3-Cl) that contain this type of hydrogen bond (Bustos, Alvarez-Thon, Barría *et al.*, 2011; Bustos, Alvarez-Thon, Cárcamo, Garland & Sánchez, 2011; Bustos, Alvarez-Thon, Cárcamo, Ibañez & Sánchez, 2011). We now present the crystal structure of a second monoclinic polymorph of the compound C₂₂H₁₈N₂O₃ (I), described previously in the literature (Bertolasi *et al.*, 1993), prepared using similar methodology (Yao, 1964) and recrystallizing from ethanol as orange crystals. The title compound crystallizes in space group P2₁/c whereas the previously reported structure (Bertolasi *et al.*, 1993) crystallizes in space group P2₁. The most striking structural difference between the two polymorphs is in the methoxy group which appears rotated by 180°.

In the molecular structure of the title compound (Fig. 1), the interplanar angle between the benzoyl units is 80.04 (5)°. The corresponding angles between the phenylhydrazinylidene and the benzoyl groups are 36.11 (5) and 55.77 (2)°, respectively. In the crystal, a strong resonance-assisted intramolecular hydrogen bond, N2—H1···O2, is observed (Table 1).

The structure shows no conventional intermolecular hydrogen bonds and the entire supramolecular structure is constructed only by weak interactions. For the sake of clarity, the crystal packing can be described through the formation of two inversion-related dimers: firstly, a dimer is formed *via* weak intermolecular contacts of the type C15—H15···O3ⁱ, and a π — π stacking interaction with a Cg···Cgⁱ distance of 3.6088 (8) Å, where Cg is the centroid of the C16—C21 ring (Fig. 2) [symmetry code: (i) -x + 1, -y + 1, -z + 2]. Secondly, a dimer is formed *via* weak contacts of the type C21—H21···O1ⁱⁱ [3.3441 (15) Å] (Fig. 3) [symmetry code: (ii) -x + 1, -y, -z + 2].

S2. Experimental

Chemicals: 1,3-diphenylpropane-1,3-dione, *p*-anisidine and sodium nitrite were procured from Sigma-Aldrich and sodium hydroxide, hydrochloric acid, sodium acetate and solvents from Merck. These chemicals were used without previous purification.

Procedure: A solution of 2.29 g (0.01 mole) of 1,3-diphenylpropane-1,3-dione (98%) in 100 ml of ethanol solution containing 0.4 g (0.01 mole) of sodium hydroxide was buffered by adding 4.80 g of sodium acetate trihydrate. The resulting β -diketonate solution was diluted with water to a volume of about 220 ml and stirred and cooled to -5 °C. A diazonium ion solution was prepared adding 1.24 g (0.01 mole) of *p*-anisidine (99%) in 8 ml of hydrochloric acid (5 mol/L), cooled to -5 °C, and a saturated aqueous solution containing 0.69 g (0.01 mole) of sodium nitrite was added dropwise. This solution was then added dropwise with vigorous stirring into the buffered β -diketonate solution. During the addition a yellow precipitate was formed which was filtered by suction and washed with an abundant quantity of water. Yield: 95% of crude product. Orange single crystals suitable for X-ray studies were obtained by recrystallization from a concentrated solution of the compound in ethanol.

S3. Refinement

All hydrogen atoms were found in difference Fourier maps. The hydrogen attached to N2 was refined freely against the diffraction data, but all other H atoms were placed in geometrically idealized positions and constrained to ride on their parent atoms, with aromatic C—H = 0.95 Å, methyl C—H = 0.98 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}$ (aromatic C) or $1.5U_{\text{eq}}$ (aliphatic C).






Figure 1

View of the title compound with the atom numbering scheme. Displacement ellipsoids for non-H atoms are drawn at the 50% probability level. The strong intramolecular hydrogen bond (N2—H1 \cdots O2) is depicted with a dashed line.

**Figure 2**

Part of the crystal packing showing the inversion-related interaction *via* C15—H15 \cdots O3 $^{\text{i}}$ weak contacts (dashed lines) and a π - π stacking interaction Cg \cdots Cg $^{\text{i}}$ (solid line) [symmetry code: (i) $-x + 1, -y + 1, -z + 2$].

**Figure 3**

Part of the crystal packing showing the inversion-related interaction *via* C21—H21···O1ⁱⁱ contacts (dashed lines) [(ii) -x + 1, -y, -z + 2]].

2-[2-(4-methoxyphenyl)hydrazinylidene]-1,3-diphenylpropane-1,3-dione

Crystal data

C₂₂H₁₈N₂O₃
 $M_r = 358.38$
 Monoclinic, P2₁/c
 Hall symbol: -P 2ybc
 $a = 12.3045 (11)$ Å
 $b = 11.055 (1)$ Å
 $c = 14.2435 (13)$ Å
 $\beta = 113.683 (1)$ °
 $V = 1774.3 (3)$ Å³
 $Z = 4$

$F(000) = 752$
 $D_x = 1.342 \text{ Mg m}^{-3}$
 Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
 Cell parameters from 999 reflections
 $\theta = 1.8\text{--}26.4$ °
 $\mu = 0.09 \text{ mm}^{-1}$
 $T = 150$ K
 Polyhedron, orange
 $0.52 \times 0.26 \times 0.15$ mm

Data collection

Bruker D8 Discover
 diffractometer with SMART CCD area detector
 Radiation source: fine-focus sealed tube

Graphite monochromator
 φ and ω scans
 10646 measured reflections

3600 independent reflections
 2894 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.036$
 $\theta_{\text{max}} = 26.4^\circ$, $\theta_{\text{min}} = 1.8^\circ$

$h = -14 \rightarrow 15$
 $k = -13 \rightarrow 13$
 $l = -17 \rightarrow 17$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.039$
 $wR(F^2) = 0.105$
 $S = 1.04$
 3600 reflections
 249 parameters
 0 restraints
 Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map
 Hydrogen site location: inferred from neighbouring sites
 H atoms treated by a mixture of independent and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.061P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.001$
 $\Delta\rho_{\text{max}} = 0.26 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.18 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. Bond distances, angles etc. have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement on F^2 for ALL reflections except those flagged by the user for potential systematic errors. Weighted R -factors wR and all goodnesses of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The observed criterion of $F^2 > \sigma(F^2)$ is used only for calculating - R -factor-obs etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.23194 (8)	-0.07143 (8)	0.87203 (6)	0.0383 (3)
O2	0.49696 (7)	0.03714 (8)	1.12916 (6)	0.0323 (3)
O3	0.72539 (8)	0.62118 (8)	0.93267 (6)	0.0344 (3)
N1	0.39904 (9)	0.18280 (9)	0.94790 (7)	0.0266 (3)
N2	0.50300 (9)	0.22036 (9)	1.01218 (8)	0.0266 (3)
C1	0.31024 (11)	-0.05076 (11)	1.09863 (9)	0.0270 (4)
C2	0.35221 (12)	-0.15824 (11)	1.15131 (9)	0.0300 (4)
C3	0.27939 (13)	-0.22881 (12)	1.18252 (10)	0.0376 (4)
C4	0.16617 (13)	-0.18947 (13)	1.16436 (10)	0.0410 (5)
C5	0.12488 (12)	-0.08089 (13)	1.11449 (10)	0.0390 (5)
C6	0.19590 (11)	-0.01201 (12)	1.08006 (9)	0.0329 (4)
C7	0.39245 (11)	0.02418 (11)	1.06834 (9)	0.0266 (4)
C8	0.34628 (10)	0.08883 (11)	0.96964 (9)	0.0261 (3)
C9	0.24507 (11)	0.03874 (11)	0.88019 (9)	0.0283 (4)
C10	0.16409 (10)	0.12118 (11)	0.79933 (9)	0.0285 (4)
C11	0.10610 (11)	0.07481 (12)	0.69998 (9)	0.0341 (4)
C12	0.02797 (12)	0.14692 (14)	0.62317 (10)	0.0401 (5)
C13	0.00466 (12)	0.26359 (14)	0.64446 (10)	0.0431 (5)
C14	0.06158 (12)	0.31001 (13)	0.74235 (10)	0.0401 (5)
C15	0.14183 (11)	0.23937 (12)	0.81968 (10)	0.0322 (4)

C16	0.55535 (10)	0.32336 (11)	0.98885 (9)	0.0246 (4)
C17	0.49391 (10)	0.39449 (10)	0.90350 (9)	0.0260 (4)
C18	0.54880 (11)	0.49364 (11)	0.88215 (9)	0.0277 (4)
C19	0.66431 (11)	0.52350 (11)	0.94650 (9)	0.0272 (4)
C20	0.72508 (11)	0.45238 (11)	1.03236 (9)	0.0292 (4)
C21	0.67143 (11)	0.35237 (11)	1.05282 (9)	0.0282 (4)
C22	0.66773 (12)	0.69076 (11)	0.84126 (9)	0.0338 (4)
H1	0.5436 (12)	0.1775 (13)	1.0677 (11)	0.041 (4)*
H2	0.43120	-0.18360	1.16610	0.0360*
H3	0.30730	-0.30380	1.21620	0.0450*
H4	0.11640	-0.23720	1.18620	0.0490*
H5	0.04760	-0.05330	1.10370	0.0470*
H6	0.16650	0.06130	1.04400	0.0390*
H11	0.12040	-0.00600	0.68540	0.0410*
H12	-0.00980	0.11610	0.55550	0.0480*
H13	-0.05060	0.31210	0.59180	0.0520*
H14	0.04570	0.39040	0.75660	0.0480*
H15	0.18160	0.27180	0.88660	0.0390*
H17	0.41430	0.37510	0.85980	0.0310*
H18	0.50720	0.54150	0.82310	0.0330*
H20	0.80400	0.47290	1.07710	0.0350*
H21	0.71390	0.30320	1.11070	0.0340*
H22A	0.64540	0.63760	0.78130	0.0510*
H22B	0.72190	0.75360	0.83730	0.0510*
H22C	0.59640	0.72870	0.84260	0.0510*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0453 (6)	0.0278 (5)	0.0359 (5)	-0.0048 (4)	0.0103 (4)	0.0012 (4)
O2	0.0266 (5)	0.0370 (5)	0.0308 (5)	0.0004 (4)	0.0089 (4)	0.0049 (4)
O3	0.0341 (5)	0.0326 (5)	0.0361 (5)	-0.0065 (4)	0.0137 (4)	0.0022 (4)
N1	0.0258 (6)	0.0266 (6)	0.0273 (5)	0.0003 (4)	0.0105 (5)	-0.0019 (4)
N2	0.0252 (6)	0.0283 (6)	0.0248 (5)	-0.0007 (4)	0.0086 (5)	0.0016 (4)
C1	0.0305 (7)	0.0269 (7)	0.0237 (6)	-0.0019 (5)	0.0111 (5)	-0.0031 (5)
C2	0.0370 (7)	0.0303 (7)	0.0242 (6)	-0.0004 (5)	0.0139 (6)	-0.0023 (5)
C3	0.0543 (9)	0.0317 (7)	0.0291 (6)	-0.0062 (6)	0.0191 (6)	-0.0013 (6)
C4	0.0473 (9)	0.0464 (9)	0.0353 (7)	-0.0178 (7)	0.0228 (7)	-0.0053 (6)
C5	0.0316 (8)	0.0520 (9)	0.0355 (7)	-0.0072 (6)	0.0156 (6)	-0.0072 (6)
C6	0.0314 (7)	0.0344 (7)	0.0319 (7)	-0.0002 (6)	0.0117 (6)	-0.0010 (6)
C7	0.0278 (7)	0.0251 (6)	0.0279 (6)	0.0028 (5)	0.0122 (6)	-0.0016 (5)
C8	0.0263 (6)	0.0249 (6)	0.0280 (6)	0.0011 (5)	0.0119 (5)	0.0007 (5)
C9	0.0294 (7)	0.0289 (7)	0.0289 (6)	-0.0019 (5)	0.0142 (5)	0.0018 (5)
C10	0.0223 (6)	0.0343 (7)	0.0285 (6)	-0.0033 (5)	0.0099 (5)	0.0045 (5)
C11	0.0305 (7)	0.0371 (8)	0.0327 (7)	-0.0066 (6)	0.0107 (6)	0.0014 (6)
C12	0.0322 (8)	0.0541 (9)	0.0295 (7)	-0.0070 (6)	0.0076 (6)	0.0046 (6)
C13	0.0303 (8)	0.0575 (10)	0.0386 (8)	0.0069 (7)	0.0107 (6)	0.0189 (7)
C14	0.0362 (8)	0.0423 (8)	0.0449 (8)	0.0098 (6)	0.0194 (7)	0.0104 (6)

C15	0.0296 (7)	0.0372 (8)	0.0317 (7)	0.0017 (6)	0.0143 (6)	0.0037 (6)
C16	0.0261 (7)	0.0251 (7)	0.0251 (6)	0.0007 (5)	0.0128 (5)	-0.0022 (5)
C17	0.0236 (6)	0.0281 (7)	0.0251 (6)	0.0003 (5)	0.0086 (5)	-0.0032 (5)
C18	0.0296 (7)	0.0283 (7)	0.0255 (6)	0.0022 (5)	0.0113 (5)	0.0011 (5)
C19	0.0285 (7)	0.0265 (6)	0.0305 (6)	-0.0015 (5)	0.0158 (5)	-0.0036 (5)
C20	0.0228 (6)	0.0329 (7)	0.0300 (6)	0.0000 (5)	0.0086 (5)	-0.0030 (5)
C21	0.0278 (7)	0.0310 (7)	0.0244 (6)	0.0039 (5)	0.0090 (5)	0.0012 (5)
C22	0.0379 (8)	0.0295 (7)	0.0382 (7)	-0.0017 (6)	0.0197 (6)	0.0027 (6)

Geometric parameters (\AA , $^\circ$)

O1—C9	1.2280 (15)	C16—C21	1.3884 (19)
O2—C7	1.2368 (16)	C16—C17	1.3890 (17)
O3—C19	1.3743 (16)	C17—C18	1.3839 (18)
O3—C22	1.4311 (15)	C18—C19	1.3866 (19)
N1—N2	1.3062 (15)	C19—C20	1.3922 (17)
N1—C8	1.3259 (16)	C20—C21	1.3776 (18)
N2—C16	1.4120 (16)	C2—H2	0.9500
N2—H1	0.884 (14)	C3—H3	0.9500
C1—C6	1.391 (2)	C4—H4	0.9500
C1—C7	1.4988 (19)	C5—H5	0.9500
C1—C2	1.3893 (17)	C6—H6	0.9500
C2—C3	1.389 (2)	C11—H11	0.9500
C3—C4	1.381 (2)	C12—H12	0.9500
C4—C5	1.383 (2)	C13—H13	0.9500
C5—C6	1.389 (2)	C14—H14	0.9500
C7—C8	1.4725 (17)	C15—H15	0.9500
C8—C9	1.4853 (18)	C17—H17	0.9500
C9—C10	1.4927 (17)	C18—H18	0.9500
C10—C11	1.4012 (17)	C20—H20	0.9500
C10—C15	1.3892 (18)	C21—H21	0.9500
C11—C12	1.3826 (19)	C22—H22A	0.9800
C12—C13	1.381 (2)	C22—H22B	0.9800
C13—C14	1.3833 (19)	C22—H22C	0.9800
C14—C15	1.3872 (19)		
C19—O3—C22	116.93 (10)	C18—C19—C20	119.50 (12)
N2—N1—C8	121.26 (10)	C19—C20—C21	120.34 (12)
N1—N2—C16	120.15 (10)	C16—C21—C20	120.04 (11)
N1—N2—H1	119.9 (10)	C1—C2—H2	120.00
C16—N2—H1	119.7 (10)	C3—C2—H2	120.00
C2—C1—C6	119.63 (13)	C2—C3—H3	120.00
C2—C1—C7	118.74 (13)	C4—C3—H3	120.00
C6—C1—C7	121.55 (11)	C3—C4—H4	120.00
C1—C2—C3	120.37 (14)	C5—C4—H4	120.00
C2—C3—C4	119.68 (13)	C4—C5—H5	120.00
C3—C4—C5	120.28 (14)	C6—C5—H5	120.00
C4—C5—C6	120.25 (14)	C1—C6—H6	120.00

C1—C6—C5	119.74 (12)	C5—C6—H6	120.00
O2—C7—C8	120.42 (12)	C10—C11—H11	120.00
O2—C7—C1	119.64 (11)	C12—C11—H11	120.00
C1—C7—C8	119.80 (11)	C11—C12—H12	120.00
N1—C8—C9	114.35 (10)	C13—C12—H12	120.00
N1—C8—C7	124.47 (11)	C12—C13—H13	120.00
C7—C8—C9	120.47 (11)	C14—C13—H13	120.00
O1—C9—C10	120.72 (11)	C13—C14—H14	120.00
C8—C9—C10	120.27 (11)	C15—C14—H14	120.00
O1—C9—C8	118.95 (11)	C10—C15—H15	120.00
C11—C10—C15	119.45 (12)	C14—C15—H15	120.00
C9—C10—C11	117.85 (11)	C16—C17—H17	120.00
C9—C10—C15	122.66 (11)	C18—C17—H17	120.00
C10—C11—C12	119.89 (12)	C17—C18—H18	120.00
C11—C12—C13	120.29 (12)	C19—C18—H18	120.00
C12—C13—C14	120.13 (13)	C19—C20—H20	120.00
C13—C14—C15	120.15 (13)	C21—C20—H20	120.00
C10—C15—C14	120.06 (12)	C16—C21—H21	120.00
C17—C16—C21	119.89 (11)	C20—C21—H21	120.00
N2—C16—C17	121.54 (11)	O3—C22—H22A	109.00
N2—C16—C21	118.56 (11)	O3—C22—H22B	109.00
C16—C17—C18	119.91 (12)	O3—C22—H22C	109.00
C17—C18—C19	120.30 (11)	H22A—C22—H22B	110.00
O3—C19—C20	115.97 (11)	H22A—C22—H22C	109.00
O3—C19—C18	124.52 (11)	H22B—C22—H22C	109.00
C22—O3—C19—C18	-4.76 (18)	C7—C8—C9—O1	-30.36 (19)
C22—O3—C19—C20	175.85 (11)	C7—C8—C9—C10	152.35 (12)
C8—N1—N2—C16	-177.67 (11)	O1—C9—C10—C11	-24.5 (2)
N2—N1—C8—C7	4.91 (19)	O1—C9—C10—C15	153.45 (14)
N2—N1—C8—C9	-165.34 (11)	C8—C9—C10—C11	152.71 (13)
N1—N2—C16—C17	6.82 (18)	C8—C9—C10—C15	-29.3 (2)
N1—N2—C16—C21	-172.50 (12)	C9—C10—C11—C12	178.39 (13)
C6—C1—C2—C3	2.05 (18)	C15—C10—C11—C12	0.3 (2)
C7—C1—C2—C3	178.65 (11)	C9—C10—C15—C14	-177.10 (13)
C2—C1—C6—C5	0.13 (18)	C11—C10—C15—C14	0.9 (2)
C7—C1—C6—C5	-176.37 (11)	C10—C11—C12—C13	-1.5 (2)
C2—C1—C7—O2	-41.22 (17)	C11—C12—C13—C14	1.6 (2)
C2—C1—C7—C8	143.16 (12)	C12—C13—C14—C15	-0.4 (2)
C6—C1—C7—O2	135.31 (13)	C13—C14—C15—C10	-0.9 (2)
C6—C1—C7—C8	-40.31 (17)	N2—C16—C17—C18	-178.94 (12)
C1—C2—C3—C4	-2.43 (19)	C21—C16—C17—C18	0.37 (19)
C2—C3—C4—C5	0.6 (2)	N2—C16—C21—C20	-179.88 (12)
C3—C4—C5—C6	1.6 (2)	C17—C16—C21—C20	0.79 (19)
C4—C5—C6—C1	-1.95 (19)	C16—C17—C18—C19	-1.03 (19)
O2—C7—C8—N1	-16.2 (2)	C17—C18—C19—O3	-178.83 (12)
O2—C7—C8—C9	153.45 (12)	C17—C18—C19—C20	0.53 (19)
C1—C7—C8—N1	159.35 (12)	O3—C19—C20—C21	-179.95 (12)

C1—C7—C8—C9	−30.96 (18)	C18—C19—C20—C21	0.6 (2)
N1—C8—C9—O1	140.32 (13)	C19—C20—C21—C16	−1.3 (2)
N1—C8—C9—C10	−36.97 (18)		

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
N2—H1···O2	0.884 (14)	1.977 (15)	2.6427 (14)	131.0 (14)
C15—H15···O3 ⁱ	0.95	2.64	3.5859 (16)	174
C21—H21···O1 ⁱⁱ	0.95	2.63	3.3441 (15)	132

Symmetry codes: (i) $-x+1, -y+1, -z+2$; (ii) $-x+1, -y, -z+2$.