

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Pyridine-2,3-diamine

Richard Betz,* Thomas Gerber, Eric Hosten and Henk Schalekamp

Nelson Mandela Metropolitan University, Summerstrand Campus, Department of Chemistry, University Way, Summerstrand, PO Box 77000, Port Elizabeth 6031, South Africa

Correspondence e-mail: richard.betz@webmail.co.za


Received 12 July 2011; accepted 20 July 2011

Key indicators: single-crystal X-ray study; $T = 200$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.036; wR factor = 0.085; data-to-parameter ratio = 8.5.

The molecule of the title pyridine derivative, $\text{C}_5\text{H}_7\text{N}_3$, shows approximately non-crystallographic C_s symmetry. Intracyclic angles cover the range 117.50 (14)– 123.03 (15)°. In the crystal, $\text{N}-\text{H}\cdots\text{N}$ hydrogen bonds connect molecules into a three-dimensional network. The closest intercentroid distance between two π -systems occurs with the c -axis repeat at 3.9064 (12) Å.

Related literature

For the crystal structure of the dihydrochloride of the title compound, see: Hemamalini & Fun (2010). For the crystal structures of Zn complexes of the title compound, see: de Cires-Mejias *et al.* (2004). For graph-set analysis of hydrogen bonds, see: Etter *et al.* (1990); Bernstein *et al.* (1995).



Experimental

Crystal data

$\text{C}_5\text{H}_7\text{N}_3$
 $M_r = 109.14$
 Tetragonal, $P4_2bc$
 $a = 16.4670$ (3) Å

$c = 3.9064$ (12) Å
 $V = 1059.3$ (3) Å³
 $Z = 8$
 Mo $K\alpha$ radiation

$\mu = 0.09$ mm⁻¹
 $T = 200$ K

$0.48 \times 0.16 \times 0.11$ mm

Data collection

Bruker APEXII CCD
 diffractometer
 9864 measured reflections

754 independent reflections
 706 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.047$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.036$
 $wR(F^2) = 0.085$
 $S = 1.10$
 754 reflections
 89 parameters
 1 restraint

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.23$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.17$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N2}-\text{H21}\cdots\text{N1}^{\text{i}}$	0.87 (2)	2.32 (2)	3.153 (2)	161.2 (19)
$\text{N2}-\text{H22}\cdots\text{N2}^{\text{ii}}$	0.85 (2)	2.58 (2)	3.4369 (16)	175.9 (18)
$\text{N3}-\text{H31}\cdots\text{N1}^{\text{iii}}$	0.86 (2)	2.32 (2)	3.115 (2)	156 (2)
$\text{N3}-\text{H32}\cdots\text{N3}^{\text{iv}}$	0.89 (3)	2.47 (2)	3.359 (2)	175 (2)

Symmetry codes: (i) $-y, x, z + \frac{1}{2}$; (ii) $y, -x, z + \frac{1}{2}$; (iii) $-y, x, z - \frac{1}{2}$; (iv) $-y + \frac{1}{2}, -x + \frac{1}{2}, z - \frac{1}{2}$.

Data collection: *APEX2* (Bruker, 2010); cell refinement: *SAINT* (Bruker, 2010); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *Mercury* (Macrae *et al.*, 2008); software used to prepare material for publication: *SHELXL97* and *PLATON* (Spek, 2009).

The authors thank Mrs Phyllis Atkinson for helpful discussions.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: OM2449).

References

- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2010). *APEX2* and *SAINT* Bruker AXS Inc., Madison, Wisconsin, USA.
- Cires-Mejias, C. de, Tanase, S., Reedijk, J., Gonzalez-Vilchez, F., Vilaplana, R., Mills, A. M., Kooijman, H. & Spek, A. L. (2004). *Inorg. Chim. Acta*, **357**, 1494–1498.
- Etter, M. C., MacDonald, J. C. & Bernstein, J. (1990). *Acta Cryst.* **B46**, 256–262.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Hemamalini, M. & Fun, H.-K. (2010). *Acta Cryst.* **E66**, o513–o514.
- Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

supporting information

Acta Cryst. (2011). E67, o2154 [doi:10.1107/S1600536811029412]

Pyridine-2,3-diamine

Richard Betz, Thomas Gerber, Eric Hosten and Henk Schalekamp

S1. Comment

Chelate ligands have found widespread use in coordination chemistry due to the enhanced thermodynamic stability of resultant coordination compounds in relation to coordination compounds exclusively applying comparable monodentate ligands. Combining identical donor atoms in different states of hybridization, a molecular set-up to accommodate a large variety of metal centers of variable Lewis acidity is at hand. In this aspect, the title compound seemed interesting due to its use as strictly neutral or – depending on the pH value – as anionic or cationic ligand. Furthermore, thanks to the presence of three possible donor atoms, the title compound might serve as a building block in the formation of metal-organic framework structures. For the title compound, two zinc-supported polymers have been reported whose crystal structure analysis shows the absence of chelate-type building motifs (de Cires-Mejias *et al.*, 2004). At the beginning of a more comprehensive study to elucidate the formation of coordination polymers exclusively featuring nitrogen-containing ligands, we determined the structure of the title compound to enable comparative studies of metrical parameters in envisioned coordination compounds. Information about the molecular and crystal structure of the dihydrochloride of the title compound is apparent in the literature (Hemamalini & Fun, 2010).

Intracyclic angles cover a range of 117.50 (14)–123.03 (15) ° with the smallest angle found on the carbon atom bearing the amino group in *meta* position to the intracyclic nitrogen atom and the biggest angle found on the carbon atom bearing a hydrogen atom in *ortho* position to the intracyclic nitrogen atom. Apart from the hydrogen atoms of the amino groups which point to opposite sides of the plane defined the aromatic system, all atoms are essentially residing in one common plane (r.m.s. deviation of all fitted non-hydrogen atoms = 0.0152 Å). The amino groups are not planar, the least-squares planes defined by the NH₂ groups subtend angles of 40.2 (2) ° and 79.5 (2) ° with the least-squares plane defined by the atoms of the heterocycle (Fig. 1).

The crystal structure of the title compound is marked by a hydrogen bonding system involving all hydrogen atoms of both amino groups as donors and the intracyclic as well as the exocyclic nitrogen atoms as acceptors. The intracyclic nitrogen atom serves as a twofold acceptor for one of the hydrogen atoms of each of the two different amino groups. The remaining hydrogen atom on each amino group gives rise to a cooperative chain of hydrogen bonds, respectively. The latter ones are antiprotonic. In terms of graph-set analysis (Etter *et al.*, 1990; Bernstein *et al.*, 1995), the descriptor for this hydrogen bonding system on the unitary level is C¹₁(2)C¹₁(2)C¹₁(4)C¹₁(5). In total, the molecules are connected to a three-dimensional network (Fig. 2). The closest intercentroid distance between two aromatic systems follows the c-axis repeat at 3.9064 (12) Å.

The packing of the title compound is shown in Figure 3.

S2. Experimental

The compound was obtained commercially (Aldrich). Crystals suitable for the X-ray diffraction study were taken directly from the provided compound.

S3. Refinement

Carbon-bound H atoms were placed in calculated positions (C—H 0.95 Å) and were included in the refinement in the riding model approximation, with $U(\text{H})$ set to $1.2U_{\text{eq}}(\text{C})$. The H atoms of the amine groups were located on a difference Fourier map and refined with individual thermal parameters. Due to the absence of a strong anomalous scatterer, the Flack parameter is meaningless. Thus, Friedel opposites (2407 pairs) have been merged and the item was removed from the CIF.






Figure 1

The molecular structure of the title compound, with atom labels and anisotropic displacement ellipsoids (drawn at 50% probability level).

**Figure 2**

Intermolecular contacts, viewed approximately along [0 0 1]. Symmetry operators: ⁱ $y, -x, z - 1/2$; ⁱⁱ $y, -x, z + 1/2$; ⁱⁱⁱ $-y, x, z - 1/2$; ^{iv} $-y, x, z + 1/2$; ^v $-y + 1/2, -x + 1/2, z - 1/2$; ^{vi} $-y + 1/2, -x + 1/2, z + 1/2$.

**Figure 3**

Molecular packing of the title compound, viewed along $[0\ 0\ -1]$ (anisotropic displacement ellipsoids drawn at 50% probability level).

Pyridine-2,3-diamine

Crystal data

$C_5H_7N_3$

$M_r = 109.14$

Tetragonal, $P4_2bc$

Hall symbol: $P\ 4c\ -2ab$

$a = 16.4670\ (3)\ \text{\AA}$

$c = 3.9064\ (12)\ \text{\AA}$

$V = 1059.3\ (3)\ \text{\AA}^3$

$Z = 8$

$F(000) = 464$

$D_x = 1.369\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 5917 reflections

$\theta = 2.5\text{--}28.3^\circ$

$\mu = 0.09\ \text{mm}^{-1}$

$T = 200\ \text{K}$

Needle, brown

$0.48 \times 0.16 \times 0.11\ \text{mm}$

Data collection

Bruker APEXII CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

φ and ω scans

9864 measured reflections

754 independent reflections

706 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.047$
 $\theta_{\text{max}} = 28.3^\circ$, $\theta_{\text{min}} = 1.8^\circ$
 $h = -21 \rightarrow 20$

$k = -21 \rightarrow 21$
 $l = -5 \rightarrow 5$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.036$
 $wR(F^2) = 0.085$
 $S = 1.10$
 754 reflections
 89 parameters
 1 restraint
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H atoms treated by a mixture of independent
 and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0428P)^2 + 0.2489P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} < 0.001$
 $\Delta\rho_{\text{max}} = 0.23 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.17 \text{ e } \text{\AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.20596 (8)	-0.05610 (8)	0.9243 (5)	0.0268 (3)
N2	0.11296 (8)	0.04458 (9)	1.0621 (5)	0.0271 (3)
H21	0.1094 (13)	0.0921 (14)	1.156 (8)	0.040 (6)*
H22	0.0934 (12)	0.0063 (14)	1.184 (7)	0.037 (6)*
N3	0.21809 (9)	0.16454 (9)	0.7974 (5)	0.0302 (4)
H31	0.1677 (13)	0.1730 (12)	0.757 (7)	0.033 (5)*
H32	0.2471 (12)	0.1946 (12)	0.652 (8)	0.038 (6)*
C1	0.18758 (9)	0.02266 (9)	0.9286 (5)	0.0225 (4)
C2	0.23965 (9)	0.08211 (9)	0.7834 (6)	0.0240 (3)
C3	0.31167 (10)	0.05570 (10)	0.6425 (5)	0.0283 (4)
H3	0.3478	0.0938	0.5417	0.034*
C4	0.33173 (10)	-0.02631 (11)	0.6469 (6)	0.0307 (4)
H4	0.3819	-0.0449	0.5556	0.037*
C5	0.27702 (10)	-0.07971 (10)	0.7870 (6)	0.0300 (4)
H5	0.2900	-0.1359	0.7869	0.036*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0256 (7)	0.0250 (7)	0.0298 (8)	0.0002 (5)	-0.0024 (7)	0.0000 (7)
N2	0.0250 (7)	0.0249 (7)	0.0315 (8)	-0.0003 (5)	0.0025 (6)	-0.0013 (7)
N3	0.0276 (7)	0.0256 (7)	0.0374 (9)	-0.0023 (5)	-0.0009 (8)	0.0039 (8)
C1	0.0214 (7)	0.0252 (7)	0.0209 (8)	-0.0013 (5)	-0.0042 (7)	-0.0005 (7)
C2	0.0238 (7)	0.0267 (7)	0.0217 (7)	-0.0032 (5)	-0.0040 (7)	0.0003 (8)
C3	0.0261 (8)	0.0357 (8)	0.0231 (8)	-0.0065 (6)	-0.0014 (8)	0.0009 (8)
C4	0.0242 (7)	0.0411 (9)	0.0268 (9)	0.0024 (6)	0.0012 (8)	-0.0044 (9)
C5	0.0288 (8)	0.0278 (8)	0.0334 (9)	0.0045 (6)	-0.0023 (10)	-0.0021 (9)

Geometric parameters (Å, °)

N1—C1	1.332 (2)	C1—C2	1.420 (2)
N1—C5	1.345 (2)	C2—C3	1.378 (2)
N2—C1	1.383 (2)	C3—C4	1.390 (2)
N2—H21	0.87 (2)	C3—H3	0.9500
N2—H22	0.85 (2)	C4—C5	1.373 (3)
N3—C2	1.404 (2)	C4—H4	0.9500
N3—H31	0.86 (2)	C5—H5	0.9500
N3—H32	0.89 (3)		
C1—N1—C5	118.97 (15)	C3—C2—C1	117.50 (14)
C1—N2—H21	117.1 (15)	N3—C2—C1	119.89 (16)
C1—N2—H22	110.7 (14)	C2—C3—C4	120.41 (16)
H21—N2—H22	114 (2)	C2—C3—H3	119.8
C2—N3—H31	113.3 (13)	C4—C3—H3	119.8
C2—N3—H32	112.2 (14)	C5—C4—C3	118.11 (16)
H31—N3—H32	108 (2)	C5—C4—H4	120.9
N1—C1—N2	117.44 (15)	C3—C4—H4	120.9
N1—C1—C2	121.95 (15)	N1—C5—C4	123.03 (15)
N2—C1—C2	120.49 (14)	N1—C5—H5	118.5
C3—C2—N3	122.57 (16)	C4—C5—H5	118.5
C5—N1—C1—N2	178.05 (17)	N3—C2—C3—C4	177.43 (19)
C5—N1—C1—C2	1.9 (3)	C1—C2—C3—C4	-0.4 (3)
N1—C1—C2—C3	-1.4 (3)	C2—C3—C4—C5	1.6 (3)
N2—C1—C2—C3	-177.42 (18)	C1—N1—C5—C4	-0.6 (3)
N1—C1—C2—N3	-179.3 (2)	C3—C4—C5—N1	-1.1 (3)
N2—C1—C2—N3	4.7 (3)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N2—H21...N1 ⁱ	0.87 (2)	2.32 (2)	3.153 (2)	161.2 (19)
N2—H22...N2 ⁱⁱ	0.85 (2)	2.58 (2)	3.4369 (16)	175.9 (18)
N3—H31...N1 ⁱⁱⁱ	0.86 (2)	2.32 (2)	3.115 (2)	156 (2)
N3—H32...N3 ^{iv}	0.89 (3)	2.47 (2)	3.359 (2)	175 (2)

Symmetry codes: (i) $-y, x, z+1/2$; (ii) $y, -x, z+1/2$; (iii) $-y, x, z-1/2$; (iv) $-y+1/2, -x+1/2, z-1/2$.