

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

exo-4-[(1*H*-Benzimidazol-2-yl)methyl]-10-oxa-4-azatricyclo[5.2.1.0^{2,6}]decane-3,5-dione

Shi-Kun Li,^{a,b} Fan Zhang,^{a,b} Tian-Xi Lv^b and Qiu-Yue Lin^{a,b*}

^aZhejiang Key Laboratory for Reactive Chemistry on Solid Surfaces, Institute of Physical Chemistry, Zhejiang Normal University, Jinhua 321004, Zhejiang, People's Republic of China, and ^bCollege of Chemistry and Life Science, Zhejiang Normal University, Jinhua 321004, Zhejiang, People's Republic of China
Correspondence e-mail: sky51@zjnu.cn


Received 29 April 2011; accepted 4 July 2011

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.035; wR factor = 0.087; data-to-parameter ratio = 10.8.

In the title compound, $\text{C}_{16}\text{H}_{15}\text{N}_3\text{O}_3$, the dihedral angle between the approximately planar benzimidazolyl group (r.m.s. deviation = 0.010 Å) and the pyrrolidine ring is 78.20 (6)°. The C—C—N bond angle of the bridging CH_2 group is 112.14 (16)°. In the crystal, molecules are linked *via* N—H...N hydrogen bonds, forming infinite chains parallel to [101] and [10 $\bar{1}$].

Related literature

For the bioactivity of norcantharidin (systematic name 4,10-dioxatricyclo[5.2.1.0^{2,6}]decane-3,5-dione), see: Wang (1989). For the use of norcantharidin in synthesis, see: Hill *et al.* (2007). For a related structure, see: Zhu & Lin (2009).



Experimental

Crystal data

 $\text{C}_{16}\text{H}_{15}\text{N}_3\text{O}_3$ $M_r = 297.31$ Orthorhombic, $Fdd2$ $a = 17.4294$ (2) Å $b = 48.2746$ (6) Å $c = 6.7947$ (1) Å $V = 5717.04$ (13) Å³ $Z = 16$ Mo $K\alpha$ radiation $\mu = 0.10$ mm⁻¹ $T = 296$ K $0.24 \times 0.17 \times 0.10$ mm

Data collection

Bruker SMART APEXII CCD diffractometer

Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996) $T_{\min} = 0.981$, $T_{\max} = 0.990$

22741 measured reflections

2159 independent reflections

1690 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.038$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.035$ $wR(F^2) = 0.087$ $S = 1.04$

2159 reflections

199 parameters

1 restraint

H-atom parameters constrained

 $\Delta\rho_{\max} = 0.10$ e Å⁻³ $\Delta\rho_{\min} = -0.17$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1A}\cdots\text{N2}^i$	0.86	1.97	2.827 (2)	175

Symmetry code: (i) $x + \frac{1}{4}, -y + \frac{1}{4}, z + \frac{1}{4}$.

Data collection: *APEX2* (Bruker, 2006); cell refinement: *SAINT* (Bruker, 2006); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

The authors thank for the financial support from the Natural Science Foundation of Zhejiang Province, China (Grant No. Y407301).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: QK2008).

References

- Bruker (2006). *SAINT* and *SMART*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Hill, T. A., Stewart, S. G., Ackland, S. P., Sauer, B., Gilbert, J., Ackland, S. P., Sakoff, J. A. & McCluskey, A. (2007). *Bioorg. Med. Chem.* **15**, 6126–6134.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Wang, G.-S. (1989). *J. Ethnopharmacol.* **26**, 147–162.
- Zhu, W.-Z. & Lin, Q.-Y. (2009). *Acta Cryst.* **E65**, o287.

supporting information

Acta Cryst. (2011). E67, o1974 [doi:10.1107/S1600536811026602]

**exo-4-[(1*H*-Benzimidazol-2-yl)methyl]-10-oxa-4-azatricyclo-
[5.2.1.0^{2,6}]decane-3,5-dione****Shi-Kun Li, Fan Zhang, Tian-Xi Lv and Qiu-Yue Lin****S1. Comment**

Norcantharidin is an important compound that has a variety of bioactivities, such as protein kinase inhibition and antitumor activity (Wang, 1989). As a contribution to structure-activity studies in this field, the title compound, (I), a norcantharidin imide (Hill *et al.*, 2007), was synthesized and investigated by X-ray crystallography. A related norcantharidin imide was reported by Zhu & Lin (2009).


X-ray crystallography confirmed the anticipated molecular structure and the atom connectivity for the title compound, as illustrated in Fig. 1. The norcantharidin imide and the benzimidazole moiety of (I) are rigid and feature usual bond lengths and angles. The dihedral angle between the flat benzimidazolyl group (C9—C15,N1,N2) and the pyrrolidine ring (C1,C2,C7,C8,N3) is 78.20 (6)°. The bond angle of the bridging CH₂ group is C15—C16—N3 = 112.1 (2)°. In the crystal structure the molecules are linked *via* the hydrogen bond N1—H1A···N2ⁱ, N1···N2ⁱ = 2.827 (2) Å ((i): x+1/4, -y+1/4, z+1/4), to form infinite hydrogen bonded chains parallel to [101] and [10 $\bar{1}$].

S2. Experimental

A mixture of 0.5 mmol norcantharidin, 0.5 mmol 2-(aminomethyl)benzimidazole, 0.5 mmol cadmium acetate as a promoter, and 10 mL distilled water was sealed in a 25 mL Teflon-lined stainless vessel and heated at 433 K for 3 d, then cooled slowly to room temperature. The solution was filtered and gave then block-like transparent crystals.

S3. Refinement

The structure was solved by direct methods and successive Fourier difference synthesis. The H atoms were positioned geometrically and refined using a riding model [aromatic C—H = 0.93 Å, aliphatic C—H = 0.97 Å, and N—H = 0.86 Å, $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}, \text{N})$]. In the absence of significant anomalous scattering, the Friedel pairs of the polar structure were merged.

**Figure 1**

A view of the molecule of (I) showing the atom-labelling scheme with displacement ellipsoids drawn at 30% probability.

exo-8-[(1*H*-Benzimidazol-2-yl)methyl]-10-oxa-8-azatricyclo[4.3.0.1^{2,5}]decane-7,9-dione

Crystal data

$C_{16}H_{15}N_3O_3$

$M_r = 297.31$

Orthorhombic, *Fdd2*

Hall symbol: *F* 2 -2*d*

$a = 17.4294$ (2) Å

$b = 48.2746$ (6) Å

$c = 6.7947$ (1) Å

$V = 5717.04$ (13) Å³

$Z = 16$

$F(000) = 2496$

$D_x = 1.382$ Mg m⁻³

Mo *K*α radiation, $\lambda = 0.71073$ Å

Cell parameters from 5637 reflections

$\theta = 2.5$ – 29.8°

$\mu = 0.10$ mm⁻¹

$T = 296$ K

Block, colourless

$0.24 \times 0.17 \times 0.10$ mm

Data collection

Bruker SMART APEXII CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

ω scans

Absorption correction: multi-scan

(*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.981$, $T_{\max} = 0.990$

22741 measured reflections

2159 independent reflections

1690 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.038$

$\theta_{\max} = 29.8^\circ$, $\theta_{\min} = 2.5^\circ$

$h = -24 \rightarrow 24$

$k = -67 \rightarrow 66$

$l = -9 \rightarrow 8$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.035$

$wR(F^2) = 0.087$

$S = 1.04$

2159 reflections

199 parameters

1 restraint

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0448P)^2 + 1.1097P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.10$ e Å⁻³

$\Delta\rho_{\min} = -0.17$ e Å⁻³

Absolute structure: 1753 Friedel pairs were merged

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
N2	-0.06276 (8)	0.10833 (3)	0.2792 (3)	0.0454 (4)
N1	0.04528 (8)	0.11596 (3)	0.4450 (3)	0.0428 (4)
H1A	0.0893	0.1229	0.4740	0.051*
N3	-0.02809 (8)	0.16452 (3)	0.1037 (3)	0.0431 (4)
O2	-0.07386 (9)	0.14915 (3)	-0.1930 (3)	0.0629 (4)
O1	-0.00312 (11)	0.18963 (4)	0.3811 (3)	0.0765 (5)
O3	-0.02560 (7)	0.22474 (3)	-0.0973 (2)	0.0556 (4)
C1	-0.07375 (10)	0.16595 (3)	-0.0624 (3)	0.0439 (4)
C2	-0.12089 (10)	0.19215 (3)	-0.0513 (3)	0.0431 (4)
H2B	-0.1761	0.1887	-0.0634	0.052*
C3	-0.09207 (12)	0.21403 (4)	-0.1970 (3)	0.0527 (5)
H3A	-0.0813	0.2068	-0.3288	0.063*
C4	-0.14630 (14)	0.23883 (4)	-0.1956 (5)	0.0713 (7)
H4A	-0.1995	0.2330	-0.2016	0.086*
H4B	-0.1357	0.2513	-0.3041	0.086*
C5	-0.12741 (14)	0.25219 (4)	0.0024 (4)	0.0700 (7)
H5A	-0.1088	0.2710	-0.0139	0.084*
H5B	-0.1716	0.2523	0.0892	0.084*
C6	-0.06469 (12)	0.23307 (4)	0.0784 (4)	0.0552 (5)
H6A	-0.0311	0.2417	0.1763	0.066*
C7	-0.09930 (11)	0.20533 (4)	0.1463 (3)	0.0464 (5)
H7A	-0.1434	0.2077	0.2341	0.056*
C8	-0.03878 (12)	0.18644 (4)	0.2306 (3)	0.0492 (5)
C16	0.02831 (10)	0.14281 (4)	0.1375 (4)	0.0519 (5)
H16A	0.0760	0.1511	0.1814	0.062*
H16B	0.0384	0.1333	0.0145	0.062*
C15	0.00198 (10)	0.12236 (3)	0.2871 (3)	0.0410 (4)
C9	-0.06167 (9)	0.09153 (3)	0.4452 (3)	0.0391 (4)
C10	-0.11496 (11)	0.07221 (4)	0.5106 (3)	0.0488 (5)
H10A	-0.1597	0.0687	0.4402	0.059*
C11	-0.09925 (12)	0.05847 (4)	0.6828 (4)	0.0571 (5)
H11A	-0.1339	0.0453	0.7288	0.068*
C12	-0.03280 (13)	0.06372 (5)	0.7903 (4)	0.0600 (6)
H12A	-0.0247	0.0543	0.9078	0.072*
C13	0.02144 (11)	0.08263 (4)	0.7270 (3)	0.0526 (5)

H13A	0.0660	0.0861	0.7984	0.063*
C14	0.00597 (10)	0.09617 (3)	0.5518 (3)	0.0395 (4)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N2	0.0332 (7)	0.0520 (8)	0.0510 (10)	-0.0018 (6)	-0.0111 (7)	0.0068 (8)
N1	0.0308 (7)	0.0445 (7)	0.0532 (10)	-0.0010 (6)	-0.0124 (8)	-0.0003 (7)
N3	0.0394 (8)	0.0408 (7)	0.0490 (10)	0.0022 (6)	-0.0058 (7)	0.0065 (7)
O2	0.0768 (10)	0.0467 (7)	0.0652 (11)	0.0086 (7)	-0.0172 (9)	-0.0142 (8)
O1	0.0876 (11)	0.0950 (13)	0.0468 (11)	0.0081 (9)	-0.0214 (10)	-0.0072 (9)
O3	0.0472 (7)	0.0489 (7)	0.0708 (11)	-0.0026 (6)	0.0051 (8)	0.0100 (7)
C1	0.0451 (9)	0.0351 (8)	0.0514 (12)	-0.0021 (7)	-0.0074 (9)	0.0015 (8)
C2	0.0380 (9)	0.0387 (8)	0.0525 (11)	0.0014 (7)	-0.0062 (9)	-0.0002 (8)
C3	0.0624 (12)	0.0469 (10)	0.0489 (13)	0.0087 (9)	-0.0070 (11)	0.0047 (9)
C4	0.0740 (15)	0.0457 (10)	0.094 (2)	0.0098 (10)	-0.0194 (15)	0.0099 (13)
C5	0.0703 (13)	0.0407 (9)	0.099 (2)	0.0100 (10)	0.0017 (15)	-0.0043 (12)
C6	0.0576 (12)	0.0427 (9)	0.0654 (15)	-0.0041 (8)	-0.0085 (11)	-0.0086 (10)
C7	0.0447 (10)	0.0481 (9)	0.0466 (12)	0.0029 (8)	0.0035 (9)	-0.0034 (9)
C8	0.0493 (11)	0.0568 (11)	0.0414 (12)	-0.0011 (9)	-0.0015 (10)	0.0022 (9)
C16	0.0386 (9)	0.0511 (10)	0.0660 (15)	0.0073 (8)	-0.0002 (10)	0.0147 (10)
C15	0.0324 (8)	0.0416 (8)	0.0490 (12)	0.0040 (7)	-0.0080 (8)	0.0041 (8)
C9	0.0334 (8)	0.0393 (8)	0.0444 (11)	0.0033 (6)	-0.0079 (8)	-0.0022 (8)
C10	0.0394 (9)	0.0502 (10)	0.0570 (14)	-0.0041 (8)	-0.0060 (9)	-0.0017 (9)
C11	0.0535 (11)	0.0506 (10)	0.0671 (15)	-0.0001 (8)	0.0027 (12)	0.0114 (11)
C12	0.0610 (12)	0.0633 (12)	0.0558 (14)	0.0089 (10)	-0.0030 (11)	0.0175 (11)
C13	0.0470 (11)	0.0628 (11)	0.0480 (12)	0.0076 (9)	-0.0156 (10)	0.0025 (10)
C14	0.0333 (8)	0.0388 (8)	0.0463 (12)	0.0042 (7)	-0.0079 (8)	-0.0021 (8)

Geometric parameters (Å, °)

N2—C15	1.317 (2)	C5—C6	1.521 (3)
N2—C9	1.389 (2)	C5—H5A	0.9700
N1—C15	1.348 (3)	C5—H5B	0.9700
N1—C14	1.381 (2)	C6—C7	1.540 (3)
N1—H1A	0.8600	C6—H6A	0.9800
N3—C8	1.378 (3)	C7—C8	1.507 (3)
N3—C1	1.383 (3)	C7—H7A	0.9800
N3—C16	1.455 (2)	C16—C15	1.490 (3)
O2—C1	1.202 (2)	C16—H16A	0.9700
O1—C8	1.206 (3)	C16—H16B	0.9700
O3—C6	1.432 (3)	C9—C10	1.389 (3)
O3—C3	1.438 (3)	C9—C14	1.402 (2)
C1—C2	1.510 (2)	C10—C11	1.373 (3)
C2—C3	1.532 (3)	C10—H10A	0.9300
C2—C7	1.533 (3)	C11—C12	1.393 (3)
C2—H2B	0.9800	C11—H11A	0.9300
C3—C4	1.525 (3)	C12—C13	1.383 (3)

C3—H3A	0.9800	C12—H12A	0.9300
C4—C5	1.528 (4)	C13—C14	1.385 (3)
C4—H4A	0.9700	C13—H13A	0.9300
C4—H4B	0.9700		
C15—N2—C9	104.80 (15)	C5—C6—H6A	113.7
C15—N1—C14	107.39 (14)	C7—C6—H6A	113.7
C15—N1—H1A	126.3	C8—C7—C2	104.70 (15)
C14—N1—H1A	126.3	C8—C7—C6	111.46 (16)
C8—N3—C1	113.23 (15)	C2—C7—C6	101.23 (17)
C8—N3—C16	123.08 (17)	C8—C7—H7A	112.9
C1—N3—C16	123.63 (17)	C2—C7—H7A	112.9
C6—O3—C3	96.35 (15)	C6—C7—H7A	112.9
O2—C1—N3	124.71 (16)	O1—C8—N3	124.0 (2)
O2—C1—C2	126.96 (19)	O1—C8—C7	127.3 (2)
N3—C1—C2	108.33 (16)	N3—C8—C7	108.75 (17)
C1—C2—C3	111.50 (17)	N3—C16—C15	112.14 (16)
C1—C2—C7	104.94 (16)	N3—C16—H16A	109.2
C3—C2—C7	101.50 (14)	C15—C16—H16A	109.2
C1—C2—H2B	112.7	N3—C16—H16B	109.2
C3—C2—H2B	112.7	C15—C16—H16B	109.2
C7—C2—H2B	112.7	H16A—C16—H16B	107.9
O3—C3—C4	102.38 (16)	N2—C15—N1	113.22 (17)
O3—C3—C2	101.96 (16)	N2—C15—C16	125.23 (18)
C4—C3—C2	109.49 (19)	N1—C15—C16	121.52 (16)
O3—C3—H3A	113.9	C10—C9—N2	129.97 (16)
C4—C3—H3A	113.9	C10—C9—C14	120.28 (17)
C2—C3—H3A	113.9	N2—C9—C14	109.74 (15)
C3—C4—C5	101.72 (19)	C11—C10—C9	117.63 (18)
C3—C4—H4A	111.4	C11—C10—H10A	121.2
C5—C4—H4A	111.4	C9—C10—H10A	121.2
C3—C4—H4B	111.4	C10—C11—C12	121.7 (2)
C5—C4—H4B	111.4	C10—C11—H11A	119.2
H4A—C4—H4B	109.3	C12—C11—H11A	119.2
C6—C5—C4	101.42 (18)	C13—C12—C11	121.7 (2)
C6—C5—H5A	111.5	C13—C12—H12A	119.1
C4—C5—H5A	111.5	C11—C12—H12A	119.1
C6—C5—H5B	111.5	C12—C13—C14	116.48 (19)
C4—C5—H5B	111.5	C12—C13—H13A	121.8
H5A—C5—H5B	109.3	C14—C13—H13A	121.8
O3—C6—C5	103.2 (2)	N1—C14—C13	132.96 (17)
O3—C6—C7	101.06 (15)	N1—C14—C9	104.84 (16)
C5—C6—C7	110.35 (16)	C13—C14—C9	122.19 (18)
O3—C6—H6A	113.7		
C8—N3—C1—O2	179.8 (2)	C16—N3—C8—O1	-0.5 (3)
C16—N3—C1—O2	2.6 (3)	C1—N3—C8—C7	0.7 (2)
C8—N3—C1—C2	0.9 (2)	C16—N3—C8—C7	177.93 (16)

C16—N3—C1—C2	-176.28 (16)	C2—C7—C8—O1	176.4 (2)
O2—C1—C2—C3	-71.9 (2)	C6—C7—C8—O1	67.8 (3)
N3—C1—C2—C3	106.99 (19)	C2—C7—C8—N3	-2.0 (2)
O2—C1—C2—C7	179.06 (19)	C6—C7—C8—N3	-110.60 (19)
N3—C1—C2—C7	-2.10 (19)	C8—N3—C16—C15	77.7 (2)
C6—O3—C3—C4	55.9 (2)	C1—N3—C16—C15	-105.4 (2)
C6—O3—C3—C2	-57.41 (16)	C9—N2—C15—N1	0.1 (2)
C1—C2—C3—O3	-77.96 (19)	C9—N2—C15—C16	178.43 (18)
C7—C2—C3—O3	33.32 (17)	C14—N1—C15—N2	-0.4 (2)
C1—C2—C3—C4	174.13 (18)	C14—N1—C15—C16	-178.74 (17)
C7—C2—C3—C4	-74.6 (2)	N3—C16—C15—N2	55.2 (3)
O3—C3—C4—C5	-35.0 (2)	N3—C16—C15—N1	-126.70 (19)
C2—C3—C4—C5	72.7 (2)	C15—N2—C9—C10	-179.28 (19)
C3—C4—C5—C6	1.0 (2)	C15—N2—C9—C14	0.14 (19)
C3—O3—C6—C5	-55.61 (16)	N2—C9—C10—C11	-179.60 (19)
C3—O3—C6—C7	58.61 (15)	C14—C9—C10—C11	1.0 (3)
C4—C5—C6—O3	33.5 (2)	C9—C10—C11—C12	0.6 (3)
C4—C5—C6—C7	-73.8 (2)	C10—C11—C12—C13	-1.3 (4)
C1—C2—C7—C8	2.40 (19)	C11—C12—C13—C14	0.4 (3)
C3—C2—C7—C8	-113.79 (16)	C15—N1—C14—C13	-178.2 (2)
C1—C2—C7—C6	118.36 (16)	C15—N1—C14—C9	0.44 (19)
C3—C2—C7—C6	2.16 (17)	C12—C13—C14—N1	179.8 (2)
O3—C6—C7—C8	73.7 (2)	C12—C13—C14—C9	1.3 (3)
C5—C6—C7—C8	-177.5 (2)	C10—C9—C14—N1	179.13 (16)
O3—C6—C7—C2	-37.14 (17)	N2—C9—C14—N1	-0.36 (19)
C5—C6—C7—C2	71.6 (2)	C10—C9—C14—C13	-2.0 (3)
C1—N3—C8—O1	-177.7 (2)	N2—C9—C14—C13	178.48 (17)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1A \cdots N2 ⁱ	0.86	1.97	2.827 (2)	175

Symmetry code: (i) $x+1/4, -y+1/4, z+1/4$.