

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

8-(Naphthalen-1-yl)quinoline

Godwin Kanu,^a Roger A. Lalancette^b and Dale E. Vitale^{c*}^aDepartment of Chemistry and Biochemistry, UCLA, 607 Charles E. Young Drive East, Box 951569, Los Angeles, CA 90095, USA, ^bDepartment of Chemistry, Rutgers University-Newark, 73 Warren Street, Newark, NJ 07102-1811, USA, and^cDepartment of Chemistry, Kean University, Union, NJ 07083, USACorrespondence e-mail: dvitale@kean.edu


Received 28 July 2011; accepted 19 August 2011

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.044; wR factor = 0.131; data-to-parameter ratio = 11.4.

In the title molecule, $\text{C}_{19}\text{H}_{13}\text{N}$, the angle between the mean planes of the naphthalene and quinoline ring systems is $68.59(2)^\circ$. The compound is of interest with respect to its potential for spontaneous resolution. In the crystal structure, the R and S isomers are arranged in alternating homochiral layers. The molecules of a given layer are oriented with their major axes (*i.e.* the axis perpendicular to the interannular bond) in the same direction and their naphthalene and quinoline ring systems are arranged parallel. Like the configurations, this orientation alternates in adjacent layers.

Related literature

For spontaneous-resolution experiments, see: Asakura & Plasson (2006); Kondipudi *et al.* (1999); Kranz *et al.* (1993); Sainz-Diaz *et al.* (2005); Wilson & Pincock (1974). For related structures, see: Kerr & Robertson (1969); Kuroda & Manson (1981). For details of the synthesis, see: Huff *et al.* (1998).



Experimental

Crystal data

 $\text{C}_{19}\text{H}_{13}\text{N}$ $M_r = 255.30$ Triclinic, $P\bar{1}$ $a = 6.1778(1)$ Å $b = 10.0392(2)$ Å $c = 10.8828(2)$ Å $\alpha = 104.537(1)^\circ$ $\beta = 106.435(1)^\circ$ $\gamma = 90.002(1)^\circ$ $V = 624.80(2)$ Å³ $Z = 2$ Cu $K\alpha$ radiation $\mu = 0.61$ mm⁻¹ $T = 100$ K $0.37 \times 0.20 \times 0.10$ mm

Data collection

Bruker SMART CCD APEXII diffractometer

Absorption correction: numerical (*SADABS*; Sheldrick, 2008) $T_{\min} = 0.806$, $T_{\max} = 0.942$

5783 measured reflections

2058 independent reflections

1678 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.011$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.044$ $wR(F^2) = 0.131$ $S = 1.06$

2058 reflections

181 parameters

H-atom parameters constrained

 $\Delta\rho_{\max} = 0.38$ e Å⁻³ $\Delta\rho_{\min} = -0.34$ e Å⁻³

Data collection: *APEX2* (Bruker, 2006); cell refinement: *APEX2*; data reduction: *SAINT* (Bruker, 2005); program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *Mercury* (Macrae *et al.*, 2008); software used to prepare material for publication: *SHELXTL*.

This work was partly supported by the SpF program of Kean University. The authors acknowledge support by NSF-CRIF Grant No. 0443538 for the X-ray diffractometer.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH5298).

References

- Asakura, K. & Plasson, R. (2006). *Chaos*, **16**, 1–7.
 Bruker (2005). *SAINT* Version 7.23a. Bruker AXS Inc., Madison, Wisconsin, USA.
 Bruker (2006). *APEX 2* Version 2.0-2. Bruker AXS Inc., Madison, Wisconsin, USA.
 Huff, B. E., Koenig, T. M., Mitchell, D. & Staszak, M. A. (1998). *Org. Synth.* **75**, 53–56.
 Kerr, K. A. & Robertson, J. M. (1969). *J. Chem. Soc. B*, pp. 1146–1149.
 Kondipudi, D. K., Laudadio, J. & Asakura, K. (1999). *J. Am. Chem. Soc.* **121**, 1448–1451.
 Kranz, M., Clark, T. & von Rague Schleyer, P. J. (1993). *J. Org. Chem.* **58**, 3317–3325.
 Kuroda, R. & Manson, S. F. (1981). *J. Chem. Soc. Perkin Trans. 2*, pp. 167–170.
 Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
 Sainz-Diaz, C. II, Martin-Islan, A. P. & Cartwright, J. H. E. (2005). *J. Phys. Chem.* **109**, 18758–18764.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Wilson, K. R. & Pincock, R. E. (1974). *J. Am. Chem. Soc.* **97**, 1474–1478.

supporting information

Acta Cryst. (2011). E67, o2522 [doi:10.1107/S1600536811034052]

8-(Naphthalen-1-yl)quinoline

Godwin Kanu, Roger A. Lalancette and Dale E. Vitale

S1. Comment

The structural similarities between the title compound (81QNNP) and 1,1'-binaphthalenyl (11BNP) suggest that, like the hydrocarbon, 81QNNP might exhibit spontaneous resolution. The characteristics that afford spontaneous resolution of the racemic compound of 11BNP are its dimorphism and moderate barrier to rotation about the C1—C1' interannular bond (Kranz *et al.*, 1993). The barrier is large enough to prevent racemization of resolved forms below about 351 K, but small enough to afford rapid interconversion of the enantiomers in the melt. The dimorphs consist of an optically inactive racemic compound (Kerr & Robertson, 1969) that is the more stable form at temperatures below 351 K and a conglomerate of single crystals of the *R* and *S* isomers (Kuroda & Manson, 1981) that is the more stable above this temperature. Accordingly, heating of the racemic compound above the melting point of the conglomerate (431 K) followed by supercooling to 423 K can produce an optically active solid (Kondipudi *et al.*, 1999).


The molecular structure of the title compound is shown in Fig. 1. The title compound closely resembles 1,1'-binaphthalenyl in molecular structure, crystal structure, and thermal behavior (Wilson & Pincock, 1974). On the molecular level, the two compounds differ only in the substitution of a nitrogen in 81QNNP for the C8 carbon in 11BNP. Likewise, the room temperature solid of 81QNNP is an optically inactive racemic compound with the components of an enantiotopic pair of molecules in the unit cell. Moreover, while spontaneous development of optical activity has not been demonstrated in the title compound, preliminary DSC results suggest that it is polymorphic. This means that, in addition to the racemic compound described herein, 81QNNP may also exist as a conglomerate. If so, it has the potential for spontaneous resolution *via* a mechanism similar to that of 11BNP (Asakura & Plasson, 2006; Sainz-Diaz *et al.*, 2005). In the crystal, the *R* and *S* isomers are arranged in alternating homochiral layers. The molecules of a given layer are oriented with their major axes in the same direction and their naphthalene and quinoline ring systems are arranged parallel. Like the configurations this orientation alternates in adjacent layers (Fig. 2).

S2. Experimental

The synthesis was carried out according to a literature procedure (Huff *et al.*, 1998). 8-(Naphthalen-1-yl)quinoline was synthesized in 58% yield from 8-bromoquinoline (Frontier Chemical) and 1-naphthalenylboronic acid (Sigma–Aldrich) using a modification of the Suzuki coupling reaction (Huff *et al.*, 1998) and crystallized from 1-propanol; m.p. 436.0–437.5 K, ¹³C NMR (CDCl₃): DEPT-CH δ 121.3, 125.6, 125.9, 126.0, 126.4, 127.0, 128.1, 128.2, 128.3, 128.6, 131.9, 136.4, 150.9 p.p.m. IR (KBr ν cm⁻¹): 3041 (CH_{ar}) 1592 and 1491 (C=C and C=N), 1379, 1310, 1204, 1064, 1015, 944, 828, 797, 782, 772, 677, 617, 517.

S3. Refinement

All H atoms were found in electron density difference maps. These were placed in geometrically idealized positions and constrained to ride on their parent C atoms with C—H distances of 0.95 Å, and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

**Figure 1**

The molecular structure of the *S* enantiomer of (I) with atom numbering scheme. Displacement ellipsoids are drawn at 50% probability level.




Figure 2

Alternating layers of *R* and *S* isomers viewed along *c* axis; *R*, *S*, *R*, *S* from top of figure.

8-(Naphthalen-1-yl)quinoline

Crystal data

$C_{19}H_{13}N$

$M_r = 255.30$

Triclinic, $P\bar{1}$

Hall symbol: $-P\ 1$

$a = 6.1778$ (1) Å

$b = 10.0392$ (2) Å

$c = 10.8828$ (2) Å

$\alpha = 104.537$ (1)°

$\beta = 106.435$ (1)°

$\gamma = 90.002$ (1)°

$V = 624.80$ (2) Å³

$Z = 2$

$F(000) = 268$

$D_x = 1.357$ Mg m⁻³

Melting point = 436.0–437.5 K

Cu $K\alpha$ radiation, $\lambda = 1.54178$ Å

Cell parameters from 3661 reflections

$\theta = 4.4$ – 66.7 °

$\mu = 0.61 \text{ mm}^{-1}$
 $T = 100 \text{ K}$

Plate, colourless
 $0.37 \times 0.20 \times 0.10 \text{ mm}$

Data collection

Bruker SMART CCD APEXII
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 φ and ω scans
 Absorption correction: numerical
 (SADABS; Sheldrick, 2008)
 $T_{\min} = 0.806$, $T_{\max} = 0.942$

5783 measured reflections
 2058 independent reflections
 1678 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.011$
 $\theta_{\max} = 67.5^\circ$, $\theta_{\min} = 4.4^\circ$
 $h = -7 \rightarrow 7$
 $k = -11 \rightarrow 11$
 $l = -12 \rightarrow 12$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.044$
 $wR(F^2) = 0.131$
 $S = 1.06$
 2058 reflections
 181 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0654P)^2 + 0.348P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.38 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.34 \text{ e } \text{\AA}^{-3}$

Special details

Experimental. crystal mounted on a Cryoloop using Paratone-N

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.9761 (3)	0.89896 (16)	1.10848 (15)	0.0304 (4)
C2	1.1692 (3)	0.96964 (18)	1.19213 (17)	0.0269 (4)
H2	1.2610	1.0176	1.1579	0.032*
C3	1.2400 (3)	0.97528 (18)	1.32817 (17)	0.0268 (4)
H3	1.3785	1.0249	1.3839	0.032*
C4	1.1084 (3)	0.90897 (17)	1.37947 (16)	0.0233 (4)
H4	1.1549	0.9126	1.4713	0.028*
C4A	0.9019 (3)	0.83446 (16)	1.29589 (15)	0.0207 (4)
C5	0.7566 (3)	0.76657 (17)	1.34510 (16)	0.0230 (4)
H5	0.7968	0.7702	1.4369	0.028*
C6	0.5598 (3)	0.69612 (17)	1.26156 (16)	0.0243 (4)
H6	0.4636	0.6510	1.2954	0.029*
C7	0.4982 (3)	0.68992 (17)	1.12520 (16)	0.0236 (4)

H7	0.3602	0.6406	1.0684	0.028*
C8	0.6333 (3)	0.75369 (17)	1.07236 (16)	0.0211 (4)
C8A	0.8399 (3)	0.82928 (16)	1.15855 (15)	0.0197 (4)
C1'	0.5605 (3)	0.74639 (17)	0.92755 (16)	0.0210 (4)
C2'	0.3732 (3)	0.81036 (17)	0.87514 (16)	0.0234 (4)
H2'	0.2924	0.8601	0.9321	0.028*
C3'	0.2981 (3)	0.80380 (17)	0.73857 (16)	0.0244 (4)
H3'	0.1677	0.8487	0.7048	0.029*
C4'	0.4114 (3)	0.73338 (17)	0.65465 (16)	0.0229 (4)
H4'	0.3598	0.7297	0.5628	0.027*
C4A'	0.6059 (3)	0.66569 (16)	0.70394 (15)	0.0207 (4)
C5'	0.7289 (3)	0.59114 (17)	0.62054 (16)	0.0234 (4)
H5'	0.6837	0.5877	0.5287	0.028*
C6'	0.9118 (3)	0.52436 (18)	0.67111 (17)	0.0263 (4)
H6'	0.9937	0.4743	0.6150	0.032*
C7'	0.9774 (3)	0.53047 (18)	0.80697 (17)	0.0274 (4)
H7'	1.1039	0.4830	0.8413	0.033*
C8'	0.8677 (2)	0.60111 (16)	0.89035 (15)	0.0173 (4)
H8'	0.9166	0.6033	0.9819	0.021*
C8A'	0.6816 (3)	0.67103 (16)	0.84112 (15)	0.0204 (4)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0366 (9)	0.0240 (8)	0.0316 (8)	0.0013 (6)	0.0141 (7)	0.0046 (6)
C2	0.0307 (9)	0.0211 (9)	0.0313 (9)	-0.0011 (7)	0.0153 (8)	0.0041 (7)
C3	0.0238 (9)	0.0213 (9)	0.0309 (10)	0.0001 (7)	0.0059 (7)	0.0016 (7)
C4	0.0276 (9)	0.0201 (9)	0.0197 (8)	0.0061 (7)	0.0044 (7)	0.0039 (7)
C4A	0.0249 (8)	0.0154 (9)	0.0210 (8)	0.0051 (6)	0.0070 (7)	0.0034 (6)
C5	0.0314 (9)	0.0201 (9)	0.0191 (8)	0.0057 (7)	0.0093 (7)	0.0058 (6)
C6	0.0285 (9)	0.0217 (9)	0.0261 (9)	0.0020 (7)	0.0128 (7)	0.0071 (7)
C7	0.0226 (8)	0.0221 (9)	0.0250 (9)	0.0006 (6)	0.0064 (7)	0.0049 (7)
C8	0.0240 (8)	0.0173 (9)	0.0220 (9)	0.0042 (6)	0.0074 (7)	0.0043 (6)
C8A	0.0227 (8)	0.0163 (9)	0.0211 (8)	0.0039 (6)	0.0085 (7)	0.0043 (6)
C1'	0.0215 (8)	0.0195 (9)	0.0214 (8)	-0.0026 (6)	0.0061 (7)	0.0046 (6)
C2'	0.0236 (8)	0.0229 (9)	0.0235 (9)	0.0015 (7)	0.0082 (7)	0.0043 (7)
C3'	0.0224 (8)	0.0230 (10)	0.0258 (9)	0.0017 (7)	0.0024 (7)	0.0081 (7)
C4'	0.0272 (9)	0.0209 (9)	0.0189 (8)	-0.0029 (7)	0.0035 (7)	0.0060 (6)
C4A'	0.0237 (8)	0.0162 (9)	0.0211 (8)	-0.0036 (6)	0.0046 (7)	0.0052 (6)
C5'	0.0295 (9)	0.0213 (9)	0.0197 (8)	-0.0041 (7)	0.0079 (7)	0.0051 (7)
C6'	0.0274 (9)	0.0225 (10)	0.0288 (9)	0.0004 (7)	0.0116 (7)	0.0028 (7)
C7'	0.0247 (9)	0.0210 (10)	0.0323 (10)	0.0024 (7)	0.0025 (7)	0.0061 (7)
C8'	0.0193 (8)	0.0148 (8)	0.0147 (7)	0.0006 (6)	0.0011 (6)	0.0026 (6)
C8A'	0.0222 (8)	0.0159 (9)	0.0217 (8)	-0.0033 (6)	0.0048 (7)	0.0044 (6)

Geometric parameters (Å, °)

N1—C2	1.348 (2)	C1'—C2'	1.374 (2)
N1—C8A	1.395 (2)	C1'—C8A'	1.434 (2)
C2—C3	1.406 (2)	C2'—C3'	1.410 (2)
C2—H2	0.9500	C2'—H2'	0.9500
C3—C4	1.364 (2)	C3'—C4'	1.364 (2)
C3—H3	0.9500	C3'—H3'	0.9500
C4—C4A	1.420 (2)	C4'—C4A'	1.418 (2)
C4—H4	0.9500	C4'—H4'	0.9500
C4A—C5	1.418 (2)	C4A'—C5'	1.418 (2)
C4A—C8A	1.421 (2)	C4A'—C8A'	1.419 (2)
C5—C6	1.363 (2)	C5'—C6'	1.364 (2)
C5—H5	0.9500	C5'—H5'	0.9500
C6—C7	1.409 (2)	C6'—C7'	1.403 (2)
C6—H6	0.9500	C6'—H6'	0.9500
C7—C8	1.376 (2)	C7'—C8'	1.346 (2)
C7—H7	0.9500	C7'—H7'	0.9500
C8—C8A	1.432 (2)	C8'—C8A'	1.393 (2)
C8—C1'	1.494 (2)	C8'—H8'	0.9500
C2—N1—C8A	118.89 (15)	C2'—C1'—C8	120.02 (15)
N1—C2—C3	122.60 (15)	C8A'—C1'—C8	120.85 (14)
N1—C2—H2	118.7	C1'—C2'—C3'	121.44 (15)
C3—C2—H2	118.7	C1'—C2'—H2'	119.3
C4—C3—C2	119.48 (15)	C3'—C2'—H2'	119.3
C4—C3—H3	120.3	C4'—C3'—C2'	120.43 (15)
C2—C3—H3	120.3	C4'—C3'—H3'	119.8
C3—C4—C4A	120.15 (15)	C2'—C3'—H3'	119.8
C3—C4—H4	119.9	C3'—C4'—C4A'	120.27 (15)
C4A—C4—H4	119.9	C3'—C4'—H4'	119.9
C5—C4A—C4	122.25 (15)	C4A'—C4'—H4'	119.9
C5—C4A—C8A	119.47 (15)	C5'—C4A'—C4'	122.28 (15)
C4—C4A—C8A	118.27 (15)	C5'—C4A'—C8A'	118.09 (15)
C6—C5—C4A	120.44 (15)	C4'—C4A'—C8A'	119.63 (15)
C6—C5—H5	119.8	C6'—C5'—C4A'	120.54 (15)
C4A—C5—H5	119.8	C6'—C5'—H5'	119.7
C5—C6—C7	120.34 (15)	C4A'—C5'—H5'	119.7
C5—C6—H6	119.8	C5'—C6'—C7'	119.23 (15)
C7—C6—H6	119.8	C5'—C6'—H6'	120.4
C8—C7—C6	121.54 (15)	C7'—C6'—H6'	120.4
C8—C7—H7	119.2	C8'—C7'—C6'	122.44 (15)
C6—C7—H7	119.2	C8'—C7'—H7'	118.8
C7—C8—C8A	119.05 (15)	C6'—C7'—H7'	118.8
C7—C8—C1'	119.94 (15)	C7'—C8'—C8A'	119.33 (15)
C8A—C8—C1'	120.99 (14)	C7'—C8'—H8'	120.3
N1—C8A—C4A	120.58 (15)	C8A'—C8'—H8'	120.3
N1—C8A—C8	120.26 (14)	C8'—C8A'—C4A'	120.35 (15)

C4A—C8A—C8	119.15 (15)	C8'—C8A'—C1'	120.53 (14)
C2'—C1'—C8A'	119.12 (15)	C4A'—C8A'—C1'	119.10 (15)
C8A—N1—C2—C3	0.4 (3)	C7—C8—C1'—C8A'	-112.47 (18)
N1—C2—C3—C4	-1.1 (3)	C8A—C8—C1'—C8A'	69.1 (2)
C2—C3—C4—C4A	0.2 (2)	C8A'—C1'—C2'—C3'	-0.2 (2)
C3—C4—C4A—C5	-178.48 (15)	C8—C1'—C2'—C3'	-179.08 (15)
C3—C4—C4A—C8A	1.1 (2)	C1'—C2'—C3'—C4'	-0.2 (2)
C4—C4A—C5—C6	179.99 (15)	C2'—C3'—C4'—C4A'	0.1 (2)
C8A—C4A—C5—C6	0.4 (2)	C3'—C4'—C4A'—C5'	179.92 (15)
C4A—C5—C6—C7	0.0 (2)	C3'—C4'—C4A'—C8A'	0.3 (2)
C5—C6—C7—C8	0.2 (3)	C4'—C4A'—C5'—C6'	-178.28 (15)
C6—C7—C8—C8A	-0.7 (2)	C8A'—C4A'—C5'—C6'	1.4 (2)
C6—C7—C8—C1'	-179.14 (15)	C4A'—C5'—C6'—C7'	-0.1 (2)
C2—N1—C8A—C4A	1.0 (2)	C5'—C6'—C7'—C8'	-0.6 (3)
C2—N1—C8A—C8	179.75 (15)	C6'—C7'—C8'—C8A'	0.1 (2)
C5—C4A—C8A—N1	177.87 (14)	C7'—C8'—C8A'—C4A'	1.2 (2)
C4—C4A—C8A—N1	-1.7 (2)	C7'—C8'—C8A'—C1'	179.58 (14)
C5—C4A—C8A—C8	-0.9 (2)	C5'—C4A'—C8A'—C8'	-1.9 (2)
C4—C4A—C8A—C8	179.47 (14)	C4'—C4A'—C8A'—C8'	177.77 (14)
C7—C8—C8A—N1	-177.73 (14)	C5'—C4A'—C8A'—C1'	179.69 (13)
C1'—C8—C8A—N1	0.7 (2)	C4'—C4A'—C8A'—C1'	-0.7 (2)
C7—C8—C8A—C4A	1.1 (2)	C2'—C1'—C8A'—C8'	-177.81 (14)
C1'—C8—C8A—C4A	179.50 (14)	C8—C1'—C8A'—C8'	1.1 (2)
C7—C8—C1'—C2'	66.4 (2)	C2'—C1'—C8A'—C4A'	0.6 (2)
C8A—C8—C1'—C2'	-112.04 (18)	C8—C1'—C8A'—C4A'	179.48 (14)
