

1-[(4-[(2-Oxo-1,2-dihydronaphthalen-1-ylidene)methyl]amino)anilino)methylidene]naphthalen-2(1H)-one dihydrate

Anita Blagus^{a*} and Branko Kaitner^b

^aDepartment of Chemistry, J.J. Strossmayer University, Osijek, Franje Kuhača 20, HR-31000 Osijek, Croatia, and ^bLaboratory of General and Inorganic Chemistry, Department of Chemistry, Faculty of Science, University of Zagreb, Horvatovac 102a, HR-10002 Zagreb, Croatia

Correspondence e-mail: ablagus@kemija.unios.hr


Received 29 September 2011; accepted 11 October 2011

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.061; wR factor = 0.190; data-to-parameter ratio = 15.1.

The title compound, $\text{C}_{28}\text{H}_{20}\text{N}_2\text{O}_2 \cdot 2\text{H}_2\text{O}$, comprises a Schiff base molecule with an imposed inversion centre in the middle of *p*-phenylenediamine unit and water molecules of crystallization. In the structure, the Schiff base molecule is present as the keto–amino tautomer with a strong intramolecular N–H···O hydrogen bond. The Schiff base molecules and water molecules of crystallization create infinite [010] columns through O–H···O hydrogen bonds. Intermolecular attractions within columns are through additional π – π interactions [centroid–centroid distance = 3.352 (1) Å] between parallel Schiff base molecules. The columns are joined into infinite (011) layers through weak C–H···O hydrogen bonds. The layers pack in an assembly by van der Waals attractions, only being effective between bordering non-polar naphthalene ring systems.

Related literature

For general background to Schiff bases, see: Blagus *et al.* (2010). The stereochemistry of intrinsic Schiff bases differs significantly, see: Inabe *et al.* (1994). For the quinoid effect in 2-oxy-naphthaldimine Schiff base derivatives, see: Gavrančić *et al.* (1996); Friščić *et al.* (1998). For the herringbone packing motif in fused aromatic systems, see: Desiraju & Gavezzotti (1989).



Experimental

Crystal data

$\text{C}_{28}\text{H}_{20}\text{N}_2\text{O}_2 \cdot 2\text{H}_2\text{O}$
 $M_r = 452.49$
 Monoclinic, $P2_1/c$
 $a = 17.4222$ (11) Å
 $b = 4.4686$ (5) Å
 $c = 15.9374$ (10) Å
 $\beta = 116.30$ (1)°

$V = 1112.3$ (2) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 0.09$ mm⁻¹
 $T = 298$ K
 $0.5 \times 0.2 \times 0.1$ mm

Data collection

Oxford Diffraction Xcalibur CCD diffractometer
 14006 measured reflections

2423 independent reflections
 1351 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.054$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.061$
 $wR(F^2) = 0.190$
 $S = 1.05$
 2423 reflections
 160 parameters
 3 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.20$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.27$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1}\cdots\text{O1}$	0.86	1.86	2.560 (3)	138
$\text{O1W}-\text{H1B}\cdots\text{O1}$	0.83	2.27	3.090 (4)	169
$\text{O1W}-\text{H1A}\cdots\text{O1}^{\text{i}}$	0.84	2.01	2.826 (4)	165
$\text{C13}-\text{H13}\cdots\text{O1W}^{\text{ii}}$	0.93	2.33	3.247 (5)	170

Symmetry codes: (i) $x, y + 1, z$; (ii) $x, -y + \frac{3}{2}, z - \frac{1}{2}$.

Data collection: *CrysAlis CCD* (Oxford Diffraction, 2003); cell refinement: *CrysAlis CCD*; data reduction: *CrysAlis RED* (Oxford Diffraction, 2003); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999), *PARST97* (Nardelli, 1995) and *Mercury* (Macrae *et al.*, 2006).

Financial support by the Ministry of Science, Education and Sport of the Republic of Croatia is gratefully acknowledged (grant No. 119–1193079–3069).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: KP2357).

References

- Blagus, A., Cinčić, D., Friščić, T., Kaitner, B. & Stilinović, V. (2010). *Maced. J. Chem. Chem. Eng.* **29**, 117–138.
- Desiraju, G. R. & Gavezzotti, A. (1989). *Acta Cryst.* **B45**, 473–482.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Friščić, T., Kaitner, B. & Meštrović, E. (1998). *Croat. Chem. Acta*, **71**, 87–98.
- Gavrančić, M., Kaitner, B. & Meštrović, E. (1996). *J. Chem. Crystallogr.* **26**, 23–28.
- Inabe, T., Luneau, I., Mitani, T., Maruyama, Y. & Takeda, S. (1994). *Bull. Chem. Soc. Jpn*, **67**, 612–621.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- Nardelli, M. (1995). *J. Appl. Cryst.* **28**, 659.
- Oxford Diffraction (2003). *CrysAlis CCD* and *CrysAlis RED*. Oxford Diffraction Ltd, Wrocław, Poland.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supporting information

Acta Cryst. (2011). E67, o2958–o2959 [doi:10.1107/S1600536811041870]

1-[(4-[(2-Oxo-1,2-dihydronaphthalen-1-ylidene)methyl]amino)anilino)methylidene]naphthalen-2(1*H*)-one dihydrate

Anita Blagus and Branko Kaitner

S1. Comment

Desiraju and Gavezzotti presented classification of packing arrangements for polynuclear aromatic hydrocarbons depending on the number and positions of C and H atoms in molecules (Desiraju & Gavezzotti, 1989). As a result of significant planarity, Schiff bases derived from *p*-phenylenediamine have significant aromatic-aromatic C \cdots C interactions compared to the number of intermolecular C–H hydrogen bonds and usually show a herringbone motif of intermolecular assembly (Blagus *et al.*, 2010).

We report here the crystal structure of the title compound (I) as a crystal hydrate. The stereochemistry of intrinsic 1,4-bis(2-hydroxy-1-naphthylmethylideneamino)benzene was earlier determined and reported (Inabe *et al.*, 1994).

Distinctively planar Schiff base molecule possesses crystallographic inversion centre in the middle of *p*-phenylenediamine moiety with *anti* arrangements of chelate rings (Fig. 1). Interplanar angle between naphthalene moiety and the central aromatic ring is 1.7 (2)°. Schiff base molecule does not deviate significantly from planarity in contrast to the structure by Inabe *et al.* with corresponding interplanar angle being 22.3°. In both structures molecules possess internal (molecular) symmetry with inversion centre in the middle of *p*-phenylenediamine moiety.

Bond distances C2–O1 [1.277 (4) Å] and C11–N1 [1.322 (4) Å] indicate keto-amino tautomeric form of (I). This is confirmed by a formation of strong intramolecular hydrogen bond N–H \cdots O [N \cdots O = 2.560 (3) Å]. Short C3-to-C4 bond distance [1.345 (5) Å] with O1 oxygen atom at C2 position of naphthalene core indicates the presence of quinoid effect (Gavranić *et al.*, 1996; Friščić *et al.*, 1998). Water molecules play crucial role in crystal packing: a) as bridging media pilling up Schiff base molecules at the separation characteristic for graphite in the form of infinite [010] columns and b) as bridging molecule connecting neighboring [010] columns into infinite (011) layers. Hydrogen bonds effective for columns formation are: a) O1W–H1A \cdots O1ⁱ 2.826 (4) Å [(i): *x*, *y* + 1, *z*] and O1W–H1B \cdots O1 3.090 (4) Å while for layers formation is effective b) C13–H13 \cdots OW1ⁱⁱ 3.247 (5) Å [(ii): *x*, –*y* + 3/2, *z*] (Fig. 2 and Table 1).

Along with intermolecular contacts *via* hydrogen bonds the linking between molecules amplifies through π – π interactions with offset (Fig. 2). Schiff base molecules arrange parallel to each other with their middle *p*-phenylenediamine moiety being separated at distances characteristic for layer separation in graphite. The shortest separations corresponding to the sum of van der Waals radii are: C12 \cdots C12ⁱⁱⁱ 3.362 (3) Å [(iii): –*x*, –*y* + 1, –*z*] and C11 \cdots C13^{iv} 3.374 (3) Å [(iv): *x*, *y* – 1, *z*]. π \cdots π interactions are also characterised by perpendicular Cgⁱⁱⁱ \cdots Cg^v distance 3.352 (1) Å [(v): –*x*, 3 – *y*, –*z*] and slippage of 2.955 Å. There is a space between each pair of neighboring columns large enough to accommodate water molecules (Fig. 3). Connection between neighboring parallel layers is accomplished through bordering non-polar naphthalene core by the standard van der Waals attractions. The rather planar Schiff base molecules of (I) reveal characteristic herringbone motif of packing arrangement (Fig. 4).

S2. Experimental

The crystals of (I), (1,4-bis(2-hydroxy-1-naphthylmethylideneamino)benzene) as crystal hydrate were obtained during an unsuccessful attempt to synthesise the nickel complex of corresponding Schiff base. Schiff base itself was prepared separately in standard way by condensation of 2-hydroxy-naphthaldehyde and *p*-phenylenediamine in ethanol solution in molar ratio 2:1 and used as a ligand in metal complex synthesis. The 1:1 mixture of 0.1 mmol DMSO solutions of Schiff base and 0.2 mmol nickel salt, NiCl₂ · 6H₂O was stirred under reflux for two h at 373 K. Preparation of nickel complex failed. The crystals of title compound crystallised from mother liquor after cooling to RT and mechanically separated from nickel salt.

S3. Refinement

Hydrogen atoms were refined in two different ways.

For hydrogen atoms bonded to C and N atoms benzene type riding mode was used with C-to-H and N-to-H bond distances taken as 0.93 and 0.86 Å, respectively.

Due to somewhat higher values of anisotropic thermal parameters of O1W oxygen atom, implying to certain disorder of water molecule, bond distances O1W to H1A and H1B, respectively, as well as bond distance H1A–H1B were restrained to the values accepted for water molecule. Bond distances O1W to H1A and H1B, respectively were fixed to 0.82 (1) Å and H1A to H1B to 1.30 (1) Å and the position of hydrogen atoms were re-calculated in consecutive refinement cycles. Isotropic thermal parameters for hydrogen atoms were estimated as 1.2 times of equivalent isotropic thermal parameter of corresponding C, N and O atoms.





Figure 1

A general overview of (I) showing numbering scheme with anisotropic thermal ellipsoids pictured at 30% probability level. Thin lines display intra N–H···O, and the two intermolecular hydrogen bonds O–H···O and C–H···O. Hydrogen atoms are drawn as spheres of arbitrary radius.

**Figure 2**

The display of three intermolecular contacts of water molecule with nearby Schiff base molecules. Two Schiff base molecules connected by π - π interactions are additionally bridged with a water molecule through two strong hydrogen bonds O1W-H1A...O1ⁱ [(i): $x, y + 1, z$] and O1W-H1B...O1, respectively. The third intermolecular contact of water molecule is *via* C13-H13...OW1ⁱⁱ 3.247 (5) Å [(ii): $x, -y + 3/2, z$] interaction.





Figure 3

Water molecules of crystallisation occupy the cylindrical voids formed by assembling Schiff base molecules into the [010] columns.

**Figure 4**

The display of herringbone packing arrangement of Schiff base molecules viewed down *c*-axes. Water molecules and all hydrogen atoms were omitted for clarity.

1-[(4-[(2-Oxo-1,2-dihydronaphthalen-1-ylidene)methyl]amino)anilino)methylidene]naphthalen-2(1H)-one dihydrate

Crystal data

$C_{28}H_{20}N_2O_2 \cdot 2H_2O$

$M_r = 452.49$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 17.4222$ (11) Å

$b = 4.4686$ (5) Å

$c = 15.9374$ (10) Å

$\beta = 116.30$ (1)°

$V = 1112.3$ (2) Å³

$Z = 2$

$F(000) = 476$

$D_x = 1.351$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 2423 reflections

$\theta = 4-27^\circ$

$\mu = 0.09$ mm⁻¹

$T = 298$ K

Prism, green

$0.5 \times 0.2 \times 0.1$ mm

Data collection

Oxford Diffraction Xcalibur CCD
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

ω scans

14006 measured reflections

2423 independent reflections

1351 reflections with $I > 2\sigma(I)$

$R_{int} = 0.054$

$\theta_{max} = 27.0^\circ$, $\theta_{min} = 3.9^\circ$

$h = -22 \rightarrow 21$

$k = -5 \rightarrow 5$

$l = -20 \rightarrow 20$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.061$
 $wR(F^2) = 0.190$
 $S = 1.05$
 2423 reflections
 160 parameters
 3 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H atoms treated by a mixture of independent
 and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0914P)^2 + 0.2128P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.009$
 $\Delta\rho_{\max} = 0.20 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.27 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.19806 (12)	0.2820 (4)	0.25053 (12)	0.0537 (6)
N1	0.12888 (13)	0.5855 (4)	0.09888 (14)	0.0420 (5)
H1	0.1321	0.5349	0.1524	0.050*
C1	0.24546 (15)	0.2518 (5)	0.13184 (17)	0.0383 (6)
C2	0.25000 (16)	0.1743 (5)	0.22164 (17)	0.0416 (6)
C3	0.31492 (17)	-0.0329 (6)	0.27967 (18)	0.0486 (7)
H3	0.3186	-0.0853	0.3378	0.058*
C4	0.37040 (18)	-0.1525 (6)	0.25139 (19)	0.0495 (7)
H4	0.4116	-0.2855	0.2910	0.059*
C5	0.36876 (15)	-0.0839 (5)	0.16289 (18)	0.0427 (6)
C6	0.42800 (18)	-0.2111 (6)	0.1358 (2)	0.0527 (7)
H6	0.4696	-0.3410	0.1764	0.063*
C7	0.42563 (19)	-0.1479 (7)	0.0517 (2)	0.0583 (8)
H7	0.4654	-0.2326	0.0345	0.070*
C8	0.36356 (19)	0.0436 (6)	-0.0084 (2)	0.0568 (8)
H8	0.3616	0.0850	-0.0665	0.068*
C9	0.30524 (18)	0.1733 (6)	0.01493 (19)	0.0498 (7)
H9	0.2643	0.3014	-0.0274	0.060*
C10	0.30542 (15)	0.1174 (5)	0.10235 (17)	0.0387 (6)
C11	0.18466 (15)	0.4573 (5)	0.07490 (17)	0.0399 (6)
H11	0.1832	0.5066	0.0175	0.048*
C12	0.06465 (15)	0.7953 (5)	0.04716 (18)	0.0396 (6)
C13	0.05037 (17)	0.8940 (6)	-0.04079 (19)	0.0466 (7)
H13	0.0838	0.8227	-0.0685	0.056*

C14	-0.01402 (16)	1.0993 (6)	-0.08723 (18)	0.0450 (6)
H14	-0.0233	1.1673	-0.1461	0.054*
O1W	0.1500 (2)	0.8211 (6)	0.3392 (2)	0.0919 (9)
H1A	0.160 (3)	0.978 (4)	0.317 (3)	0.110*
H1B	0.170 (3)	0.687 (5)	0.318 (3)	0.110*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0561 (12)	0.0580 (12)	0.0539 (12)	0.0074 (10)	0.0306 (10)	0.0081 (9)
N1	0.0396 (12)	0.0385 (11)	0.0463 (12)	0.0040 (10)	0.0177 (10)	0.0037 (10)
C1	0.0330 (13)	0.0333 (12)	0.0435 (14)	-0.0026 (11)	0.0122 (11)	-0.0004 (11)
C2	0.0396 (14)	0.0392 (14)	0.0462 (15)	-0.0029 (12)	0.0193 (13)	0.0003 (12)
C3	0.0490 (16)	0.0481 (15)	0.0434 (15)	0.0008 (13)	0.0156 (13)	0.0080 (12)
C4	0.0414 (15)	0.0442 (15)	0.0511 (16)	0.0031 (13)	0.0097 (13)	0.0079 (13)
C5	0.0323 (13)	0.0402 (13)	0.0495 (15)	-0.0007 (11)	0.0124 (12)	-0.0009 (12)
C6	0.0394 (15)	0.0461 (16)	0.0653 (19)	0.0066 (12)	0.0166 (14)	0.0016 (14)
C7	0.0535 (18)	0.0555 (17)	0.074 (2)	0.0034 (15)	0.0361 (17)	-0.0068 (16)
C8	0.0609 (19)	0.0574 (18)	0.0573 (18)	0.0021 (15)	0.0309 (16)	-0.0027 (14)
C9	0.0511 (17)	0.0498 (16)	0.0480 (16)	0.0090 (13)	0.0214 (14)	0.0045 (12)
C10	0.0345 (13)	0.0333 (12)	0.0442 (14)	-0.0025 (11)	0.0138 (11)	-0.0025 (11)
C11	0.0373 (14)	0.0352 (13)	0.0449 (14)	-0.0005 (11)	0.0162 (12)	-0.0029 (11)
C12	0.0365 (14)	0.0328 (13)	0.0451 (14)	0.0008 (11)	0.0142 (12)	-0.0017 (11)
C13	0.0433 (15)	0.0454 (14)	0.0536 (16)	0.0058 (13)	0.0238 (13)	0.0002 (13)
C14	0.0454 (15)	0.0457 (14)	0.0427 (14)	0.0052 (13)	0.0185 (13)	0.0039 (12)
O1W	0.113 (2)	0.0952 (18)	0.0958 (19)	0.0004 (18)	0.0719 (17)	0.0129 (17)

Geometric parameters (Å, °)

O1—C2	1.277 (3)	C6—H6	0.9300
N1—C11	1.322 (3)	C7—C8	1.378 (4)
N1—C12	1.412 (3)	C7—H7	0.9300
N1—H1	0.8600	C8—C9	1.357 (4)
C1—C11	1.392 (3)	C8—H8	0.9300
C1—C2	1.440 (3)	C9—C10	1.414 (4)
C1—C10	1.452 (3)	C9—H9	0.9300
C2—C3	1.437 (3)	C11—H11	0.9300
C3—C4	1.345 (4)	C12—C13	1.383 (4)
C3—H3	0.9300	C13—C14	1.383 (4)
C4—C5	1.431 (4)	C13—H13	0.9300
C4—H4	0.9300	C14—H14	0.9300
C5—C6	1.404 (4)	O1W—H1A	0.836 (10)
C5—C10	1.419 (3)	O1W—H1B	0.831 (10)
C6—C7	1.353 (4)		
C11—N1—C12	127.8 (2)	C6—C7—C8	119.2 (3)
C11—N1—H1	116.1	C6—C7—H7	120.4
C12—N1—H1	116.1	C8—C7—H7	120.4

C11—C1—C2	119.8 (2)	C9—C8—C7	121.9 (3)
C11—C1—C10	120.7 (2)	C9—C8—H8	119.1
C2—C1—C10	119.5 (2)	C7—C8—H8	119.1
O1—C2—C3	119.5 (2)	C8—C9—C10	121.3 (3)
O1—C2—C1	122.1 (2)	C8—C9—H9	119.4
C3—C2—C1	118.4 (2)	C10—C9—H9	119.4
C4—C3—C2	121.2 (2)	C9—C10—C5	116.3 (2)
C4—C3—H3	119.4	C9—C10—C1	123.9 (2)
C2—C3—H3	119.4	C5—C10—C1	119.8 (2)
C3—C4—C5	122.8 (2)	N1—C11—C1	122.8 (2)
C3—C4—H4	118.6	N1—C11—H11	118.6
C5—C4—H4	118.6	C1—C11—H11	118.6
C6—C5—C10	120.3 (2)	C13—C12—N1	122.9 (2)
C6—C5—C4	121.3 (2)	C12—C13—C14	119.5 (2)
C10—C5—C4	118.4 (2)	C12—C13—H13	120.2
C7—C6—C5	121.1 (3)	C14—C13—H13	120.2
C7—C6—H6	119.4	C13—C14—H14	119.6
C5—C6—H6	119.4	H1A—O1W—H1B	103 (2)
C11—C1—C2—O1	2.0 (4)	C8—C9—C10—C1	179.3 (2)
C10—C1—C2—O1	-178.9 (2)	C6—C5—C10—C9	1.3 (4)
C11—C1—C2—C3	-178.5 (2)	C4—C5—C10—C9	-178.7 (2)
C10—C1—C2—C3	0.6 (3)	C6—C5—C10—C1	-178.9 (2)
O1—C2—C3—C4	179.6 (2)	C4—C5—C10—C1	1.1 (3)
C1—C2—C3—C4	0.0 (4)	C11—C1—C10—C9	-2.3 (4)
C2—C3—C4—C5	-0.1 (4)	C2—C1—C10—C9	178.6 (2)
C3—C4—C5—C6	179.6 (3)	C11—C1—C10—C5	177.9 (2)
C3—C4—C5—C10	-0.5 (4)	C2—C1—C10—C5	-1.2 (3)
C10—C5—C6—C7	-0.7 (4)	C12—N1—C11—C1	-179.9 (2)
C4—C5—C6—C7	179.3 (3)	C2—C1—C11—N1	-0.9 (3)
C5—C6—C7—C8	-0.3 (4)	C10—C1—C11—N1	-179.9 (2)
C6—C7—C8—C9	0.7 (4)	C11—N1—C12—C13	1.3 (4)
C7—C8—C9—C10	-0.1 (4)	N1—C12—C13—C14	179.8 (2)
C8—C9—C10—C5	-0.9 (4)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N1—H1...O1	0.86	1.86	2.560 (3)	138
O1W—H1B...O1	0.83	2.27	3.090 (4)	169
O1W—H1A...O1 ⁱ	0.84	2.01	2.826 (4)	165
C13—H13...O1W ⁱⁱ	0.93	2.33	3.247 (5)	170

Symmetry codes: (i) *x*, *y*+1, *z*; (ii) *x*, -*y*+3/2, *z*-1/2.