

## Methyl pyrido[2,3-*b*]pyrazine-3-carboxylate

Hoong-Kun Fun,<sup>a\*</sup>‡ Madhukar Hemamalini,<sup>a</sup> Anita Hazra<sup>b</sup> and Shyamaprosad Goswami<sup>b</sup>

<sup>a</sup>X-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, and <sup>b</sup>Department of Chemistry, Bengal Engineering and Science University, Shibpur, Howrah 711 103, India  
Correspondence e-mail: hkfun@usm.my


Received 11 October 2011; accepted 25 October 2011

Key indicators: single-crystal X-ray study;  $T = 100\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.001\text{ \AA}$ ;  $R$  factor = 0.041;  $wR$  factor = 0.115; data-to-parameter ratio = 19.1.

The asymmetric unit of the title compound,  $\text{C}_9\text{H}_7\text{N}_3\text{O}_2$ , is composed of two independent molecules. The crystal structure is stabilized by  $\text{C}-\text{H}\cdots\text{O}$  and  $\text{C}-\text{H}\cdots\text{N}$  hydrogen bonds, forming a three-dimensional network. The crystal structure also features pyrazine–pyrazine  $\pi\cdots\pi$  interactions [centroid–centroid distance =  $3.6994(5)\text{ \AA}$ ] and also pyridine–pyrazine  $\pi\cdots\pi$  interactions [centroid–centroid distance =  $3.6374(5)\text{ \AA}$ ].

### Related literature

For details of heterocyclic esters, see: Listvan *et al.* (2002); Li *et al.* (2007); Goswami & Hazra (2009); Goswami *et al.* (2011). For reference bond-length data, see: Allen *et al.* (1987). For the stability of the temperature controller used in the data collection, see: Cosier & Glazer (1986).



### Experimental

#### Crystal data

$\text{C}_9\text{H}_7\text{N}_3\text{O}_2$	$V = 1654.24(4)\text{ \AA}^3$
$M_r = 189.18$	$Z = 8$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 9.5135(1)\text{ \AA}$	$\mu = 0.11\text{ mm}^{-1}$
$b = 26.9042(3)\text{ \AA}$	$T = 100\text{ K}$
$c = 6.7837(1)\text{ \AA}$	$0.22 \times 0.20 \times 0.19\text{ mm}$
$\beta = 107.686(1)^\circ$	

#### Data collection

Bruker SMART APEXII CCD area-detector diffractometer	18421 measured reflections
Absorption correction: multi-scan ( <i>SADABS</i> ; Bruker, 2009)	4876 independent reflections
$T_{\min} = 0.975$ , $T_{\max} = 0.979$	4118 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.025$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.041$	255 parameters
$wR(F^2) = 0.115$	H-atom parameters constrained
$S = 1.04$	$\Delta\rho_{\max} = 0.38\text{ e \AA}^{-3}$
4876 reflections	$\Delta\rho_{\min} = -0.30\text{ e \AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C3A—H3AA $\cdots$ O2A <sup>i</sup>	0.93	2.54	3.2401 (13)	133
C4A—H4AA $\cdots$ N2B <sup>ii</sup>	0.93	2.55	3.3311 (14)	141
C9A—H9AA $\cdots$ N1B	0.96	2.62	3.4741 (14)	149
C3B—H3BA $\cdots$ O2B <sup>ii</sup>	0.93	2.53	3.2496 (14)	135
C4B—H4BA $\cdots$ N2A <sup>iv</sup>	0.93	2.51	3.3350 (14)	147
C9B—H9BA $\cdots$ N1A	0.96	2.57	3.4266 (14)	149

Symmetry codes: (i)  $x - 1, y, z$ ; (ii)  $-x + 1, y - \frac{1}{2}, -z + \frac{3}{2}$ ; (iii)  $x + 1, y, z$ ; (iv)  $-x + 2, y + \frac{1}{2}, -z + \frac{3}{2}$ .

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL* and *PLATON* (Spek, 2009).

HKF and MH thank the Malaysian Government and Universiti Sains Malaysia for the Research University Grant No. 1001/PFIZIK/811160. MH also thanks Universiti Sains Malaysia for a post-doctoral research fellowship. SG and AH thank the CSIR [No. 01 (2292)/09/EMR-II], Government of India, for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: WN2453).

### References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Bruker (2009). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cosier, J. & Glazer, A. M. (1986). *J. Appl. Cryst.* **19**, 105–107.
- Goswami, S. & Hazra, A. (2009). *Chem. Lett.* **38**, 484–485.
- Goswami, S., Hazra, A., Goh, J. H. & Fun, H.-K. (2011). *Synth. Commun.* **41**, 2505–2510.
- Li, D. Z., Li, Y., Chen, X. G., Zhu, C. G., Yang, J., Liu, H. Y. & Pan, X. D. (2007). *Chin. Chem. Lett.* **18**, 1335–1338.
- Listvan, V. N., Listvan, V. V. & Skekel, A. N. (2002). *Chem. Heterocycl. Compd.* **38**, 1480–1483.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.

‡ Thomson Reuters ResearcherID: A-3561-2009.

# supporting information

*Acta Cryst.* (2011). E67, o3120 [doi:10.1107/S1600536811044412]

## Methyl pyrido[2,3-*b*]pyrazine-3-carboxylate

**Hoong-Kun Fun, Madhukar Hemamalini, Anita Hazra and Shyamaprosad Goswami**

### S1. Comment

Heterocyclic esters are important compounds in respect of their biological and pharmaceutical characteristics (Listvan *et al.*, 2002; Li *et al.*, 2007). Recently we have developed a mild methodology for the synthesis of heterocyclic esters from their corresponding aldehydes (Goswami & Hazra, 2009; Goswami *et al.*, 2011). Here we report the crystal structure of methyl pyrido[2,3-*b*]pyrazine-3-carboxylate.

The asymmetric unit of the title compound consists of two crystallographically independent methyl pyrido[2,3-*b*]pyrazine-3-carboxylate molecules, (A & B), as shown in Fig. 1. The bond lengths of molecules A and B agree with each other and are within normal ranges (Allen *et al.*, 1987).

In the crystal structure (Fig. 2), the molecules are linked through intermolecular C—H···O and C—H···N hydrogen bonds (Table 1), forming a three-dimensional network. Furthermore, the crystal structure is stabilized by the following  $\pi$ – $\pi$  interactions: (a) between pyrazine rings (N1A,N3A/C1A,C2A/C6A,C7A, centroid Cg1) Cg1···Cg1(1-x, -y, 1-z) 3.6994 (5) Å and (b) between pyrazine (N1B,N3B/C1B,C2B/C6B,C7B, centroid Cg4) and pyridine (N2B/C2B–C6B, centroid Cg5) rings Cg4···Cg5( x, 1/2-y, 1/2+z) 3.6374 (5) Å


The two independent molecules in the asymmetric unit are strikingly similar in respect of their bond distances, bond angles and thermal parameters. For example, for bond distances the average  $\delta/\sigma = 0.88$ . The symmetry between molecules A and B is almost that of an inversion centre.

### S2. Experimental


Methyl pyrido[2,3-*b*]pyrazine-3-carboxylate was synthesized from pyrido[2,3-*b*] pyrazine-3-carbaldehyde by our recently developed techniques (Goswami & Hazra, 2009; Goswami *et al.*, 2011). Single crystals were grown by slow evaporation of a chloroform solution of the compound, Mp 155–156°C.

### S3. Refinement

All hydrogen atoms were positioned geometrically, with  $Csp^2$ —H = 0.93 Å and C(methyl)—H = 0.96 Å; they were refined using a riding model, with  $U_{iso}(\text{H}) = xU_{eq}(\text{C})$ , where x = 1.5 for methyl H and 1.2 for all other H atoms. A rotating group model was applied to the methyl groups.

**Figure 1**

The asymmetric unit of the title compound, showing 50% probability displacement ellipsoids. Dashed lines indicate hydrogen bonds.

**Figure 2**

The crystal packing of the title compound. Dashed lines indicate hydrogen bonds.

**Methyl pyrido[2,3-*b*]pyrazine-3-carboxylate***Crystal data*

$C_9H_7N_3O_2$   
 $M_r = 189.18$   
Monoclinic,  $P2_1/c$   
Hall symbol: -P 2ybc  
 $a = 9.5135 (1)$  Å  
 $b = 26.9042 (3)$  Å  
 $c = 6.7837 (1)$  Å  
 $\beta = 107.686 (1)^\circ$   
 $V = 1654.24 (4)$  Å<sup>3</sup>  
 $Z = 8$

$F(000) = 784$   
 $D_x = 1.519 \text{ Mg m}^{-3}$   
Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å  
Cell parameters from 6855 reflections  
 $\theta = 2.3\text{--}30.1^\circ$   
 $\mu = 0.11 \text{ mm}^{-1}$   
 $T = 100$  K  
Block, brown  
 $0.22 \times 0.20 \times 0.19$  mm

*Data collection*

Bruker SMART APEXII CCD area-detector  
diffractometer  
Radiation source: fine-focus sealed tube  
Graphite monochromator  
 $\varphi$  and  $\omega$  scans  
Absorption correction: multi-scan  
(SADABS; Bruker, 2009)  
 $T_{\min} = 0.975$ ,  $T_{\max} = 0.979$

18421 measured reflections  
4876 independent reflections  
4118 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.025$   
 $\theta_{\max} = 30.1^\circ$ ,  $\theta_{\min} = 2.3^\circ$   
 $h = -13 \rightarrow 12$   
 $k = -36 \rightarrow 37$   
 $l = -8 \rightarrow 9$

*Refinement*

Refinement on  $F^2$   
Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.041$   
 $wR(F^2) = 0.115$   
 $S = 1.04$   
4876 reflections  
255 parameters  
0 restraints  
Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map  
Hydrogen site location: inferred from  
neighbouring sites  
H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0645P)^2 + 0.3345P]$   
where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} < 0.001$   
 $\Delta\rho_{\max} = 0.38 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.30 \text{ e } \text{\AA}^{-3}$

*Special details*

**Experimental.** The crystal was placed in the cold stream of an Oxford Cryosystems Cobra open-flow nitrogen cryostat (Cosier & Glazer, 1986) operating at 100.0 (1) K.

**Geometry.** All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted R-factor  $wR$  and goodness of fit S are based on  $F^2$ , conventional R-factors R are based on F, with F set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\sigma(F^2)$  is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on  $F^2$  are statistically about twice as large as those based on F, and R-factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å<sup>2</sup>)*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1A	0.93918 (7)	0.07303 (3)	0.70643 (12)	0.01812 (16)
O2A	1.02425 (8)	-0.00559 (3)	0.76407 (12)	0.02050 (17)

N1A	0.53239 (9)	0.03988 (3)	0.73953 (13)	0.01616 (17)
N2A	0.58303 (9)	-0.09571 (3)	0.79786 (14)	0.02002 (18)
N3A	0.74901 (9)	-0.03476 (3)	0.77626 (13)	0.01595 (17)
C1A	0.66889 (10)	0.05049 (4)	0.74572 (15)	0.01556 (19)
H1AA	0.6958	0.0836	0.7411	0.019*
C2A	0.50066 (10)	-0.00925 (3)	0.75265 (14)	0.01429 (18)
C3A	0.35658 (10)	-0.02404 (4)	0.74564 (15)	0.01738 (19)
H3AA	0.2817	-0.0007	0.7284	0.021*
C4A	0.32995 (11)	-0.07339 (4)	0.76471 (16)	0.0193 (2)
H4AA	0.2361	-0.0843	0.7595	0.023*
C5A	0.44732 (11)	-0.10778 (4)	0.79259 (16)	0.0209 (2)
H5AA	0.4274	-0.1411	0.8082	0.025*
C6A	0.60995 (10)	-0.04658 (4)	0.77600 (15)	0.01502 (18)
C7A	0.77626 (10)	0.01276 (3)	0.75932 (14)	0.01413 (18)
C8A	0.92792 (10)	0.02481 (4)	0.74628 (15)	0.01526 (19)
C9A	1.07833 (11)	0.08837 (4)	0.67914 (17)	0.0203 (2)
H9AA	1.0750	0.1233	0.6488	0.031*
H9AB	1.0951	0.0701	0.5668	0.031*
H9AC	1.1570	0.0819	0.8038	0.031*
O1B	0.52796 (7)	0.17332 (3)	0.64501 (12)	0.01820 (16)
O2B	0.45239 (8)	0.25302 (3)	0.63114 (13)	0.02204 (17)
N1B	0.93475 (9)	0.20808 (3)	0.61052 (13)	0.01529 (17)
N2B	0.89543 (9)	0.34431 (3)	0.62179 (13)	0.01837 (18)
N3B	0.72566 (9)	0.28254 (3)	0.61794 (13)	0.01472 (16)
C1B	0.79969 (10)	0.19717 (3)	0.61024 (15)	0.01489 (18)
H1BA	0.7715	0.1640	0.6057	0.018*
C2B	0.96972 (10)	0.25733 (3)	0.61735 (14)	0.01400 (18)
C3B	1.11389 (11)	0.27225 (4)	0.62514 (15)	0.0176 (2)
H3BA	1.1861	0.2489	0.6259	0.021*
C4B	1.14411 (11)	0.32193 (4)	0.63151 (16)	0.0199 (2)
H4BA	1.2379	0.3330	0.6372	0.024*
C5B	1.03117 (11)	0.35652 (4)	0.62938 (16)	0.0196 (2)
H5BA	1.0545	0.3901	0.6336	0.024*
C6B	0.86440 (10)	0.29472 (3)	0.61752 (14)	0.01401 (18)
C7B	0.69557 (10)	0.23467 (3)	0.61664 (14)	0.01338 (18)
C8B	0.54412 (10)	0.22230 (4)	0.62971 (15)	0.01495 (18)
C9B	0.39006 (11)	0.15787 (4)	0.67601 (18)	0.0215 (2)
H9BA	0.3901	0.1224	0.6927	0.032*
H9BB	0.3093	0.1673	0.5581	0.032*
H9BC	0.3795	0.1736	0.7977	0.032*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O1A	0.0143 (3)	0.0134 (3)	0.0275 (4)	-0.0008 (2)	0.0076 (3)	0.0009 (3)
O2A	0.0153 (3)	0.0168 (3)	0.0304 (4)	0.0025 (3)	0.0085 (3)	0.0018 (3)
N1A	0.0162 (4)	0.0156 (4)	0.0176 (4)	0.0015 (3)	0.0065 (3)	0.0003 (3)
N2A	0.0175 (4)	0.0140 (4)	0.0271 (4)	-0.0009 (3)	0.0047 (3)	0.0023 (3)

N3A	0.0133 (4)	0.0139 (4)	0.0195 (4)	0.0006 (3)	0.0034 (3)	0.0012 (3)
C1A	0.0165 (4)	0.0129 (4)	0.0183 (4)	0.0012 (3)	0.0068 (3)	0.0008 (3)
C2A	0.0142 (4)	0.0154 (4)	0.0132 (4)	0.0008 (3)	0.0040 (3)	0.0001 (3)
C3A	0.0146 (4)	0.0224 (5)	0.0156 (4)	0.0010 (3)	0.0053 (3)	0.0002 (4)
C4A	0.0155 (4)	0.0243 (5)	0.0181 (4)	-0.0044 (4)	0.0050 (3)	0.0000 (4)
C5A	0.0206 (5)	0.0181 (5)	0.0231 (5)	-0.0045 (4)	0.0051 (4)	0.0014 (4)
C6A	0.0142 (4)	0.0137 (4)	0.0164 (4)	0.0006 (3)	0.0035 (3)	0.0012 (3)
C7A	0.0133 (4)	0.0133 (4)	0.0153 (4)	0.0012 (3)	0.0035 (3)	0.0005 (3)
C8A	0.0144 (4)	0.0144 (4)	0.0165 (4)	-0.0004 (3)	0.0040 (3)	-0.0009 (3)
C9A	0.0165 (4)	0.0168 (5)	0.0298 (5)	-0.0020 (3)	0.0101 (4)	0.0011 (4)
O1B	0.0143 (3)	0.0148 (3)	0.0272 (4)	-0.0014 (2)	0.0087 (3)	0.0005 (3)
O2B	0.0166 (3)	0.0183 (4)	0.0338 (4)	0.0029 (3)	0.0116 (3)	0.0028 (3)
N1B	0.0147 (4)	0.0154 (4)	0.0166 (4)	0.0012 (3)	0.0061 (3)	-0.0005 (3)
N2B	0.0211 (4)	0.0140 (4)	0.0215 (4)	-0.0020 (3)	0.0085 (3)	-0.0009 (3)
N3B	0.0146 (4)	0.0145 (4)	0.0159 (4)	0.0010 (3)	0.0060 (3)	0.0004 (3)
C1B	0.0155 (4)	0.0127 (4)	0.0172 (4)	0.0004 (3)	0.0062 (3)	0.0001 (3)
C2B	0.0134 (4)	0.0156 (4)	0.0132 (4)	0.0008 (3)	0.0043 (3)	0.0000 (3)
C3B	0.0138 (4)	0.0217 (5)	0.0177 (4)	-0.0004 (3)	0.0056 (3)	0.0000 (4)
C4B	0.0157 (4)	0.0245 (5)	0.0195 (5)	-0.0054 (4)	0.0052 (4)	-0.0006 (4)
C5B	0.0225 (5)	0.0170 (5)	0.0196 (5)	-0.0053 (4)	0.0068 (4)	-0.0007 (4)
C6B	0.0144 (4)	0.0143 (4)	0.0139 (4)	0.0002 (3)	0.0051 (3)	-0.0002 (3)
C7B	0.0122 (4)	0.0146 (4)	0.0138 (4)	0.0013 (3)	0.0047 (3)	0.0007 (3)
C8B	0.0142 (4)	0.0159 (4)	0.0150 (4)	-0.0004 (3)	0.0048 (3)	0.0007 (3)
C9B	0.0152 (4)	0.0189 (5)	0.0324 (6)	-0.0024 (4)	0.0102 (4)	0.0040 (4)

Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )

O1A—C8A	1.3361 (11)	O1B—C8B	1.3344 (11)
O1A—C9A	1.4512 (11)	O1B—C9B	1.4511 (11)
O2A—C8A	1.2066 (12)	O2B—C8B	1.2041 (12)
N1A—C1A	1.3177 (12)	N1B—C1B	1.3174 (12)
N1A—C2A	1.3649 (12)	N1B—C2B	1.3637 (12)
N2A—C5A	1.3212 (13)	N2B—C5B	1.3183 (13)
N2A—C6A	1.3630 (12)	N2B—C6B	1.3648 (12)
N3A—C7A	1.3163 (12)	N3B—C7B	1.3187 (12)
N3A—C6A	1.3601 (12)	N3B—C6B	1.3608 (12)
C1A—C7A	1.4230 (13)	C1B—C7B	1.4239 (13)
C1A—H1AA	0.9300	C1B—H1BA	0.9300
C2A—C3A	1.4141 (13)	C2B—C3B	1.4146 (13)
C2A—C6A	1.4194 (13)	C2B—C6B	1.4200 (13)
C3A—C4A	1.3650 (14)	C3B—C4B	1.3650 (14)
C3A—H3AA	0.9300	C3B—H3BA	0.9300
C4A—C5A	1.4178 (15)	C4B—C5B	1.4183 (15)
C4A—H4AA	0.9300	C4B—H4BA	0.9300
C5A—H5AA	0.9300	C5B—H5BA	0.9300
C7A—C8A	1.5074 (13)	C7B—C8B	1.5074 (13)
C9A—H9AA	0.9600	C9B—H9BA	0.9600
C9A—H9AB	0.9600	C9B—H9BB	0.9600

C9A—H9AC	0.9600	C9B—H9BC	0.9600
C8A—O1A—C9A	115.74 (7)	C8B—O1B—C9B	115.14 (8)
C1A—N1A—C2A	116.29 (8)	C1B—N1B—C2B	116.38 (8)
C5A—N2A—C6A	116.72 (9)	C5B—N2B—C6B	116.57 (9)
C7A—N3A—C6A	116.38 (8)	C7B—N3B—C6B	116.36 (8)
N1A—C1A—C7A	121.92 (9)	N1B—C1B—C7B	121.96 (9)
N1A—C1A—H1AA	119.0	N1B—C1B—H1BA	119.0
C7A—C1A—H1AA	119.0	C7B—C1B—H1BA	119.0
N1A—C2A—C3A	120.11 (9)	N1B—C2B—C3B	120.05 (9)
N1A—C2A—C6A	121.57 (8)	N1B—C2B—C6B	121.57 (8)
C3A—C2A—C6A	118.31 (9)	C3B—C2B—C6B	118.38 (9)
C4A—C3A—C2A	118.45 (9)	C4B—C3B—C2B	118.13 (9)
C4A—C3A—H3AA	120.8	C4B—C3B—H3BA	120.9
C2A—C3A—H3AA	120.8	C2B—C3B—H3BA	120.9
C3A—C4A—C5A	119.08 (9)	C3B—C4B—C5B	119.37 (9)
C3A—C4A—H4AA	120.5	C3B—C4B—H4BA	120.3
C5A—C4A—H4AA	120.5	C5B—C4B—H4BA	120.3
N2A—C5A—C4A	124.53 (10)	N2B—C5B—C4B	124.55 (9)
N2A—C5A—H5AA	117.7	N2B—C5B—H5BA	117.7
C4A—C5A—H5AA	117.7	C4B—C5B—H5BA	117.7
N3A—C6A—N2A	116.20 (8)	N3B—C6B—N2B	116.04 (8)
N3A—C6A—C2A	120.92 (9)	N3B—C6B—C2B	120.96 (9)
N2A—C6A—C2A	122.88 (9)	N2B—C6B—C2B	122.98 (9)
N3A—C7A—C1A	122.81 (9)	N3B—C7B—C1B	122.73 (8)
N3A—C7A—C8A	115.56 (8)	N3B—C7B—C8B	115.13 (8)
C1A—C7A—C8A	121.59 (8)	C1B—C7B—C8B	122.11 (8)
O2A—C8A—O1A	124.97 (9)	O2B—C8B—O1B	125.19 (9)
O2A—C8A—C7A	124.05 (9)	O2B—C8B—C7B	123.84 (9)
O1A—C8A—C7A	110.96 (8)	O1B—C8B—C7B	110.94 (8)
O1A—C9A—H9AA	109.5	O1B—C9B—H9BA	109.5
O1A—C9A—H9AB	109.5	O1B—C9B—H9BB	109.5
H9AA—C9A—H9AB	109.5	H9BA—C9B—H9BB	109.5
O1A—C9A—H9AC	109.5	O1B—C9B—H9BC	109.5
H9AA—C9A—H9AC	109.5	H9BA—C9B—H9BC	109.5
H9AB—C9A—H9AC	109.5	H9BB—C9B—H9BC	109.5
C2A—N1A—C1A—C7A	1.75 (14)	C2B—N1B—C1B—C7B	0.02 (13)
C1A—N1A—C2A—C3A	-179.51 (9)	C1B—N1B—C2B—C3B	177.88 (9)
C1A—N1A—C2A—C6A	1.27 (13)	C1B—N1B—C2B—C6B	-1.92 (13)
N1A—C2A—C3A—C4A	-178.29 (9)	N1B—C2B—C3B—C4B	179.91 (9)
C6A—C2A—C3A—C4A	0.95 (14)	C6B—C2B—C3B—C4B	-0.28 (14)
C2A—C3A—C4A—C5A	0.57 (14)	C2B—C3B—C4B—C5B	-0.22 (14)
C6A—N2A—C5A—C4A	0.43 (16)	C6B—N2B—C5B—C4B	0.56 (15)
C3A—C4A—C5A—N2A	-1.35 (16)	C3B—C4B—C5B—N2B	0.09 (16)
C7A—N3A—C6A—N2A	-178.69 (9)	C7B—N3B—C6B—N2B	-179.12 (8)
C7A—N3A—C6A—C2A	1.72 (14)	C7B—N3B—C6B—C2B	-0.48 (13)
C5A—N2A—C6A—N3A	-178.34 (9)	C5B—N2B—C6B—N3B	177.50 (8)

C5A—N2A—C6A—C2A	1.24 (15)	C5B—N2B—C6B—C2B	−1.11 (14)
N1A—C2A—C6A—N3A	−3.16 (14)	N1B—C2B—C6B—N3B	2.25 (14)
C3A—C2A—C6A—N3A	177.61 (8)	C3B—C2B—C6B—N3B	−177.55 (9)
N1A—C2A—C6A—N2A	177.28 (9)	N1B—C2B—C6B—N2B	−179.20 (9)
C3A—C2A—C6A—N2A	−1.94 (14)	C3B—C2B—C6B—N2B	0.99 (14)
C6A—N3A—C7A—C1A	1.31 (14)	C6B—N3B—C7B—C1B	−1.43 (13)
C6A—N3A—C7A—C8A	−176.28 (8)	C6B—N3B—C7B—C8B	176.59 (8)
N1A—C1A—C7A—N3A	−3.24 (15)	N1B—C1B—C7B—N3B	1.77 (15)
N1A—C1A—C7A—C8A	174.20 (9)	N1B—C1B—C7B—C8B	−176.12 (9)
C9A—O1A—C8A—O2A	1.37 (14)	C9B—O1B—C8B—O2B	−3.23 (14)
C9A—O1A—C8A—C7A	−176.73 (8)	C9B—O1B—C8B—C7B	175.04 (8)
N3A—C7A—C8A—O2A	−5.04 (14)	N3B—C7B—C8B—O2B	3.54 (14)
C1A—C7A—C8A—O2A	177.34 (9)	C1B—C7B—C8B—O2B	−178.43 (9)
N3A—C7A—C8A—O1A	173.08 (8)	N3B—C7B—C8B—O1B	−174.76 (8)
C1A—C7A—C8A—O1A	−4.54 (12)	C1B—C7B—C8B—O1B	3.27 (12)

*Hydrogen-bond geometry (Å, °)*

D—H···A	D—H	H···A	D···A	D—H···A
C3A—H3AA···O2A <sup>i</sup>	0.93	2.54	3.2401 (13)	133
C4A—H4AA···N2B <sup>ii</sup>	0.93	2.55	3.3311 (14)	141
C9A—H9AA···N1B	0.96	2.62	3.4741 (14)	149
C3B—H3BA···O2B <sup>iii</sup>	0.93	2.53	3.2496 (14)	135
C4B—H4BA···N2A <sup>iv</sup>	0.93	2.51	3.3350 (14)	147
C9B—H9BA···N1A	0.96	2.57	3.4266 (14)	149

Symmetry codes: (i)  $x-1, y, z$ ; (ii)  $-x+1, y-1/2, -z+3/2$ ; (iii)  $x+1, y, z$ ; (iv)  $-x+2, y+1/2, -z+3/2$ .