

**2,5-Dioxopyrrolidin-1-yl 3-(furan-2-yl)-acrylate****Haifeng Zhu, Miao Guo, Han Chen, Haixiao Jin and Hongze Liang\***Faculty of Materials Science and Chemical Engineering, Ningbo University, Ningbo, Zhejiang 315211, People's Republic of China  
Correspondence e-mail: lianghongze@nbu.edu.cn


Received 2 September 2011; accepted 3 October 2011

Key indicators: single-crystal X-ray study;  $T = 296\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$ ;  $R$  factor = 0.035;  $wR$  factor = 0.082; data-to-parameter ratio = 15.6.

The title compound,  $\text{C}_{11}\text{H}_9\text{NO}_5$ , was prepared by the reaction of 2-furanacrylic acid and *N*-hydroxysuccinimide. The molecule consists of two approximately planar moieties, *viz.* a succinimide group and the rest of the molecule [the largest deviations from the least-squares planes are 0.120 (1) and 0.210 (1)  $\text{\AA}$ , respectively]. The dihedral angle between these fragments is 63.70 (5) $^\circ$ . In the crystal, molecules are linked by  $\text{C}-\text{H}\cdots\text{O}$  hydrogen bonds into two-dimensional nets.

**Related literature**

For derivatives of *N*-hydroxysuccinimide, see: Anderson *et al.* (1964); Blumberg & Vallee (1975); Brown *et al.* (2005); Cheng *et al.* (2007); Jones (2003).

**Experimental***Crystal data*
 $\text{C}_{11}\text{H}_9\text{NO}_5$   
 $M_r = 235.19$   
Orthorhombic,  $Pbca$ 
 $a = 10.3054(13)\text{ \AA}$   
 $b = 9.2376(12)\text{ \AA}$   
 $c = 21.892(3)\text{ \AA}$ 
 $V = 2084.0(5)\text{ \AA}^3$   
 $Z = 8$   
Mo  $K\alpha$  radiation  
 $\mu = 0.12\text{ mm}^{-1}$   
 $T = 296\text{ K}$   
 $0.30 \times 0.30 \times 0.10\text{ mm}$ 
*Data collection*
Bruker APEXII CCD area-detector diffractometer  
Absorption correction: multi-scan (*SADABS*; Bruker, 2005)  
 $T_{\min} = 0.977$ ,  $T_{\max} = 0.977$   
16939 measured reflections  
2399 independent reflections  
1900 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.041$ 
*Refinement*
 $R[F^2 > 2\sigma(F^2)] = 0.035$   
 $wR(F^2) = 0.082$   
 $S = 1.02$   
2399 reflections  
154 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.20\text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.24\text{ e \AA}^{-3}$ 

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C12—H12A $\cdots$ O1 <sup>i</sup>	0.93	2.48	3.3533 (17)	156
C14—H14A $\cdots$ O5 <sup>ii</sup>	0.93	2.45	3.3672 (18)	169

Symmetry codes: (i)  $x - \frac{1}{2}, y, -z + \frac{1}{2}$ ; (ii)  $x - \frac{1}{2}, y - 1, -z + \frac{1}{2}$ .

Data collection: *APEX2* (Bruker, 2005); cell refinement: *SAINT* (Bruker, 2005); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

This project was supported by Zhejiang Provincial Natural Science Foundation of China (grant No. Y4100113), the National Natural Science Foundation of China (grant No. 20903058), the Natural Science Foundation of Ningbo City (grant Nos. 2009 A610047, 2010 A610025, 2010 A610027) and the K. C. Wong Magna Fund of Ningbo University.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: YK2020).

**References**

- Anderson, G. W., Callahan, F. M. & Zimmerman, J. E. (1964). *J. Am. Chem. Soc.* **86**, 1839–1842.
- Blumberg, S. & Vallee, B. L. (1975). *Biochemistry*, **14**, 2410–2419.
- Brown, C. L., Atkinson, S. J. & Healy, P. C. (2005). *Acta Cryst. E* **61**, o1203–o1204.
- Bruker (2005). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cheng, F., Gamble, L. J. & Grainger, D. W. (2007). *Anal. Chem.* **79**, 8781–8788.
- Jones, P. G. (2003). *Acta Cryst. E* **59**, o1951–o1952.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

# supporting information

*Acta Cryst.* (2011). E67, o2863 [doi:10.1107/S1600536811040566]

## 2,5-Dioxopyrrolidin-1-yl 3-(furan-2-yl)acrylate

Haifeng Zhu, Miao Guo, Han Chen, Haixiao Jin and Hongze Liang


### S1. Comment

*N*-Hydroxysuccinimide is frequently used in organic chemistry as an activating reagent, which readily form crystalline adducts with amines or acids (Jones, 2003). *N*-Hydroxysuccinimide esters are widely used as leaving groups to activate carboxylic acids (Cheng *et al.*, 2007). The title compound *N*-(2-furanacryloyl)-succinimide ester is an intermediate of FAPGG which is the substrate of diagnostic reagent. We have used a simple procedure to synthesize the title compound in reasonable yields (Blumberg & Vallee, 1975; Brown *et al.*, 2005).

The molecular structure of the title compound (I) is shown in Fig. 1. In the molecule, the dihedral angle between the furan and succinimide rings is 56.26 (43)°. In the crystal structure, molecules are linked by the C12—H12···O1<sup>i</sup> hydrogen bonds to form chains along *a* and C14—H14···O5<sup>ii</sup> hydrogen bonds to form chains along *b* directions.

### S2. Experimental

2-Furanacrylic acid (13.81 g, 0.10 mol), *N*-hydroxysuccinimide (11.51 g, 0.10 mol) and dicyclohexylcarbodiimide (20.63 g, 0.10 mol) were added to 200 ml dioxane in a round flask. This mixture was stirred at 4°C for 14 h before the dicyclohexylurea was removed by filtration. Then the resulting dark brown filtrate was evaporated in vacuum to give the dark brown residue. Slight brown crystals were obtained by recrystallization from 2-propanol (15.29 g, 65%). <sup>1</sup>H NMR(400 MHz, CDCl<sub>3</sub>): δ 7.62 (d, J=16 Hz, 1H), 7.56 (d, J=1.6 Hz, 1H), 6.77 (d, J=3.6 Hz, 1H), 6.53–6.52 (dd, J=3.6 Hz, J=1.6 Hz, 1H), 6.47 (d, J=16 Hz, 1H), 2.87 (s, 4H).



**Figure 1**

The molecular structure of (I) showing displacement ellipsoids at the 30% probability level.

**2,5-Dioxopyrrolidin-1-yl 3-(furan-2-yl)acrylate***Crystal data*

$C_{11}H_9NO_5$	$F(000) = 976$
$M_r = 235.19$	$D_x = 1.499 \text{ Mg m}^{-3}$
Orthorhombic, $Pbca$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -p 2ac 2ab	Cell parameters from 3779 reflections
$a = 10.3054 (13) \text{ \AA}$	$\theta = 2.7\text{--}27.4^\circ$
$b = 9.2376 (12) \text{ \AA}$	$\mu = 0.12 \text{ mm}^{-1}$
$c = 21.892 (3) \text{ \AA}$	$T = 296 \text{ K}$
$V = 2084.0 (5) \text{ \AA}^3$	Needle, brown
$Z = 8$	$0.30 \times 0.30 \times 0.10 \text{ mm}$

*Data collection*

Bruker APEXII CCD area-detector diffractometer	16939 measured reflections
Radiation source: fine-focus sealed tube	2399 independent reflections
Graphite monochromator	1900 reflections with $I > 2\sigma(I)$
Detector resolution: 0 pixels $\text{mm}^{-1}$	$R_{\text{int}} = 0.041$
$\varphi$ and $\omega$ scans	$\theta_{\text{max}} = 27.6^\circ, \theta_{\text{min}} = 1.9^\circ$
Absorption correction: multi-scan (SADABS; Bruker, 2005)	$h = -13 \rightarrow 13$
$T_{\text{min}} = 0.977, T_{\text{max}} = 0.977$	$k = -12 \rightarrow 12$
	$l = -28 \rightarrow 28$

*Refinement*

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.035$	H-atom parameters constrained
$wR(F^2) = 0.082$	$w = 1/[\sigma^2(F_o^2) + (0.0285P)^2 + 0.9232P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.02$	$(\Delta/\sigma)_{\text{max}} < 0.001$
2399 reflections	$\Delta\rho_{\text{max}} = 0.20 \text{ e \AA}^{-3}$
154 parameters	$\Delta\rho_{\text{min}} = -0.24 \text{ e \AA}^{-3}$
0 restraints	
Primary atom site location: structure-invariant direct methods	

*Special details*

**Geometry.** All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^* / U_{\text{eq}}$
O1	0.80658 (9)	0.22692 (10)	0.28201 (4)	0.0261 (2)
O2	0.89760 (9)	0.55049 (10)	0.10826 (4)	0.0273 (2)
O3	0.67881 (9)	0.58836 (11)	0.11454 (5)	0.0314 (2)
O4	0.81461 (10)	0.44760 (11)	-0.00431 (5)	0.0356 (3)

O5	1.01727 (10)	0.81990 (11)	0.09170 (5)	0.0349 (3)
C6	0.68207 (13)	0.36659 (14)	0.21024 (6)	0.0237 (3)
H6A	0.6008	0.3944	0.1960	0.028*
C7	0.78634 (13)	0.42777 (14)	0.18443 (6)	0.0244 (3)
H7A	0.8686	0.4048	0.1989	0.029*
N8	0.89355 (11)	0.63213 (12)	0.05495 (5)	0.0246 (3)
C9	0.68679 (12)	0.26114 (14)	0.25831 (6)	0.0220 (3)
C10	0.77299 (13)	0.52965 (14)	0.13407 (6)	0.0233 (3)
C11	0.96337 (13)	0.76125 (14)	0.04991 (6)	0.0243 (3)
C12	0.59420 (14)	0.18035 (14)	0.28687 (6)	0.0255 (3)
H12A	0.5054	0.1829	0.2793	0.031*
C13	0.85503 (13)	0.56972 (15)	0.00020 (6)	0.0250 (3)
C14	0.65847 (14)	0.09163 (15)	0.33025 (6)	0.0270 (3)
H14A	0.6205	0.0249	0.3566	0.032*
C15	0.78534 (14)	0.12362 (15)	0.32556 (6)	0.0279 (3)
H15A	0.8503	0.0810	0.3489	0.034*
C16	0.87575 (15)	0.68378 (15)	-0.04765 (6)	0.0289 (3)
H16A	0.7934	0.7229	-0.0614	0.035*
H16B	0.9214	0.6439	-0.0826	0.035*
C17	0.95729 (14)	0.80146 (15)	-0.01663 (6)	0.0286 (3)
H17A	1.0438	0.8046	-0.0341	0.034*
H17B	0.9170	0.8956	-0.0217	0.034*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O1	0.0231 (5)	0.0287 (5)	0.0264 (5)	-0.0003 (4)	-0.0014 (4)	0.0072 (4)
O2	0.0248 (5)	0.0329 (5)	0.0242 (5)	-0.0006 (4)	-0.0008 (4)	0.0097 (4)
O3	0.0272 (5)	0.0337 (5)	0.0334 (5)	0.0032 (4)	-0.0018 (4)	0.0105 (4)
O4	0.0398 (6)	0.0254 (5)	0.0416 (6)	-0.0037 (5)	-0.0043 (5)	-0.0042 (5)
O5	0.0351 (6)	0.0322 (5)	0.0376 (6)	-0.0055 (5)	-0.0022 (5)	-0.0085 (5)
C6	0.0264 (7)	0.0222 (6)	0.0226 (6)	0.0026 (5)	-0.0019 (5)	-0.0008 (5)
C7	0.0253 (7)	0.0243 (7)	0.0235 (7)	0.0016 (6)	-0.0028 (5)	0.0022 (5)
N8	0.0284 (6)	0.0253 (6)	0.0199 (5)	-0.0037 (5)	-0.0009 (5)	0.0052 (5)
C9	0.0222 (6)	0.0232 (6)	0.0208 (6)	0.0025 (5)	-0.0012 (5)	-0.0009 (5)
C10	0.0245 (7)	0.0215 (6)	0.0238 (6)	-0.0014 (6)	-0.0009 (5)	0.0000 (5)
C11	0.0211 (6)	0.0211 (6)	0.0309 (7)	0.0008 (5)	0.0032 (6)	-0.0016 (6)
C12	0.0245 (7)	0.0251 (7)	0.0268 (7)	-0.0017 (6)	0.0018 (5)	-0.0016 (5)
C13	0.0239 (7)	0.0243 (7)	0.0269 (7)	0.0037 (6)	-0.0016 (6)	-0.0025 (5)
C14	0.0363 (8)	0.0227 (6)	0.0220 (6)	-0.0047 (6)	0.0031 (6)	0.0010 (5)
C15	0.0359 (8)	0.0253 (7)	0.0226 (6)	0.0014 (6)	-0.0030 (6)	0.0065 (6)
C16	0.0355 (8)	0.0283 (7)	0.0229 (7)	0.0055 (6)	0.0013 (6)	0.0007 (6)
C17	0.0280 (7)	0.0246 (7)	0.0333 (8)	0.0012 (6)	0.0048 (6)	0.0068 (6)

*Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )*

O1—C15	1.3664 (16)	C9—C12	1.3632 (18)
O1—C9	1.3759 (15)	C11—C17	1.5044 (19)

O2—N8	1.3900 (13)	C12—C14	1.4185 (19)
O2—C10	1.4162 (16)	C12—H12A	0.9300
O3—C10	1.1912 (16)	C13—C16	1.5011 (19)
O4—C13	1.2066 (16)	C14—C15	1.344 (2)
O5—C11	1.1996 (16)	C14—H14A	0.9300
C6—C7	1.3391 (18)	C15—H15A	0.9300
C6—C9	1.4348 (18)	C16—C17	1.533 (2)
C6—H6A	0.9300	C16—H16A	0.9700
C7—C10	1.4561 (18)	C16—H16B	0.9700
C7—H7A	0.9300	C17—H17A	0.9700
N8—C13	1.3880 (17)	C17—H17B	0.9700
N8—C11	1.3974 (17)		
C15—O1—C9	106.25 (10)	C14—C12—H12A	126.4
N8—O2—C10	112.43 (9)	O4—C13—N8	123.89 (13)
C7—C6—C9	124.66 (12)	O4—C13—C16	130.41 (12)
C7—C6—H6A	117.7	N8—C13—C16	105.69 (11)
C9—C6—H6A	117.7	C15—C14—C12	106.01 (12)
C6—C7—C10	121.09 (12)	C15—C14—H14A	127.0
C6—C7—H7A	119.5	C12—C14—H14A	127.0
C10—C7—H7A	119.5	C14—C15—O1	111.26 (12)
C13—N8—O2	120.55 (11)	C14—C15—H15A	124.4
C13—N8—C11	115.69 (11)	O1—C15—H15A	124.4
O2—N8—C11	120.93 (11)	C13—C16—C17	105.46 (11)
C12—C9—O1	109.22 (11)	C13—C16—H16A	110.6
C12—C9—C6	133.19 (13)	C17—C16—H16A	110.6
O1—C9—C6	117.58 (11)	C13—C16—H16B	110.6
O3—C10—O2	122.25 (12)	C17—C16—H16B	110.6
O3—C10—C7	130.02 (13)	H16A—C16—H16B	108.8
O2—C10—C7	107.72 (11)	C11—C17—C16	106.07 (11)
O5—C11—N8	124.31 (13)	C11—C17—H17A	110.5
O5—C11—C17	130.25 (13)	C16—C17—H17A	110.5
N8—C11—C17	105.42 (11)	C11—C17—H17B	110.5
C9—C12—C14	107.25 (12)	C16—C17—H17B	110.5
C9—C12—H12A	126.4	H17A—C17—H17B	108.7

*Hydrogen-bond geometry (Å, °)*

D—H···A	D—H	H···A	D···A	D—H···A
C12—H12A···O1 <sup>i</sup>	0.93	2.48	3.3533 (17)	156
C14—H14A···O5 <sup>ii</sup>	0.93	2.45	3.3672 (18)	169

Symmetry codes: (i)  $x-1/2, y, -z+1/2$ ; (ii)  $x-1/2, y-1, -z+1/2$ .