

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

2-(1,3-Benzoxazol-2-yl)guanidinium chloride

 Shaaban K. Mohamed,^a Peter N. Horton,^b Mahmoud A.A. El-Remaily^c and Seik Weng Ng^{d,e,*}

^aChemistry and Environmental Division, Manchester Metropolitan University, Manchester M15 6BH, England, ^bSchool of Chemistry, University of Southampton, Southampton SO17 1BJ, England, ^cDepartment of Chemistry, Faculty of Science, Sohag University, Egypt, ^dDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia, and ^eChemistry Department, King Abdulaziz University, PO Box 80203 Jeddah, Saudi Arabia

Correspondence e-mail: seikweng@um.edu.my


Received 17 October 2011; accepted 25 October 2011

 Key indicators: single-crystal X-ray study; $T = 120$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.036; wR factor = 0.089; data-to-parameter ratio = 14.2.

The non-H atoms of the cation of the title salt, $\text{C}_8\text{H}_9\text{N}_4\text{O}^+\cdot\text{Cl}^-$, are approximately co-planar (r.m.s. deviation = 0.024 Å) with one amino group forming an intramolecular hydrogen bond to the tertiary N atom of the benzoxazole fused-ring system. The cations and anions are linked by cyclic $R_2^1(6)$ N—H...Cl hydrogen-bonding associations, generating linear chains running along the a -axis direction.

Related literature

For the synthesis, see: Takahashi & Niino (1943). For the structure of a co-crystal of 2-(1,3-benzoxazol-2-yl)guanidine, see: Bishop *et al.* (2005) and for the structure of 2-(1,3-benzothiazol-2-yl)guanidine, see: Mohamed *et al.* (2011). For graph-set analysis, see: Etter *et al.* (1990).



Experimental

Crystal data

$\text{C}_8\text{H}_9\text{N}_4\text{O}^+\cdot\text{Cl}^-$
 $M_r = 212.64$
 Triclinic, $P\bar{1}$
 $a = 6.669$ (2) Å
 $b = 8.152$ (4) Å
 $c = 9.630$ (3) Å

$\alpha = 65.062$ (2)°
 $\beta = 85.020$ (2)°
 $\gamma = 73.710$ (2)°
 $V = 455.4$ (3) Å³
 $Z = 2$
 Mo $K\alpha$ radiation

$\mu = 0.39$ mm⁻¹
 $T = 120$ K

0.20 × 0.05 × 0.04 mm

Data collection

Bruker–Nonius Roper CCD camera on κ -goniostat diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.926$, $T_{\max} = 0.985$

8184 measured reflections
 2088 independent reflections
 1895 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.044$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.036$
 $wR(F^2) = 0.089$
 $S = 1.03$
 2088 reflections
 147 parameters
 5 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.31$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.26$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N2—H1...Cl1	0.88 (1)	2.18 (1)	3.047 (2)	168 (2)
N3—H2...Cl1	0.88 (1)	2.67 (2)	3.415 (2)	144 (2)
N3—H3...Cl1 ⁱ	0.87 (1)	2.53 (2)	3.297 (2)	147 (2)
N4—H4...Cl1 ⁱ	0.88 (1)	2.33 (1)	3.168 (2)	160 (2)
N4—H5...N1	0.88 (1)	2.08 (2)	2.765 (2)	134 (2)

 Symmetry code: (i) $x + 1, y, z$.

Data collection: COLLECT (Hooft, 1998); cell refinement: DENZO (Otwinowski & Minor, 1997) and COLLECT; data reduction: DENZO; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: X-SEED (Barbour, 2001); software used to prepare material for publication: publCIF (Westrip, 2010).

The use of the EPSRC X-ray crystallographic facilities at the University of Southampton, England, is gratefully acknowledged. We thank Manchester Metropolitan University, Sohag University and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: ZS2158).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
 Bishop, M. M., Coles, S. J., Lee, A. H. W., Lindoy, L. F., Parkin, A., Thorn-Seshold, O. T. & Turner, P. (2005). *Supramol. Chem.* **17**, 567–578.
 Etter, M. C., MacDonald, J. C. & Bernstein, J. (1990). *Acta Cryst.* **B46**, 256–262.
 Hooft, R. (1998). COLLECT. Nonius BV, Delft, The Netherlands.
 Mohamed, S. K., Horton, P. N., El-Remaily, M. A. A. & Ng, S. W. (2011). *Acta Cryst.* **E67**, o3132.
 Otwinowski, O. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
 Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Takahashi, T. & Niino, A. (1943). *Yakugaku Zasshi*, **63**, 249–252.
 Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2011). E67, o3133 [doi:10.1107/S1600536811044655]

2-(1,3-Benzoxazol-2-yl)guanidinium chloride

Shaaban K. Mohamed, Peter N. Horton, Mahmoud A.A. El-Remaily and Seik Weng Ng

S1. Comment


The preceding study reports 2-(1,3-benzothiazol-2-yl)guanidinium chloride (Mohamed *et al.*, 2011). Replacing the sulfur in the fused-ring by oxygen leads to the analogous compound, 2-(1,3-benzoxazol-2-yl)guanidinium chloride (Scheme 1). However, this salt and 2-(1,3-benzothiazol-2-yl)guanidinium chloride are not isostructural as they belong to different crystal systems. The non-H atoms of the cation of the title salt, $C_8H_9N_4O^+ Cl^-$ (Fig. 1), lie on a plane with one amino group forming an intramolecular hydrogen bond to the tertiary N atom of the benzoxazole fused-ring. The cations and anions are linked by cyclic $R^1_2(6) N-H\cdots Cl$ hydrogen-bonding associations [Etter *et al.*, 1990], to generate linear chains running along the *a*-axis of the triclinic unit cell (Table 1). This salt was first reported in 1943 (Takahashi & Niino, 1943). The structure of a co-crystal of 2-(1,3-benzoxazol-2-yl)guanidine has been reported (Bishop *et al.*, 2005).

S2. Experimental

2-(1,3-Benzoxazol-2-yl)guanidine was synthesized by using a literature procedure similar to that used for synthesizing 2-(1,3-benzothiazol-2-yl)guanidine (Takahashi & Niino, 1943). The guanidine (0.05 mol) was heated in ethanol (50 ml) in the presence of a few drops of hydrochloric acid for 3 h. The mixture was cooled and the product was recrystallized from ethanol to give the title compound (m.p. 538 K) in 95% yield.

S3. Refinement

Carbon-bound H-atoms were placed in calculated positions (C—H 0.95 Å) and were included in the refinement in the riding model approximation, with $U_{iso}(H)$ set to $1.2U_{eq}(C)$. The amino H-atoms were located in a difference Fourier map, and were refined with a distance restraint of N—H = 0.88 ± 0.01 Å, with their isotropic displacement parameters freely refining.

**Figure 1**

Thermal ellipsoid plot (Barbour, 2001) of $C_8H_9N_4O^+ Cl^-$ at the 70% probability level. Hydrogen atoms are drawn as spheres of arbitrary radius.

2-(1,3-Benzoxazol-2-yl)guanidinium chloride

Crystal data

$C_8H_9N_4O^+ Cl^-$

$M_r = 212.64$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 6.669 (2) \text{ \AA}$

$b = 8.152 (4) \text{ \AA}$

$c = 9.630 (3) \text{ \AA}$

$\alpha = 65.062 (2)^\circ$

$\beta = 85.020 (2)^\circ$

$\gamma = 73.710 (2)^\circ$

$V = 455.4 (3) \text{ \AA}^3$

$Z = 2$

$F(000) = 220$

$D_x = 1.551 \text{ Mg m}^{-3}$

Melting point: 538 K

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 1979 reflections

$\theta = 2.9\text{--}27.5^\circ$

$\mu = 0.39 \text{ mm}^{-1}$

$T = 120 \text{ K}$

Prism, colorless

$0.20 \times 0.05 \times 0.04 \text{ mm}$

Data collection

Bruker-Nonius Roper CCD camera on κ -goniostat diffractometer

Radiation source: Bruker-Nonius FR591 rotating anode

Graphite monochromator

Detector resolution: $9.091 \text{ pixels mm}^{-1}$

φ and ω scans

Absorption correction: multi-scan (SADABS; Sheldrick, 1996)

$T_{\min} = 0.926$, $T_{\max} = 0.985$

8184 measured reflections

2088 independent reflections

1895 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.044$

$\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 3.2^\circ$

$h = -8 \rightarrow 8$

$k = -10 \rightarrow 10$

$l = -12 \rightarrow 12$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.036$
 $wR(F^2) = 0.089$
 $S = 1.03$
 2088 reflections
 147 parameters
 5 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H atoms treated by a mixture of independent
 and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0256P)^2 + 0.4591P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.31 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.26 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	0.25918 (6)	0.37898 (6)	0.41313 (5)	0.01998 (13)
O1	0.50373 (17)	-0.01606 (16)	0.27313 (13)	0.0176 (3)
N1	0.8440 (2)	-0.0414 (2)	0.21101 (16)	0.0179 (3)
N2	0.6679 (2)	0.1505 (2)	0.34140 (17)	0.0186 (3)
N3	0.7839 (2)	0.3256 (2)	0.43342 (18)	0.0215 (3)
N4	1.0159 (2)	0.1588 (2)	0.31608 (18)	0.0212 (3)
C1	0.5575 (3)	-0.1365 (2)	0.19970 (19)	0.0171 (3)
C2	0.4318 (3)	-0.2237 (2)	0.1669 (2)	0.0197 (3)
H2A	0.2904	-0.2110	0.1960	0.024*
C3	0.5268 (3)	-0.3327 (2)	0.0875 (2)	0.0224 (4)
H3A	0.4482	-0.3975	0.0619	0.027*
C4	0.7347 (3)	-0.3490 (2)	0.0448 (2)	0.0221 (4)
H4A	0.7930	-0.4229	-0.0106	0.027*
C5	0.8592 (3)	-0.2595 (2)	0.0815 (2)	0.0210 (4)
H5A	1.0010	-0.2719	0.0533	0.025*
C6	0.7658 (3)	-0.1517 (2)	0.16100 (19)	0.0178 (3)
C7	0.6845 (2)	0.0303 (2)	0.27334 (19)	0.0171 (3)
C8	0.8270 (2)	0.2129 (2)	0.36229 (19)	0.0169 (3)
H1	0.544 (2)	0.201 (3)	0.367 (3)	0.035 (6)*
H2	0.654 (2)	0.367 (4)	0.454 (3)	0.046 (7)*
H3	0.881 (3)	0.372 (3)	0.446 (3)	0.039 (7)*
H4	1.110 (3)	0.205 (3)	0.334 (2)	0.025 (5)*
H5	1.035 (4)	0.085 (3)	0.268 (2)	0.035 (6)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl1	0.0149 (2)	0.0214 (2)	0.0280 (2)	-0.00545 (15)	0.00285 (15)	-0.01445 (17)

O1	0.0145 (5)	0.0191 (6)	0.0231 (6)	-0.0051 (4)	0.0008 (4)	-0.0122 (5)
N1	0.0169 (7)	0.0190 (7)	0.0208 (7)	-0.0048 (5)	0.0015 (5)	-0.0111 (6)
N2	0.0133 (7)	0.0216 (7)	0.0259 (8)	-0.0041 (5)	0.0015 (5)	-0.0151 (6)
N3	0.0182 (7)	0.0238 (8)	0.0295 (8)	-0.0066 (6)	0.0011 (6)	-0.0170 (7)
N4	0.0162 (7)	0.0250 (8)	0.0288 (8)	-0.0068 (6)	0.0028 (6)	-0.0169 (7)
C1	0.0188 (8)	0.0156 (8)	0.0171 (8)	-0.0032 (6)	-0.0001 (6)	-0.0076 (6)
C2	0.0180 (8)	0.0196 (8)	0.0215 (8)	-0.0062 (6)	0.0008 (6)	-0.0079 (7)
C3	0.0241 (9)	0.0211 (9)	0.0249 (9)	-0.0074 (7)	-0.0021 (7)	-0.0110 (7)
C4	0.0247 (9)	0.0208 (9)	0.0232 (9)	-0.0040 (7)	0.0007 (7)	-0.0127 (7)
C5	0.0195 (8)	0.0221 (9)	0.0225 (9)	-0.0039 (7)	0.0025 (7)	-0.0118 (7)
C6	0.0173 (8)	0.0176 (8)	0.0194 (8)	-0.0049 (6)	-0.0002 (6)	-0.0081 (7)
C7	0.0147 (7)	0.0175 (8)	0.0199 (8)	-0.0046 (6)	-0.0009 (6)	-0.0079 (7)
C8	0.0153 (7)	0.0170 (8)	0.0176 (8)	-0.0037 (6)	-0.0005 (6)	-0.0067 (7)

Geometric parameters (Å, °)

O1—C7	1.361 (2)	N4—H5	0.879 (10)
O1—C1	1.391 (2)	C1—C2	1.371 (2)
N1—C7	1.292 (2)	C1—C6	1.391 (2)
N1—C6	1.409 (2)	C2—C3	1.395 (3)
N2—C8	1.361 (2)	C2—H2A	0.9500
N2—C7	1.368 (2)	C3—C4	1.397 (3)
N2—H1	0.879 (10)	C3—H3A	0.9500
N3—C8	1.321 (2)	C4—C5	1.398 (2)
N3—H2	0.875 (10)	C4—H4A	0.9500
N3—H3	0.874 (10)	C5—C6	1.387 (2)
N4—C8	1.318 (2)	C5—H5A	0.9500
N4—H4	0.876 (9)		
C7—O1—C1	102.84 (12)	C2—C3—H3A	119.2
C7—N1—C6	102.87 (14)	C4—C3—H3A	119.2
C8—N2—C7	125.13 (14)	C3—C4—C5	121.75 (16)
C8—N2—H1	115.8 (16)	C3—C4—H4A	119.1
C7—N2—H1	118.9 (16)	C5—C4—H4A	119.1
C8—N3—H2	119.2 (18)	C6—C5—C4	116.77 (16)
C8—N3—H3	119.4 (16)	C6—C5—H5A	121.6
H2—N3—H3	121 (2)	C4—C5—H5A	121.6
C8—N4—H4	115.4 (14)	C5—C6—C1	119.91 (16)
C8—N4—H5	118.0 (16)	C5—C6—N1	131.01 (16)
H4—N4—H5	127 (2)	C1—C6—N1	109.06 (14)
C2—C1—O1	127.63 (15)	N1—C7—O1	117.68 (15)
C2—C1—C6	124.81 (16)	N1—C7—N2	129.18 (15)
O1—C1—C6	107.55 (14)	O1—C7—N2	113.14 (14)
C1—C2—C3	115.05 (16)	N4—C8—N3	122.14 (16)
C1—C2—H2A	122.5	N4—C8—N2	120.88 (15)
C3—C2—H2A	122.5	N3—C8—N2	116.97 (15)
C2—C3—C4	121.70 (16)		

C7—O1—C1—C2	-179.20 (17)	O1—C1—C6—N1	0.35 (18)
C7—O1—C1—C6	-0.35 (17)	C7—N1—C6—C5	177.86 (18)
O1—C1—C2—C3	178.07 (15)	C7—N1—C6—C1	-0.19 (18)
C6—C1—C2—C3	-0.6 (3)	C6—N1—C7—O1	0.0 (2)
C1—C2—C3—C4	-0.4 (3)	C6—N1—C7—N2	-179.71 (17)
C2—C3—C4—C5	1.1 (3)	C1—O1—C7—N1	0.26 (19)
C3—C4—C5—C6	-0.7 (3)	C1—O1—C7—N2	179.97 (14)
C4—C5—C6—C1	-0.2 (3)	C8—N2—C7—N1	-3.1 (3)
C4—C5—C6—N1	-178.12 (17)	C8—N2—C7—O1	177.25 (15)
C2—C1—C6—C5	0.9 (3)	C7—N2—C8—N4	0.0 (3)
O1—C1—C6—C5	-177.95 (15)	C7—N2—C8—N3	-178.54 (16)
C2—C1—C6—N1	179.24 (16)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N2—H1...C11	0.88 (1)	2.18 (1)	3.047 (2)	168 (2)
N3—H2...C11	0.88 (1)	2.67 (2)	3.415 (2)	144 (2)
N3—H3...C11 ⁱ	0.87 (1)	2.53 (2)	3.297 (2)	147 (2)
N4—H4...C11 ⁱ	0.88 (1)	2.33 (1)	3.168 (2)	160 (2)
N4—H5...N1	0.88 (1)	2.08 (2)	2.765 (2)	134 (2)

Symmetry code: (i) $x+1, y, z$.