

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Ethyl 2-(4-nitrophenyl)-1-[3-(2-oxopyrrolidin-1-yl)propyl]-1H-benzimidazole-5-carboxylate

 Yeong Keng Yoon,^a Mohamed Ashraf Ali,^a Tan Soo Choon,^a Safra Izuani Jama Asik^b and Ibrahim Abdul Razak^{b*‡}
^aInstitute for Research in Molecular Medicine, Universiti Sains Malaysia, Minden 11800, Penang, Malaysia, and ^bSchool of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia

Correspondence e-mail: arazaki@usm.my


Received 30 November 2011; accepted 5 December 2011

 Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.069; wR factor = 0.197; data-to-parameter ratio = 25.8.

In the title compound, $\text{C}_{23}\text{H}_{24}\text{N}_4\text{O}_5$, the essentially planar benzimidazole ring system [maximum deviation = $0.008(2)$ Å] forms a dihedral angle of $39.22(7)^\circ$ with the attached nitrobenzene ring. The pyrrolidin-2-one ring adopts an envelope conformation with a C atom as the flap. In the crystal, molecules are connected by $\text{C}-\text{H}\cdots\text{O}$ interactions, forming sheets propagating in (011). The crystal packing also features weak $\pi-\pi$ stacking interactions [centroid-centroid = $3.6746(12)$ Å].

Related literature

For applications of benzimidazole compounds, see: Rao *et al.* (2002); Ali *et al.* (2007). For the stability of the temperature controller used in the data collection, see: Cosier & Glazer (1986). For ring conformations, see: Cremer & Pople (1975).



Experimental

Crystal data

 $\text{C}_{23}\text{H}_{24}\text{N}_4\text{O}_5$
 $M_r = 436.46$
[‡] Thomson Reuters ResearcherID: A-5599-2009.

 Triclinic, $P\bar{1}$
 $a = 9.3125(1)$ Å
 $b = 10.0941(1)$ Å
 $c = 12.9955(2)$ Å
 $\alpha = 91.958(1)^\circ$
 $\beta = 107.752(1)^\circ$
 $\gamma = 114.465(1)^\circ$
 $V = 1040.67(2)$ Å³
 $Z = 2$

 Mo $K\alpha$ radiation

 $\mu = 0.10$ mm⁻¹
 $T = 100$ K

 $0.34 \times 0.20 \times 0.13$ mm

Data collection

 Bruker SMART APEXII CCD diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2009)
 $T_{\min} = 0.967$, $T_{\max} = 0.987$

 27734 measured reflections
 7466 independent reflections
 5375 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.032$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.069$
 $wR(F^2) = 0.197$
 $S = 1.04$

7466 reflections

289 parameters

H-atom parameters constrained

 $\Delta\rho_{\text{max}} = 0.50$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.30$ e Å⁻³
Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C15}-\text{H15A}\cdots\text{O3}^{\text{i}}$	0.99	2.41	3.355 (3)	159
$\text{C15}-\text{H15B}\cdots\text{O3}^{\text{ii}}$	0.99	2.38	3.186 (3)	139
$\text{C19}-\text{H19A}\cdots\text{O2}^{\text{iii}}$	0.99	2.38	3.312 (3)	156

 Symmetry codes: (i) $x - 1, y - 1, z - 1$; (ii) $-x, -y, -z + 1$; (iii) $-x - 1, -y, -z + 1$.

Data collection: APEX2 (Bruker, 2009); cell refinement: SAINT (Bruker, 2009); data reduction: SAINT; program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL and PLATON (Spek, 2009).

The authors thank the Malaysian Government and Universiti Sains Malaysia for the Research University Grant (Nos. 1001/PFIZIK/811151 and 1001/PSK/8620012). The authors also wish to express their thanks to the Pharmacogenetic and Novel Therapeutic Research, Institute for Research in Molecular Medicine, Universiti of Sains Malaysia.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB6545).

References

- Ali, M. A., Shahar Yar, M. & Siddiqui, A. A. (2007). *Eur. J. Med. Chem.* **42**, 268–275.
 Bruker (2009). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
 Cosier, J. & Glazer, A. M. (1986). *J. Appl. Cryst.* **19**, 105–107.
 Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
 Rao, A., Chimirri, A., Clercq, E. D., Monforte, A. M., Monforte, P., Pannecouque, C. & Zappala, M. (2002). *Farmaco*, **57**, 819–823.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

supporting information

Acta Cryst. (2012). E68, o59 [doi:10.1107/S1600536811052408]

Ethyl 2-(4-nitrophenyl)-1-[3-(2-oxopyrrolidin-1-yl)propyl]-1*H*-benzimidazole-5-carboxylate

Yeong Keng Yoon, Mohamed Ashraf Ali, Tan Soo Choon, Safra Izuani Jama Asik and Ibrahim Abdul Razak

S1. Comment

Benzimidazole derivatives have many clinical applications (Rao *et al.*, 2002) and are being currently evaluated for their anti-TB activities (Ali *et al.*, 2007).

In the title compound, (I), Fig. 1, the benzimidazole (N1–N2/C1–C7) ring is essentially planar with maximum deviation of 0.008 (2) for atom C3. This ring also makes dihedral angles of 77.02 (11) and 39.22 (7)° with the mean plane of pyrrolidin-2-one, (O3/N3/C20–C23) and nitrobenzene, (C8–C13) groups, respectively. The mean plane of pyrrolidin-2-one, (O3/N3/C20–C23) adopts an envelope conformation with puckering parameters $Q = 0.237$ (3) Å and $\varphi = 76.0$ (6)° (Cremer & Pople, 1975).


In the crystal (Fig. 2), the molecules are connected by C15—H15A···O3, C15—H15B···O3 and C19—H19A···O2 interactions to form (011) sheets. π – π stacking interactions are observed between the benzene (C1–C6; centroid Cg3) rings with their centroids distance of 3.6746 (12) Å [symmetry code : -x,-y,1-z].

S2. Experimental


Ethyl 3-amino-4-(3(2-oxopyrrolidin-1-yl)propylamino)benzoate (0.84 mmol) and sodium metabisulfite adduct of 4-nitrobenzaldehyde (1.68 mmol) were dissolved in DMF. The reaction mixture was reflux at 130 °C for 2 hrs. After completion, the reaction mixture was diluted in ethyl acetate (20 ml) and washed with water (20 ml). The organic layer was collected, dried over Na₂SO₄ and the evaporated in *vacuo* to yield the product. The product was recrystallised from ethyl acetate as orange blocks.

S3. Refinement

All the H atoms were positioned geometrically and refined using a riding model with with C–H = 0.95–0.99 Å. The U_{iso} values were constrained to be 1.5 U_{eq} of the carrier atom for methyl H atoms and 1.2 U_{eq} for the remaining H atoms.

**Figure 1**

The structure of the title compound, showing 50% probability displacement ellipsoids.

**Figure 2**

The crystal packing, viewed along the *a*-axis, showing two-dimensional planes parallel to (011). Hydrogen atoms that not involved in hydrogen bonding (dashed lines) are omitted for clarity.

Ethyl 2-(4-nitrophenyl)-1-[3-(2-oxopyrrolidin-1-yl)propyl]-1H-benzimidazole-5-carboxylate

Crystal data

$C_{23}H_{24}N_4O_5$	$Z = 2$
$M_r = 436.46$	$F(000) = 460$
Triclinic, $P\bar{1}$	$D_x = 1.393 \text{ Mg m}^{-3}$
Hall symbol: -P 1	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$a = 9.3125 (1) \text{ \AA}$	Cell parameters from 8871 reflections
$b = 10.0941 (1) \text{ \AA}$	$\theta = 2.5\text{--}32.6^\circ$
$c = 12.9955 (2) \text{ \AA}$	$\mu = 0.10 \text{ mm}^{-1}$
$\alpha = 91.958 (1)^\circ$	$T = 100 \text{ K}$
$\beta = 107.752 (1)^\circ$	Block, orange
$\gamma = 114.465 (1)^\circ$	$0.34 \times 0.20 \times 0.13 \text{ mm}$
$V = 1040.67 (2) \text{ \AA}^3$	

Data collection

Bruker SMART APEXII CCD diffractometer	27734 measured reflections
Radiation source: fine-focus sealed tube	7466 independent reflections
Graphite monochromator	5375 reflections with $I > 2\sigma(I)$
φ and ω scans	$R_{\text{int}} = 0.032$
Absorption correction: multi-scan (SADABS; Bruker, 2009)	$\theta_{\text{max}} = 32.6^\circ$, $\theta_{\text{min}} = 1.7^\circ$
$T_{\text{min}} = 0.967$, $T_{\text{max}} = 0.987$	$h = -14 \rightarrow 14$
	$k = -15 \rightarrow 15$
	$l = -19 \rightarrow 19$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.069$	H-atom parameters constrained
$wR(F^2) = 0.197$	$w = 1/[\sigma^2(F_o^2) + (0.0843P)^2 + 0.8488P]$
$S = 1.04$	where $P = (F_o^2 + 2F_c^2)/3$
7466 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
289 parameters	$\Delta\rho_{\text{max}} = 0.50 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.30 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	

Special details

Experimental. The crystal was placed in the cold stream of an Oxford Cryosystems Cobra open-flow nitrogen cryostat (Cosier & Glazer, 1986) operating at 100.0 (1) K.

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	-0.29047 (16)	-0.18231 (15)	0.17443 (11)	0.0271 (3)
O2	-0.45664 (18)	-0.31986 (16)	0.26240 (12)	0.0328 (3)

O3	0.2053 (2)	0.51173 (19)	0.95685 (13)	0.0385 (4)
O4	0.82578 (18)	0.95186 (18)	0.77738 (14)	0.0371 (4)
O5	0.6533 (2)	1.04206 (17)	0.78208 (14)	0.0388 (4)
N1	0.11744 (17)	0.29815 (16)	0.46007 (12)	0.0201 (3)
N2	0.00661 (17)	0.29678 (16)	0.59278 (11)	0.0193 (3)
N3	-0.0066 (2)	0.56743 (18)	0.86162 (13)	0.0245 (3)
N4	0.6825 (2)	0.93979 (18)	0.75733 (13)	0.0268 (3)
C1	-0.0276 (2)	0.16584 (19)	0.43862 (14)	0.0189 (3)
C2	-0.1050 (2)	0.04589 (19)	0.35168 (14)	0.0201 (3)
H2A	-0.0586	0.0455	0.2959	0.024*
C3	-0.2518 (2)	-0.07262 (19)	0.34957 (14)	0.0208 (3)
C4	-0.3197 (2)	-0.0732 (2)	0.43315 (15)	0.0233 (3)
H4A	-0.4194	-0.1570	0.4299	0.028*
C5	-0.2451 (2)	0.0444 (2)	0.51921 (14)	0.0224 (3)
H5A	-0.2914	0.0443	0.5751	0.027*
C6	-0.0981 (2)	0.16403 (19)	0.52027 (14)	0.0194 (3)
C7	0.1333 (2)	0.37277 (18)	0.55203 (14)	0.0185 (3)
C8	0.2713 (2)	0.52054 (19)	0.60439 (13)	0.0193 (3)
C9	0.2460 (2)	0.6325 (2)	0.65125 (14)	0.0217 (3)
H9A	0.1366	0.6135	0.6500	0.026*
C10	0.3803 (2)	0.7712 (2)	0.69957 (14)	0.0221 (3)
H10A	0.3638	0.8478	0.7310	0.026*
C11	0.5388 (2)	0.79566 (19)	0.70104 (14)	0.0216 (3)
C12	0.5674 (2)	0.6891 (2)	0.65223 (15)	0.0237 (3)
H12A	0.6764	0.7100	0.6517	0.028*
C13	0.4322 (2)	0.5512 (2)	0.60426 (14)	0.0221 (3)
H13A	0.4491	0.4763	0.5708	0.027*
C14	-0.3446 (2)	-0.2050 (2)	0.25940 (15)	0.0234 (3)
C15	-0.3754 (2)	-0.3076 (2)	0.08290 (16)	0.0292 (4)
H15A	-0.4990	-0.3501	0.0636	0.035*
H15B	-0.3399	-0.3859	0.1032	0.035*
C16	-0.3270 (3)	-0.2504 (3)	-0.01223 (19)	0.0441 (6)
H16A	-0.3815	-0.3318	-0.0754	0.066*
H16B	-0.2044	-0.2086	0.0080	0.066*
H16C	-0.3632	-0.1734	-0.0316	0.066*
C17	-0.0017 (2)	0.3308 (2)	0.70099 (13)	0.0214 (3)
H17A	0.1063	0.4151	0.7467	0.026*
H17B	-0.0149	0.2442	0.7376	0.026*
C18	-0.1456 (2)	0.3693 (2)	0.69594 (14)	0.0236 (3)
H18A	-0.2548	0.2819	0.6574	0.028*
H18B	-0.1392	0.4503	0.6540	0.028*
C19	-0.1347 (2)	0.4175 (2)	0.81236 (15)	0.0261 (4)
H19A	-0.2453	0.4099	0.8095	0.031*
H19B	-0.1099	0.3490	0.8594	0.031*
C20	-0.0400 (3)	0.6949 (2)	0.84211 (18)	0.0354 (5)
H20A	-0.1478	0.6790	0.8505	0.042*
H20B	-0.0439	0.7153	0.7676	0.042*
C21	0.1094 (4)	0.8213 (3)	0.9307 (2)	0.0443 (6)

H21A	0.1467	0.9144	0.9017	0.053*
H21B	0.0793	0.8379	0.9952	0.053*
C22	0.2468 (3)	0.7692 (2)	0.96175 (18)	0.0353 (5)
H22A	0.3248	0.8102	0.9210	0.042*
H22B	0.3121	0.7996	1.0416	0.042*
C23	0.1509 (2)	0.6021 (2)	0.92917 (15)	0.0268 (4)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0245 (6)	0.0248 (7)	0.0246 (6)	0.0055 (5)	0.0078 (5)	−0.0053 (5)
O2	0.0311 (7)	0.0247 (7)	0.0282 (7)	0.0022 (6)	0.0062 (6)	0.0020 (6)
O3	0.0361 (8)	0.0421 (9)	0.0348 (8)	0.0228 (7)	0.0031 (6)	−0.0081 (7)
O4	0.0244 (7)	0.0338 (8)	0.0417 (9)	0.0048 (6)	0.0088 (6)	0.0019 (7)
O5	0.0404 (8)	0.0261 (8)	0.0417 (9)	0.0100 (7)	0.0116 (7)	−0.0034 (6)
N1	0.0184 (6)	0.0202 (7)	0.0214 (7)	0.0080 (5)	0.0073 (5)	0.0017 (5)
N2	0.0188 (6)	0.0213 (7)	0.0165 (6)	0.0072 (5)	0.0070 (5)	0.0028 (5)
N3	0.0271 (7)	0.0251 (8)	0.0226 (7)	0.0122 (6)	0.0097 (6)	0.0024 (6)
N4	0.0276 (7)	0.0226 (8)	0.0231 (7)	0.0048 (6)	0.0089 (6)	0.0020 (6)
C1	0.0173 (7)	0.0202 (8)	0.0204 (7)	0.0086 (6)	0.0079 (6)	0.0034 (6)
C2	0.0194 (7)	0.0206 (8)	0.0200 (7)	0.0089 (6)	0.0068 (6)	0.0019 (6)
C3	0.0193 (7)	0.0205 (8)	0.0204 (7)	0.0079 (6)	0.0053 (6)	0.0028 (6)
C4	0.0199 (7)	0.0222 (8)	0.0240 (8)	0.0058 (6)	0.0078 (6)	0.0064 (6)
C5	0.0221 (7)	0.0228 (8)	0.0205 (7)	0.0069 (7)	0.0092 (6)	0.0048 (6)
C6	0.0188 (7)	0.0199 (8)	0.0178 (7)	0.0078 (6)	0.0055 (6)	0.0025 (6)
C7	0.0181 (7)	0.0177 (7)	0.0194 (7)	0.0078 (6)	0.0064 (6)	0.0028 (6)
C8	0.0195 (7)	0.0204 (8)	0.0174 (7)	0.0086 (6)	0.0063 (6)	0.0033 (6)
C9	0.0202 (7)	0.0230 (8)	0.0232 (8)	0.0105 (7)	0.0080 (6)	0.0037 (6)
C10	0.0264 (8)	0.0190 (8)	0.0207 (8)	0.0103 (7)	0.0080 (6)	0.0027 (6)
C11	0.0222 (7)	0.0194 (8)	0.0184 (7)	0.0051 (6)	0.0066 (6)	0.0028 (6)
C12	0.0200 (7)	0.0245 (9)	0.0241 (8)	0.0067 (7)	0.0093 (6)	0.0031 (7)
C13	0.0226 (8)	0.0240 (8)	0.0215 (8)	0.0104 (7)	0.0101 (6)	0.0031 (6)
C14	0.0208 (7)	0.0236 (8)	0.0225 (8)	0.0098 (7)	0.0036 (6)	0.0029 (6)
C15	0.0267 (9)	0.0276 (10)	0.0255 (9)	0.0091 (8)	0.0046 (7)	−0.0052 (7)
C16	0.0467 (13)	0.0415 (13)	0.0320 (11)	0.0082 (11)	0.0157 (10)	−0.0058 (10)
C17	0.0224 (7)	0.0263 (9)	0.0149 (7)	0.0103 (7)	0.0067 (6)	0.0033 (6)
C18	0.0196 (7)	0.0285 (9)	0.0202 (8)	0.0084 (7)	0.0072 (6)	−0.0003 (7)
C19	0.0234 (8)	0.0301 (9)	0.0238 (8)	0.0088 (7)	0.0119 (7)	0.0000 (7)
C20	0.0498 (12)	0.0293 (10)	0.0289 (10)	0.0226 (10)	0.0095 (9)	0.0032 (8)
C21	0.0650 (16)	0.0292 (11)	0.0329 (11)	0.0212 (11)	0.0094 (11)	0.0030 (9)
C22	0.0365 (10)	0.0305 (10)	0.0285 (10)	0.0038 (9)	0.0143 (8)	−0.0019 (8)
C23	0.0284 (9)	0.0302 (10)	0.0216 (8)	0.0116 (8)	0.0114 (7)	−0.0017 (7)

Geometric parameters (Å, °)

O1—C14	1.337 (2)	C10—C11	1.386 (2)
O1—C15	1.461 (2)	C10—H10A	0.9500
O2—C14	1.208 (2)	C11—C12	1.387 (3)

O3—C23	1.229 (3)	C12—C13	1.386 (3)
O4—N4	1.231 (2)	C12—H12A	0.9500
O5—N4	1.227 (2)	C13—H13A	0.9500
N1—C7	1.325 (2)	C15—C16	1.496 (3)
N1—C1	1.392 (2)	C15—H15A	0.9900
N2—C6	1.380 (2)	C15—H15B	0.9900
N2—C7	1.386 (2)	C16—H16A	0.9800
N2—C17	1.468 (2)	C16—H16B	0.9800
N3—C23	1.347 (2)	C16—H16C	0.9800
N3—C19	1.448 (2)	C17—C18	1.527 (2)
N3—C20	1.456 (3)	C17—H17A	0.9900
N4—C11	1.470 (2)	C17—H17B	0.9900
C1—C2	1.396 (2)	C18—C19	1.534 (2)
C1—C6	1.405 (2)	C18—H18A	0.9900
C2—C3	1.386 (2)	C18—H18B	0.9900
C2—H2A	0.9500	C19—H19A	0.9900
C3—C4	1.413 (2)	C19—H19B	0.9900
C3—C14	1.492 (2)	C20—C21	1.531 (3)
C4—C5	1.377 (3)	C20—H20A	0.9900
C4—H4A	0.9500	C20—H20B	0.9900
C5—C6	1.396 (2)	C21—C22	1.527 (4)
C5—H5A	0.9500	C21—H21A	0.9900
C7—C8	1.466 (2)	C21—H21B	0.9900
C8—C13	1.400 (2)	C22—C23	1.515 (3)
C8—C9	1.402 (2)	C22—H22A	0.9900
C9—C10	1.389 (3)	C22—H22B	0.9900
C9—H9A	0.9500		
C14—O1—C15	115.53 (15)	O1—C14—C3	112.44 (15)
C7—N1—C1	104.74 (14)	O1—C15—C16	107.13 (17)
C6—N2—C7	106.18 (13)	O1—C15—H15A	110.3
C6—N2—C17	124.31 (14)	C16—C15—H15A	110.3
C7—N2—C17	128.43 (14)	O1—C15—H15B	110.3
C23—N3—C19	123.98 (17)	C16—C15—H15B	110.3
C23—N3—C20	114.20 (17)	H15A—C15—H15B	108.5
C19—N3—C20	121.81 (16)	C15—C16—H16A	109.5
O5—N4—O4	123.33 (17)	C15—C16—H16B	109.5
O5—N4—C11	118.19 (16)	H16A—C16—H16B	109.5
O4—N4—C11	118.48 (16)	C15—C16—H16C	109.5
N1—C1—C2	129.65 (15)	H16A—C16—H16C	109.5
N1—C1—C6	110.09 (15)	H16B—C16—H16C	109.5
C2—C1—C6	120.25 (16)	N2—C17—C18	113.73 (14)
C3—C2—C1	117.63 (16)	N2—C17—H17A	108.8
C3—C2—H2A	121.2	C18—C17—H17A	108.8
C1—C2—H2A	121.2	N2—C17—H17B	108.8
C2—C3—C4	121.30 (16)	C18—C17—H17B	108.8
C2—C3—C14	121.58 (16)	H17A—C17—H17B	107.7
C4—C3—C14	117.12 (16)	C17—C18—C19	110.22 (14)

C5—C4—C3	121.73 (16)	C17—C18—H18A	109.6
C5—C4—H4A	119.1	C19—C18—H18A	109.6
C3—C4—H4A	119.1	C17—C18—H18B	109.6
C4—C5—C6	116.66 (16)	C19—C18—H18B	109.6
C4—C5—H5A	121.7	H18A—C18—H18B	108.1
C6—C5—H5A	121.7	N3—C19—C18	112.88 (15)
N2—C6—C5	131.67 (16)	N3—C19—H19A	109.0
N2—C6—C1	105.92 (14)	C18—C19—H19A	109.0
C5—C6—C1	122.42 (16)	N3—C19—H19B	109.0
N1—C7—N2	113.07 (15)	C18—C19—H19B	109.0
N1—C7—C8	122.98 (15)	H19A—C19—H19B	107.8
N2—C7—C8	123.96 (15)	N3—C20—C21	103.43 (18)
C13—C8—C9	119.31 (16)	N3—C20—H20A	111.1
C13—C8—C7	118.47 (15)	C21—C20—H20A	111.1
C9—C8—C7	122.19 (15)	N3—C20—H20B	111.1
C10—C9—C8	120.27 (15)	C21—C20—H20B	111.1
C10—C9—H9A	119.9	H20A—C20—H20B	109.0
C8—C9—H9A	119.9	C22—C21—C20	104.44 (18)
C11—C10—C9	118.67 (16)	C22—C21—H21A	110.9
C11—C10—H10A	120.7	C20—C21—H21A	110.9
C9—C10—H10A	120.7	C22—C21—H21B	110.9
C10—C11—C12	122.60 (16)	C20—C21—H21B	110.9
C10—C11—N4	118.71 (16)	H21A—C21—H21B	108.9
C12—C11—N4	118.68 (16)	C23—C22—C21	104.12 (18)
C13—C12—C11	118.05 (16)	C23—C22—H22A	110.9
C13—C12—H12A	121.0	C21—C22—H22A	110.9
C11—C12—H12A	121.0	C23—C22—H22B	110.9
C12—C13—C8	121.02 (16)	C21—C22—H22B	110.9
C12—C13—H13A	119.5	H22A—C22—H22B	109.0
C8—C13—H13A	119.5	O3—C23—N3	124.88 (19)
O2—C14—O1	123.64 (17)	O3—C23—C22	127.05 (19)
O2—C14—C3	123.92 (17)	N3—C23—C22	108.06 (18)
C7—N1—C1—C2	179.48 (17)	C9—C10—C11—N4	-176.78 (16)
C7—N1—C1—C6	0.23 (19)	O5—N4—C11—C10	-13.9 (3)
N1—C1—C2—C3	-179.08 (17)	O4—N4—C11—C10	166.00 (17)
C6—C1—C2—C3	0.1 (2)	O5—N4—C11—C12	166.58 (18)
C1—C2—C3—C4	-0.9 (3)	O4—N4—C11—C12	-13.5 (3)
C1—C2—C3—C14	179.30 (15)	C10—C11—C12—C13	-2.7 (3)
C2—C3—C4—C5	1.2 (3)	N4—C11—C12—C13	176.81 (16)
C14—C3—C4—C5	-179.04 (16)	C11—C12—C13—C8	0.4 (3)
C3—C4—C5—C6	-0.5 (3)	C9—C8—C13—C12	1.7 (3)
C7—N2—C6—C5	-179.83 (18)	C7—C8—C13—C12	179.80 (16)
C17—N2—C6—C5	11.2 (3)	C15—O1—C14—O2	-0.5 (3)
C7—N2—C6—C1	0.53 (18)	C15—O1—C14—C3	179.81 (15)
C17—N2—C6—C1	-168.39 (15)	C2—C3—C14—O2	167.84 (18)
C4—C5—C6—N2	-179.88 (17)	C4—C3—C14—O2	-12.0 (3)
C4—C5—C6—C1	-0.3 (3)	C2—C3—C14—O1	-12.4 (2)

N1—C1—C6—N2	-0.49 (19)	C4—C3—C14—O1	167.77 (15)
C2—C1—C6—N2	-179.81 (15)	C14—O1—C15—C16	167.07 (17)
N1—C1—C6—C5	179.83 (16)	C6—N2—C17—C18	-77.5 (2)
C2—C1—C6—C5	0.5 (3)	C7—N2—C17—C18	116.10 (19)
C1—N1—C7—N2	0.12 (19)	N2—C17—C18—C19	-174.58 (15)
C1—N1—C7—C8	-179.84 (15)	C23—N3—C19—C18	-95.5 (2)
C6—N2—C7—N1	-0.42 (19)	C20—N3—C19—C18	85.1 (2)
C17—N2—C7—N1	167.89 (16)	C17—C18—C19—N3	77.3 (2)
C6—N2—C7—C8	179.53 (15)	C23—N3—C20—C21	-13.5 (2)
C17—N2—C7—C8	-12.2 (3)	C19—N3—C20—C21	165.93 (18)
N1—C7—C8—C13	-38.3 (2)	N3—C20—C21—C22	22.3 (2)
N2—C7—C8—C13	141.76 (17)	C20—C21—C22—C23	-23.2 (2)
N1—C7—C8—C9	139.76 (18)	C19—N3—C23—O3	0.2 (3)
N2—C7—C8—C9	-40.2 (2)	C20—N3—C23—O3	179.69 (19)
C13—C8—C9—C10	-1.7 (3)	C19—N3—C23—C22	179.15 (16)
C7—C8—C9—C10	-179.69 (16)	C20—N3—C23—C22	-1.4 (2)
C8—C9—C10—C11	-0.5 (3)	C21—C22—C23—O3	-165.3 (2)
C9—C10—C11—C12	2.7 (3)	C21—C22—C23—N3	15.8 (2)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C15—H15A \cdots O3 ⁱ	0.99	2.41	3.355 (3)	159
C15—H15B \cdots O3 ⁱⁱ	0.99	2.38	3.186 (3)	139
C19—H19A \cdots O2 ⁱⁱⁱ	0.99	2.38	3.312 (3)	156

Symmetry codes: (i) $x-1, y-1, z-1$; (ii) $-x, -y, -z+1$; (iii) $-x-1, -y, -z+1$.