

# Bis{2-[{(pyridin-4-yl- $\kappa$ N)sulfanyl]-pyrazine}silver(I) tetrafluoridoborate

Zi-Jia Wang

Department of Chemistry, Capital Normal University, Beijing 100048, People's Republic of China  
 Correspondence e-mail: wanchqing@yahoo.com.cn

Received 22 November 2011; accepted 29 November 2011

Key indicators: single-crystal X-ray study;  $T = 296\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.006\text{ \AA}$ ;  $R$  factor = 0.036;  $wR$  factor = 0.116; data-to-parameter ratio = 12.4.

In the title mononuclear complex,  $[\text{Ag}(\text{C}_9\text{H}_7\text{N}_3\text{S})_2]\text{BF}_4$ , the  $\text{Ag}^{\text{I}}$  ion adopts a virtually linear coordination geometry [ $\text{N}-\text{Ag}-\text{N} = 178.06(11)^{\circ}$ ] with the two ligands bound to the metal atom *via* the pyridine N atoms. The metal-coordinated pyridine rings are almost coplanar, making a dihedral angle of  $1.5(2)^{\circ}$ , while the two pendent pyrazine rings are arranged on the same side of the  $\text{N}-\text{Ag}-\text{N}$  line. Along the  $a$  axis, the mononuclear coordination units are stacked with  $\pi-\pi$  interactions between the pyridine rings [centroid-centroid distance =  $3.569(4)\text{ \AA}$ ], leading to infinite chains. The chains are interconnected through intermolecular  $\text{N}(\text{pyrazine})\cdots\pi(\text{pyrazine})$  interactions forming layers parallel to the  $ab$  plane [ $\text{N}\cdots\text{centroid} = 3.268(5)\text{ \AA}$ ]. These layers are further stacked along the  $c$ -axis direction, furnishing a three-dimensional supramolecular framework with the tetrafluoridoborate anions embedded within the interstices.

## Related literature

For metal complexes with chalcogenobispyridines and derivatives, see: Baradello *et al.* (2004); Dunne *et al.* (1997). For the crystal structures of di-2-pyridyl sulfide and its *N*-positional isomer complexes, see: Jung *et al.* (2001, 2003). For the  $\text{N}(\text{pyrazinyl})\cdots\text{centroid}(\text{pyrazinyl})$  distance in  $\{[\text{Ni}(\text{L})-(\text{NO}_3)_2]\}_{\infty}$  ( $\text{L}$  = bis(2-pyrazylmethyl)sulfide), see: Black *et al.* (2007); For van der Waals radii, see: Bondi (1964) and for the half thickness of phenyl rings, see: Malone *et al.* (1997).

## Experimental

### Crystal data

$[\text{Ag}(\text{C}_9\text{H}_7\text{N}_3\text{S})_2]\text{BF}_4$   
 $M_r = 573.15$   
 Monoclinic,  $P2_1/n$   
 $a = 7.2232(2)\text{ \AA}$   
 $b = 16.4826(3)\text{ \AA}$   
 $c = 17.6098(4)\text{ \AA}$   
 $\beta = 91.666(1)^{\circ}$

$V = 2095.69(8)\text{ \AA}^3$   
 $Z = 4$   
 Mo  $K\alpha$  radiation  
 $\mu = 1.22\text{ mm}^{-1}$   
 $T = 296\text{ K}$   
 $0.40 \times 0.30 \times 0.20\text{ mm}$

### Data collection

Bruker SMART APEXII CCD  
 area-detector' diffractometer  
 Absorption correction: multi-scan  
 (*SADABS*; Bruker, 2007)  
 $T_{\min} = 0.616$ ,  $T_{\max} = 0.746$

14794 measured reflections  
 3597 independent reflections  
 3215 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.028$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.036$   
 $wR(F^2) = 0.116$   
 $S = 1.11$   
 3597 reflections

289 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.40\text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.62\text{ e \AA}^{-3}$


Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL* and *PLATON* (Spek, 2009).

The authors are grateful for financial support from the Beijing Municipal Education Commission.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: ZQ2141).

## References

- Baradello, L., Lo Schiavo, S., Nicolò, F., Alibrandi, G., Tresoldi, G. & Lanza, S. (2004). *Eur. J. Inorg. Chem.* pp. 3358–3369.
- Black, C. A., Hanton, L. R. & Spicer, M. D. (2007). *Inorg. Chem.* **46**, 3669–3679.
- Bondi, A. (1964). *J. Phys. Chem.* **68**, 441–451.
- Bruker (2007). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Dunne, S. J., Summers, L. A. & von Nagy-Felsobuki, E. I. (1997). *Coord. Chem. Rev.* **165**, 1–92.
- Jung, O.-S., Kim, Y. J., Lee, Y.-A., Chae, H. K., Jang, H. G. & Hong, J. (2001). *Inorg. Chem.* **40**, 2105–2110.
- Jung, O.-S., Kim, Y. J., Lee, Y.-A., Park, K.-M. & Lee, S. S. (2003). *Inorg. Chem.* **42**, 844–850.
- Malone, J. F., Murray, C. M., Charlton, M. H., Docherty, R. & Lavery, A. J. (1997). *J. Chem. Soc. Faraday Trans.* **93**, 3429–3436.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.



# supporting information

*Acta Cryst.* (2012). E68, m4 [doi:10.1107/S1600536811051270]

## Bis{2-[(pyridin-4-yl- $\kappa$ N)sulfanyl]pyrazine}silver(I) tetrafluoridoborate

Zi-Jia Wang

### S1. Comment

Chalcogenobispyridines and derivates were widely used as versatile building blocks for supramolecular assembly (Baradello *et al.*, 2004; Dunne *et al.*, 1997). The ligand, such as the di-2-pyridyl sulfide and its N-positional isomers, endowed with the rotatable C( $sp^2$ )—S bond and a variable C( $sp^2$ )—S—C( $sp^2$ ) angle (about 100°), exhibits flexible ligation modes in construction of diverse coordination motifs with unusual properties (Jung *et al.*, 2001; Jung *et al.*, 2003). Herein, we report a new silver complex derived from the 2-(pyridin-4-ylsulfanyl)pyrazine ligand.


In the mononuclear complex,  $[\text{Ag}(\text{C}_9\text{H}_7\text{N}_3\text{S})_2]^+\text{BF}_4^-$ , the silver(I) ion adopts a linear coordination geometry [N3—Ag1—N4 = 178.06 (1)°] with the two ligands bound to the metal center via the 4-pyridyl N atoms (Fig. 1). The two pyridyl rings bound to Ag<sup>I</sup> are almost coplanar, while the two pendent pyrazinyl rings are arranged on the same side of the N—Ag—N line. The dihedral angle between the mean planes of the two pendent pyrazinyl rings is 48.89 (1)°. Along the *a* axis, the mononuclear units are stacked with  $\pi$ — $\pi$  interactions between the 4-pyridyl rings [Cg1···Cg2<sup>i</sup> 3.569 (4) Å; symmetry code: (i) = *x*+1, *y*, *z*] leading to infinite chains (Cg1 = C5-C6-C7-N3-C8-C9; Cg2 = N4-C10-C11-C12-C13-C14). The formed chains interconnect through intermolecular N(pyrazinyl)··· $\pi$ (pyrazinyl) interactions forming layers parallel to the *ab* plane (Fig. 2). For the N(pyrazinyl)··· $\pi$ (pyrazinyl) contact, the N6···Cg3<sup>ii</sup> distance equals 3.268 (5) Å (Cg3 = N1-C1-C2-N2-C3-C4; (ii) = -*x*, -*y*, -*z*) which is comparable to that N(pyrazinyl)···centroid(pyrazinyl) of 3.05 Å reported by Black *et al.* (2007) in  $\{[\text{Ni}(\text{L})(\text{NO}_3)_2]\}_\infty$  (*L* = bis(2-pyrazylmethyl)sulfide). These distances are shorter than the van der Waals separation of 3.40 Å on the basis of Pauling's values for the half thickness of phenyl rings (1.85 Å) (Malone *et al.*, 1997) and the van der Waals radius of N (1.55 Å) (Bondi, 1964). The almost parallel layers are further stacked along the *c* direction to furnish a three-dimensional supramolecular framework with the tetrafluoridoborate anions embedded within the interstices (Fig. 3).

### S2. Experimental


4-Pyridyl-2-pyrazinyl sulfide was synthesized by reacting 2-chloropyrazine (0.6 g, 5.2 mmol) with sodium pyridine-4-thiolate (5 mmol) in 40 ml methanol. The mixture was refluxed with stirring for 10 hours under the protection of N<sub>2</sub>. After filtration and concentration in vacuo, the obtained crude product was further purified by chromatography on silica gel using ether acetate/dichloromethane (1:3) as the eluent, giving 0.444 g of yellow powder of 4-pyridyl-2-pyrazinyl sulfide in 47% yield. Reaction of 4-pyridyl-2-pyrazinyl sulfide (19 mg, 0.1 mmol) and AgBF<sub>4</sub> (20 mg, 0.1 mmol) in 4 ml methanol with stirring at room temperature for 3 hours. The obtained clear solution was filtrated, and the filtration was left evaporation in air. After about one week, the block-like crystals of the title complex were deposited (18.1 mg, yield 63%).

**S3. Refinement**


The H atoms were placed in idealized positions and allowed to ride on the relevant carbon atoms, with C—H = 0.93 Å and  $U_{iso}(\text{H}) = 1.2U_{eq}(\text{C})$ .

**Figure 1**

The atom-numbering scheme of the title complex. Displacement ellipsoids are drawn at the 50% probability level, while the H atoms are shown as rods of arbitrary radius.

**Figure 2**

The  $\pi$ - $\pi$  stacking and N... $\pi$ (pyrazinyl) interactions between the mononuclear units, which are respectively shown as thick red-dashed lines and thin blue-dashed lines.

**Figure 3**

The packing structure of the title complex. The  $\text{Ag}^{\text{I}}$  ions are shown as purple balls, while the B-F bonds are shown as thick bond mode for clarity.

### Bis{2-[{(pyridin-4-yl- $\kappa\text{N}$ )sulfanyl]pyrazine}silver(I) tetrafluoridoborate

#### Crystal data

$[\text{Ag}(\text{C}_9\text{H}_7\text{N}_3\text{S})_2]\text{BF}_4$

$M_r = 573.15$

Monoclinic,  $P2_1/n$

Hall symbol: -P 2yn

$a = 7.2232 (2)$  Å

$b = 16.4826 (3)$  Å

$c = 17.6098 (4)$  Å

$\beta = 91.666 (1)^\circ$

$V = 2095.69 (8)$  Å<sup>3</sup>

$Z = 4$

$F(000) = 1136$

$D_x = 1.817 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

$\mu = 1.22 \text{ mm}^{-1}$

$T = 296 \text{ K}$

Block, yellow

$0.40 \times 0.30 \times 0.20$  mm

#### Data collection

'Bruker SMART APEXII CCD area-detector'  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

$\omega$  scans

Absorption correction: multi-scan  
(*SADABS*; Bruker, 2007)

$T_{\min} = 0.616$ ,  $T_{\max} = 0.746$

14794 measured reflections

3597 independent reflections

3215 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.028$

$\theta_{\max} = 25.0^\circ$ ,  $\theta_{\min} = 2.3^\circ$

$h = -8 \rightarrow 8$

$k = -19 \rightarrow 19$

$l = -20 \rightarrow 20$

#### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.036$

$wR(F^2) = 0.116$

$S = 1.11$

3597 reflections

289 parameters

0 restraints

Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map

Hydrogen site location: inferred from  
neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0584P)^2 + 2.8465P] \quad P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.001 \quad \Delta\rho_{\max} = 0.40 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.62 \text{ e } \text{\AA}^{-3}$$

*Special details*

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted R-factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional R-factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\text{sigma}(F^2)$  is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and R-factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Ag1	0.49553 (4)	0.746244 (18)	0.040102 (19)	0.04882 (15)
S1	1.31192 (14)	0.55322 (8)	0.04143 (6)	0.0556 (3)
S2	-0.31897 (14)	0.94030 (8)	0.04687 (6)	0.0554 (3)
N1	1.3844 (4)	0.62555 (19)	0.1758 (2)	0.0456 (8)
N2	1.5853 (5)	0.4934 (2)	0.2314 (2)	0.0557 (9)
N3	0.7552 (4)	0.68164 (18)	0.04124 (17)	0.0392 (7)
N4	0.2344 (4)	0.81040 (18)	0.04305 (18)	0.0409 (7)
N5	-0.3990 (4)	0.85798 (19)	0.1733 (2)	0.0459 (8)
N6	-0.5935 (5)	0.9876 (2)	0.2337 (2)	0.0582 (10)
C1	1.4578 (6)	0.6243 (3)	0.2450 (3)	0.0532 (10)
H1A	1.4393	0.6685	0.2767	0.064*
C2	1.5609 (6)	0.5599 (3)	0.2723 (3)	0.0576 (11)
H2A	1.6150	0.5630	0.3208	0.069*
C3	1.5056 (6)	0.4919 (2)	0.1643 (2)	0.0482 (9)
H3A	1.5164	0.4456	0.1346	0.058*
C4	1.4040 (5)	0.5577 (2)	0.1353 (2)	0.0387 (8)
C5	1.0957 (5)	0.6030 (2)	0.0462 (2)	0.0378 (8)
C6	1.0216 (6)	0.6326 (3)	-0.0213 (2)	0.0499 (10)
H6A	1.0862	0.6276	-0.0660	0.060*
C7	0.8514 (6)	0.6697 (3)	-0.0217 (2)	0.0487 (10)
H7A	0.8005	0.6874	-0.0679	0.058*
C8	0.8306 (5)	0.6540 (3)	0.1058 (2)	0.0458 (9)
H8A	0.7667	0.6626	0.1502	0.055*
C9	0.9972 (5)	0.6135 (3)	0.1114 (2)	0.0479 (10)
H9A	1.0421	0.5938	0.1579	0.057*
C10	0.1515 (6)	0.8409 (3)	-0.0177 (3)	0.0569 (11)
H10A	0.2110	0.8369	-0.0637	0.068*
C11	-0.0205 (6)	0.8788 (3)	-0.0173 (2)	0.0546 (11)
H11A	-0.0779	0.8967	-0.0622	0.066*
C12	-0.1045 (5)	0.8893 (2)	0.0520 (2)	0.0398 (8)
C13	-0.0160 (5)	0.8602 (3)	0.1160 (2)	0.0472 (9)
H13A	-0.0676	0.8671	0.1634	0.057*

C14	0.1513 (5)	0.8204 (3)	0.1094 (2)	0.0462 (9)
H14A	0.2089	0.7997	0.1532	0.055*
C15	-0.4147 (5)	0.9284 (2)	0.1376 (2)	0.0373 (8)
C16	-0.4753 (6)	0.8546 (3)	0.2415 (3)	0.0560 (11)
H16A	-0.4609	0.8077	0.2703	0.067*
C17	-0.5735 (6)	0.9177 (3)	0.2701 (3)	0.0580 (11)
H17A	-0.6283	0.9115	0.3169	0.070*
C18	-0.5097 (6)	0.9934 (2)	0.1678 (2)	0.0457 (9)
H18A	-0.5148	1.0422	0.1412	0.055*
B1	0.4920 (7)	0.7584 (3)	-0.1658 (3)	0.0479 (11)
F1	0.5771 (6)	0.8120 (2)	-0.1155 (2)	0.0970 (11)
F2	0.4070 (6)	0.7015 (2)	-0.1210 (2)	0.0977 (11)
F3	0.3676 (6)	0.7987 (3)	-0.2082 (3)	0.1304 (16)
F4	0.6212 (6)	0.7205 (3)	-0.2054 (3)	0.1233 (15)

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Ag1	0.0314 (2)	0.0524 (2)	0.0627 (2)	0.01446 (12)	0.00139 (14)	-0.00343 (13)
S1	0.0364 (5)	0.0807 (8)	0.0494 (6)	0.0263 (5)	-0.0019 (4)	-0.0044 (5)
S2	0.0389 (5)	0.0792 (8)	0.0483 (6)	0.0271 (5)	0.0076 (4)	0.0090 (5)
N1	0.0355 (17)	0.0401 (17)	0.061 (2)	0.0034 (13)	0.0009 (15)	0.0044 (15)
N2	0.053 (2)	0.057 (2)	0.056 (2)	0.0147 (17)	-0.0056 (17)	0.0105 (17)
N3	0.0299 (15)	0.0427 (17)	0.0450 (17)	0.0059 (12)	-0.0009 (13)	-0.0034 (13)
N4	0.0309 (15)	0.0422 (17)	0.0496 (18)	0.0059 (12)	0.0039 (13)	0.0010 (13)
N5	0.0373 (17)	0.0398 (17)	0.061 (2)	0.0029 (13)	0.0068 (15)	-0.0034 (15)
N6	0.057 (2)	0.061 (2)	0.057 (2)	0.0122 (17)	0.0154 (18)	-0.0127 (18)
C1	0.045 (2)	0.050 (2)	0.064 (3)	0.0002 (18)	0.001 (2)	-0.0068 (19)
C2	0.048 (2)	0.071 (3)	0.054 (3)	0.004 (2)	-0.0018 (19)	0.004 (2)
C3	0.046 (2)	0.042 (2)	0.057 (2)	0.0077 (17)	0.0040 (18)	0.0045 (18)
C4	0.0239 (16)	0.042 (2)	0.051 (2)	0.0038 (14)	0.0030 (15)	0.0082 (16)
C5	0.0277 (17)	0.0403 (19)	0.045 (2)	0.0046 (14)	-0.0009 (14)	-0.0010 (15)
C6	0.042 (2)	0.068 (3)	0.040 (2)	0.0146 (19)	0.0011 (16)	-0.0026 (18)
C7	0.040 (2)	0.065 (3)	0.041 (2)	0.0171 (18)	-0.0053 (16)	0.0010 (18)
C8	0.0258 (17)	0.067 (3)	0.045 (2)	0.0037 (17)	0.0081 (15)	0.0042 (18)
C9	0.0332 (19)	0.067 (3)	0.044 (2)	0.0065 (17)	0.0023 (16)	0.0182 (18)
C10	0.052 (2)	0.066 (3)	0.054 (2)	0.023 (2)	0.023 (2)	0.010 (2)
C11	0.051 (2)	0.072 (3)	0.041 (2)	0.026 (2)	0.0096 (18)	0.015 (2)
C12	0.0314 (18)	0.0379 (19)	0.050 (2)	0.0032 (14)	0.0055 (15)	-0.0012 (15)
C13	0.0333 (19)	0.067 (3)	0.041 (2)	0.0074 (18)	0.0025 (15)	-0.0054 (18)
C14	0.0289 (18)	0.064 (2)	0.046 (2)	0.0058 (17)	-0.0004 (15)	-0.0009 (18)
C15	0.0248 (16)	0.0421 (19)	0.0450 (19)	0.0033 (14)	0.0010 (14)	-0.0050 (15)
C16	0.048 (2)	0.053 (3)	0.067 (3)	-0.0004 (19)	0.008 (2)	0.014 (2)
C17	0.052 (3)	0.075 (3)	0.048 (2)	0.009 (2)	0.0100 (19)	0.002 (2)
C18	0.042 (2)	0.042 (2)	0.054 (2)	0.0072 (16)	0.0028 (17)	-0.0039 (17)
B1	0.055 (3)	0.053 (3)	0.036 (2)	0.010 (2)	0.002 (2)	-0.0037 (18)
F1	0.132 (3)	0.073 (2)	0.085 (2)	-0.0111 (19)	-0.030 (2)	-0.0048 (16)
F2	0.129 (3)	0.073 (2)	0.093 (2)	-0.013 (2)	0.034 (2)	0.0027 (17)

F3	0.127 (4)	0.134 (4)	0.127 (4)	0.035 (3)	-0.057 (3)	0.029 (3)
F4	0.124 (3)	0.130 (3)	0.119 (3)	0.032 (3)	0.061 (3)	-0.021 (3)

*Geometric parameters ( $\text{\AA}$ ,  $^{\circ}$ )*

Ag1—N3	2.156 (3)	C5—C9	1.380 (5)
Ag1—N4	2.165 (3)	C6—C7	1.373 (6)
S1—C4	1.765 (4)	C6—H6A	0.9300
S1—C5	1.768 (3)	C7—H7A	0.9300
S2—C12	1.763 (4)	C8—C9	1.378 (5)
S2—C15	1.770 (4)	C8—H8A	0.9300
N1—C1	1.315 (6)	C9—H9A	0.9300
N1—C4	1.337 (5)	C10—C11	1.391 (6)
N2—C3	1.300 (6)	C10—H10A	0.9300
N2—C2	1.326 (6)	C11—C12	1.389 (5)
N3—C8	1.327 (5)	C11—H11A	0.9300
N3—C7	1.340 (5)	C12—C13	1.366 (6)
N4—C10	1.311 (5)	C13—C14	1.383 (5)
N4—C14	1.339 (5)	C13—H13A	0.9300
N5—C15	1.324 (5)	C14—H14A	0.9300
N5—C16	1.336 (6)	C15—C18	1.386 (5)
N6—C17	1.324 (6)	C16—C17	1.364 (6)
N6—C18	1.328 (5)	C16—H16A	0.9300
C1—C2	1.376 (6)	C17—H17A	0.9300
C1—H1A	0.9300	C18—H18A	0.9300
C2—H2A	0.9300	B1—F3	1.330 (6)
C3—C4	1.397 (5)	B1—F4	1.337 (6)
C3—H3A	0.9300	B1—F2	1.380 (6)
C5—C6	1.378 (5)	B1—F1	1.382 (6)
N3—Ag1—N4	178.06 (11)	C8—C9—C5	118.1 (4)
C4—S1—C5	104.22 (17)	C8—C9—H9A	120.9
C12—S2—C15	105.47 (18)	C5—C9—H9A	120.9
C1—N1—C4	115.8 (3)	N4—C10—C11	123.6 (4)
C3—N2—C2	116.6 (4)	N4—C10—H10A	118.2
C8—N3—C7	116.7 (3)	C11—C10—H10A	118.2
C8—N3—Ag1	120.8 (2)	C12—C11—C10	118.3 (4)
C7—N3—Ag1	122.5 (3)	C12—C11—H11A	120.9
C10—N4—C14	117.4 (3)	C10—C11—H11A	120.9
C10—N4—Ag1	123.0 (3)	C13—C12—C11	118.4 (3)
C14—N4—Ag1	119.6 (3)	C13—C12—S2	126.8 (3)
C15—N5—C16	115.5 (3)	C11—C12—S2	114.8 (3)
C17—N6—C18	116.1 (4)	C12—C13—C14	119.1 (4)
N1—C1—C2	122.4 (4)	C12—C13—H13A	120.5
N1—C1—H1A	118.8	C14—C13—H13A	120.5
C2—C1—H1A	118.8	N4—C14—C13	123.1 (4)
N2—C2—C1	121.8 (4)	N4—C14—H14A	118.4
N2—C2—H2A	119.1	C13—C14—H14A	118.4

C1—C2—H2A	119.1	N5—C15—C18	122.2 (3)
N2—C3—C4	122.1 (4)	N5—C15—S2	119.7 (3)
N2—C3—H3A	118.9	C18—C15—S2	118.1 (3)
C4—C3—H3A	118.9	N5—C16—C17	122.2 (4)
N1—C4—C3	121.1 (4)	N5—C16—H16A	118.9
N1—C4—S1	119.5 (3)	C17—C16—H16A	118.9
C3—C4—S1	119.3 (3)	N6—C17—C16	122.3 (4)
C6—C5—C9	118.4 (3)	N6—C17—H17A	118.8
C6—C5—S1	116.4 (3)	C16—C17—H17A	118.8
C9—C5—S1	125.1 (3)	N6—C18—C15	121.5 (4)
C7—C6—C5	119.2 (4)	N6—C18—H18A	119.3
C7—C6—H6A	120.4	C15—C18—H18A	119.3
C5—C6—H6A	120.4	F3—B1—F4	114.3 (5)
N3—C7—C6	123.1 (4)	F3—B1—F2	110.8 (5)
N3—C7—H7A	118.4	F4—B1—F2	108.0 (4)
C6—C7—H7A	118.4	F3—B1—F1	108.7 (4)
N3—C8—C9	124.3 (3)	F4—B1—F1	109.2 (5)
N3—C8—H8A	117.8	F2—B1—F1	105.4 (4)
C9—C8—H8A	117.8		
C4—N1—C1—C2	4.9 (6)	C14—N4—C10—C11	-3.5 (7)
C3—N2—C2—C1	-0.5 (7)	Ag1—N4—C10—C11	176.8 (4)
N1—C1—C2—N2	-3.1 (7)	N4—C10—C11—C12	3.9 (8)
C2—N2—C3—C4	2.0 (6)	C10—C11—C12—C13	-1.4 (7)
C1—N1—C4—C3	-3.4 (5)	C10—C11—C12—S2	178.4 (4)
C1—N1—C4—S1	-179.6 (3)	C15—S2—C12—C13	-11.0 (4)
N2—C3—C4—N1	-0.1 (6)	C15—S2—C12—C11	169.1 (3)
N2—C3—C4—S1	176.1 (3)	C11—C12—C13—C14	-1.1 (6)
C5—S1—C4—N1	-40.5 (3)	S2—C12—C13—C14	179.1 (3)
C5—S1—C4—C3	143.3 (3)	C10—N4—C14—C13	0.6 (6)
C4—S1—C5—C6	158.8 (3)	Ag1—N4—C14—C13	-179.6 (3)
C4—S1—C5—C9	-21.7 (4)	C12—C13—C14—N4	1.6 (7)
C9—C5—C6—C7	-1.5 (7)	C16—N5—C15—C18	-2.2 (6)
S1—C5—C6—C7	178.1 (3)	C16—N5—C15—S2	180.0 (3)
C8—N3—C7—C6	-1.4 (6)	C12—S2—C15—N5	-40.4 (3)
Ag1—N3—C7—C6	176.3 (3)	C12—S2—C15—C18	141.7 (3)
C5—C6—C7—N3	2.7 (7)	C15—N5—C16—C17	4.3 (6)
C7—N3—C8—C9	-1.1 (6)	C18—N6—C17—C16	-1.0 (7)
Ag1—N3—C8—C9	-178.8 (3)	N5—C16—C17—N6	-2.8 (8)
N3—C8—C9—C5	2.1 (7)	C17—N6—C18—C15	3.0 (7)
C6—C5—C9—C8	-0.7 (6)	N5—C15—C18—N6	-1.4 (6)
S1—C5—C9—C8	179.7 (3)	S2—C15—C18—N6	176.4 (3)