

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## 3-Methyl-4-nitrophenol

Su-Lan Dong\* and Xiaochun Cheng

College of Life Science and Chemical Engineering, Huaiyin Institute of Technology,  
Huaiyin 223003, Jiangsu, People's Republic of China  
Correspondence e-mail: dsl710221@163.com


Received 13 January 2012; accepted 20 January 2012

Key indicators: single-crystal X-ray study;  $T = 296$  K; mean  $\sigma(\text{C}-\text{C}) = 0.003$  Å;  
R factor = 0.043; wR factor = 0.148; data-to-parameter ratio = 12.7.

In the title molecule,  $\text{C}_7\text{H}_7\text{NO}_3$ , the nitro group is oriented at  $14.4$  (3)° with respect to the plane of the benzene ring. The crystal structure is stabilized by  $\text{O}-\text{H}\cdots\text{O}$  hydrogen bonds and further consolidated by  $\text{C}-\text{H}\cdots\text{O}$  interactions.

## Related literature

For applications of the title compound in organophosphorus pesticides and for the synthetic procedure, see: Yin & Shi (2005). For a related structure, see: Barve & Pant (1971).



## Experimental

## Crystal data

$\text{C}_7\text{H}_7\text{NO}_3$   
 $M_r = 153.14$   
Monoclinic,  $P2_1/c$   
 $a = 7.2993$  (14) Å  
 $b = 13.023$  (3) Å  
 $c = 7.4445$  (16) Å  
 $\beta = 91.217$  (4)°

$V = 707.5$  (2) Å<sup>3</sup>  
 $Z = 4$   
Mo  $K\alpha$  radiation  
 $\mu = 0.11$  mm<sup>-1</sup>  
 $T = 296$  K  
 $0.20 \times 0.18 \times 0.15$  mm

## Data collection

Enraf–Nonius CAD-4  
diffractometer  
Absorption correction:  $\psi$  scan  
(North *et al.*, 1968)  
 $T_{\min} = 0.978$ ,  $T_{\max} = 0.983$   
3860 measured reflections

1293 independent reflections  
1071 reflections with  $I > 2\sigma$   
 $R_{\text{int}} = 0.023$   
3 standard reflections every 200  
reflections  
intensity decay: 1%

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.043$   
 $wR(F^2) = 0.148$   
 $S = 1.00$   
1293 reflections

102 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.18$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.21$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{O3}-\text{H3}\cdots\text{O2}^{\text{i}}$	0.82	2.00	2.811 (2)	169
$\text{C5}-\text{H5}\cdots\text{O3}^{\text{ii}}$	0.93	2.58	3.437 (2)	154

Symmetry codes: (i)  $x + 1, -y + \frac{1}{2}, z - \frac{1}{2}$ ; (ii)  $-x + 2, -y + 1, -z$ .

Data collection: *CAD-4 Software* (Enraf–Nonius, 1985); cell refinement: *CAD-4 Software*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors thank the Center of Testing and Analysis, Nanjing University, for the data collection. We also thank the Contract grant sponsor, the Natural Science Foundation of Jiangsu Province of China (BK2008195), and the Science Research Foundation of Huaiyin Institute of Technology (2517045) for help.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PV2508).

## References

- Barve, J. V. & Pant, L. M. (1971). *Acta Cryst.* **B27**, 1158–1162.  
Enraf–Nonius (1985). *CAD-4 Software*. Enraf–Nonius, Delft, The Netherlands.  
Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.  
North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.  
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
Yin, W. P. & Shi, M. (2005). *Tetrahedron*, **61**, 10861–10867.

## supporting information

*Acta Cryst.* (2012). E68, o518 [doi:10.1107/S160053681200253X]

## 3-Methyl-4-nitrophenol

Su-Lan Dong and Xiaochun Cheng

### S1. Comment

The title compound is an important intermediate which can be used in many fields such as organophosphorus pesticides (Yin & Shi, 2005).


In the title molecule (Fig. 1), the nitro group (N1/O1/O2) and the benzene ring (C2—C7) are oriented at 14.4 (3) ° with respect to each other. The molecular structure is stabilized by O3—H3···O2 intermolecular hydrogen bonds and further consolidated by C5—H5···O3 type intermolecular hydrogen bonding interactions (Tab. 2 and Fig. 2). The bond lengths and angles in the title compound are in agreement with the corresponding bond lengths and angles reported for a closely related compound (Barve & Pant, 1971).

### S2. Experimental


The title compound was prepared by a method reported in the literature (Yin & Shi, 2005). A solution of sodium nitrite (5.11 g, 74 mmol) and nitric acid (13 g, 208 mmol) in dichloromethane (50 ml) was added slowly to a solution of *m*-cresol (5 g, 46.2 mmol). After stirring for 5 h at a temperature of 323 K, the solvent was evaporated on a rotary evaporator to obtain the title compound. The crystals were grown from an ethanol solution by slow evaporation of the solvent at room temperature in about 7 days.

### S3. Refinement

All H atoms were positioned geometrically and constrained to ride on their parent atoms, with O—H = 0.82 Å and C—H = 0.93 and 0.96 for aryl and alkyl H atoms, respectively, with  $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C})$ , where  $x = 1.2$  for aryl and 1.5 for other H atoms.

**Figure 1**

The molecular structure of the title compound with the atom numbering scheme. Displacement ellipsoids are drawn at the 50% probability level. H atoms are presented as small spheres of arbitrary radius.

**Figure 2**

A view of the O—H···O and C—H···O hydrogen bonds (dotted lines) in the crystal structure of the title compound.

### 3-Methyl-4-nitrophenol

#### Crystal data

$C_7H_7NO_3$

$M_r = 153.14$

Monoclinic,  $P2_1/c$

Hall symbol:  $-P 2_1bc$

$a = 7.2993$  (14) Å

$b = 13.023$  (3) Å

$c = 7.4445$  (16) Å

$\beta = 91.217$  (4)°

$V = 707.5$  (2) Å<sup>3</sup>

$Z = 4$

$F(000) = 320$

$D_x = 1.438$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 2201 reflections

$\theta = 2.8$ – $29.8$ °

$\mu = 0.11$  mm<sup>-1</sup>

$T = 296$  K

Block, colorless

$0.20 \times 0.18 \times 0.15$  mm

#### Data collection

Enraf–Nonius CAD-4  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

$\omega/2\theta$  scans

Absorption correction:  $\psi$  scan  
(North *et al.*, 1968)

$T_{\min} = 0.978$ ,  $T_{\max} = 0.983$

3860 measured reflections

1293 independent reflections

1071 reflections with  $I > 2\sigma$

$R_{\text{int}} = 0.023$

$\theta_{\max} = 25.5$ °,  $\theta_{\min} = 2.8$ °

$h = -8 \rightarrow 8$

$k = -15 \rightarrow 15$   
 $l = -9 \rightarrow 6$

3 standard reflections every 200 reflections  
 intensity decay: 1%

*Refinement*

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.043$   
 $wR(F^2) = 0.148$   
 $S = 1.00$   
 1293 reflections  
 102 parameters  
 0 restraints  
 Primary atom site location: structure-invariant  
 direct methods

Secondary atom site location: difference Fourier  
 map  
 Hydrogen site location: inferred from  
 neighbouring sites  
 H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0847P)^2 + 0.266P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} < 0.001$   
 $\Delta\rho_{\max} = 0.18 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.21 \text{ e } \text{\AA}^{-3}$

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
O3	1.09340 (18)	0.37232 (10)	0.0133 (2)	0.0532 (4)
H3	1.1635	0.3301	-0.0293	0.080*
C7	0.9309 (2)	0.21737 (13)	0.0839 (2)	0.0388 (5)
H7	1.0295	0.1779	0.0476	0.047*
C3	0.6322 (2)	0.23242 (13)	0.1996 (2)	0.0370 (4)
C5	0.8008 (3)	0.38512 (14)	0.1339 (2)	0.0417 (5)
H5	0.8101	0.4563	0.1303	0.050*
C4	0.6461 (2)	0.33913 (14)	0.1958 (2)	0.0402 (5)
H4	0.5497	0.3792	0.2356	0.048*
C1	0.7737 (3)	0.05198 (14)	0.1483 (3)	0.0542 (6)
H1A	0.7391	0.0288	0.2652	0.081*
H1B	0.8940	0.0268	0.1226	0.081*
H1C	0.6877	0.0266	0.0597	0.081*
C2	0.7745 (2)	0.16801 (13)	0.1442 (2)	0.0366 (4)
C6	0.9445 (2)	0.32372 (14)	0.0763 (2)	0.0378 (4)
N1	0.4618 (2)	0.19093 (13)	0.2651 (2)	0.0467 (4)
O2	0.3574 (2)	0.24929 (12)	0.3445 (2)	0.0635 (5)
O1	0.4233 (2)	0.10112 (12)	0.2411 (3)	0.0868 (7)

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O3	0.0443 (8)	0.0454 (8)	0.0708 (10)	-0.0081 (6)	0.0235 (7)	-0.0031 (7)

C7	0.0350 (10)	0.0421 (11)	0.0396 (9)	0.0033 (7)	0.0062 (7)	-0.0029 (7)
C3	0.0335 (10)	0.0432 (10)	0.0344 (9)	-0.0041 (7)	0.0044 (7)	0.0012 (7)
C5	0.0458 (10)	0.0327 (9)	0.0470 (10)	-0.0005 (7)	0.0102 (8)	0.0002 (7)
C4	0.0369 (9)	0.0416 (10)	0.0423 (10)	0.0039 (7)	0.0085 (7)	-0.0010 (7)
C1	0.0606 (13)	0.0382 (11)	0.0643 (13)	-0.0016 (9)	0.0113 (10)	0.0002 (9)
C2	0.0409 (9)	0.0348 (9)	0.0340 (8)	-0.0015 (7)	0.0012 (7)	-0.0006 (7)
C6	0.0367 (9)	0.0396 (10)	0.0373 (9)	-0.0061 (7)	0.0069 (7)	-0.0009 (7)
N1	0.0381 (9)	0.0517 (10)	0.0507 (9)	-0.0072 (7)	0.0075 (7)	0.0021 (7)
O2	0.0437 (8)	0.0656 (10)	0.0823 (11)	0.0005 (7)	0.0253 (8)	0.0006 (8)
O1	0.0719 (12)	0.0569 (10)	0.1333 (17)	-0.0280 (9)	0.0428 (11)	-0.0163 (10)

*Geometric parameters (Å, °)*

O3—C6	1.351 (2)	C5—C6	1.393 (3)
O3—H3	0.8200	C5—H5	0.9300
C7—C6	1.390 (3)	C4—H4	0.9300
C7—C2	1.393 (2)	C1—C2	1.511 (3)
C7—H7	0.9300	C1—H1A	0.9600
C3—C4	1.394 (3)	C1—H1B	0.9600
C3—C2	1.404 (2)	C1—H1C	0.9600
C3—N1	1.450 (2)	N1—O1	1.215 (2)
C5—C4	1.367 (2)	N1—O2	1.236 (2)
C6—O3—H3	109.5	C2—C1—H1B	109.5
C6—C7—C2	122.19 (15)	H1A—C1—H1B	109.5
C6—C7—H7	118.9	C2—C1—H1C	109.5
C2—C7—H7	118.9	H1A—C1—H1C	109.5
C4—C3—C2	122.34 (15)	H1B—C1—H1C	109.5
C4—C3—N1	116.23 (15)	C7—C2—C3	115.82 (15)
C2—C3—N1	121.42 (16)	C7—C2—C1	118.12 (16)
C4—C5—C6	118.99 (16)	C3—C2—C1	126.04 (16)
C4—C5—H5	120.5	O3—C6—C7	122.66 (15)
C6—C5—H5	120.5	O3—C6—C5	117.02 (16)
C5—C4—C3	120.33 (16)	C7—C6—C5	120.31 (15)
C5—C4—H4	119.8	O1—N1—O2	121.27 (16)
C3—C4—H4	119.8	O1—N1—C3	120.45 (16)
C2—C1—H1A	109.5	O2—N1—C3	118.28 (16)
C6—C5—C4—C3	-0.6 (3)	C2—C7—C6—O3	-178.06 (16)
C2—C3—C4—C5	1.1 (3)	C2—C7—C6—C5	1.6 (3)
N1—C3—C4—C5	-179.15 (15)	C4—C5—C6—O3	178.90 (16)
C6—C7—C2—C3	-1.0 (2)	C4—C5—C6—C7	-0.8 (3)
C6—C7—C2—C1	-179.66 (16)	C4—C3—N1—O1	165.72 (19)
C4—C3—C2—C7	-0.4 (2)	C2—C3—N1—O1	-14.6 (3)
N1—C3—C2—C7	179.95 (15)	C4—C3—N1—O2	-13.9 (2)
C4—C3—C2—C1	178.20 (17)	C2—C3—N1—O2	165.86 (17)
N1—C3—C2—C1	-1.5 (3)		

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
O3—H3···O2 <sup>i</sup>	0.82	2.00	2.811 (2)	169
C4—H4···O2	0.93	2.35	2.671 (2)	100
C5—H5···O3 <sup>ii</sup>	0.93	2.58	3.437 (2)	154

Symmetry codes: (i)  $x+1, -y+1/2, z-1/2$ ; (ii)  $-x+2, -y+1, -z$ .