

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

2-(4,5-Diphenyl-2-*p*-tolyl-1*H*-imidazol-1-yl)-3-phenylpropan-1-ol

Yongmei Xiao, Liangru Yang, Kun He, Jinwei Yuan and Pu Mao*

School of Chemistry and Chemical Engineering, Henan University of Technology, Zhengzhou 450001, People's Republic of China
Correspondence e-mail: henangongda@yahoo.com


Received 8 December 2011; accepted 20 December 2011

Key indicators: single-crystal X-ray study; $T = 291$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.037; wR factor = 0.098; data-to-parameter ratio = 14.2.

In the title compound, $\text{C}_{31}\text{H}_{28}\text{N}_2\text{O}$, the dihedral angles formed by the imidazole ring with the three aryl substituents are 18.52 (8) and 85.56 (7) and 85.57 (7)°, respectively. In the crystal, molecules are linked by $\text{O}-\text{H}\cdots\text{N}$ and $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds into chains parallel to the a axis.

Related literature

For the synthesis and properties of chiral ionic liquids, see: Olivier-Bourbigou *et al.* (2010); Chen *et al.* (2008); Mao *et al.* (2010).



Experimental

Crystal data

 $\text{C}_{31}\text{H}_{28}\text{N}_2\text{O}$ $M_r = 444.55$ Orthorhombic, $P2_12_12_1$ $a = 9.3413$ (7) Å $b = 13.7402$ (11) Å $c = 19.6296$ (14) Å $V = 2519.5$ (3) Å³ $Z = 4$ Cu $K\alpha$ radiation $\mu = 0.55$ mm⁻¹ $T = 291$ K

0.25 × 0.20 × 0.20 mm

Data collection

Agilent Xcalibur Eos Gemini diffractometer
Absorption correction: multi-scan (*CrysAlis PRO*; Agilent, 2011)
 $T_{\min} = 0.866$, $T_{\max} = 1.000$

9302 measured reflections
4441 independent reflections
4007 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.027$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.037$ $wR(F^2) = 0.098$ $S = 1.03$

4441 reflections

313 parameters

H atoms treated by a mixture of independent and constrained refinement

 $\Delta\rho_{\max} = 0.12$ e Å⁻³ $\Delta\rho_{\min} = -0.13$ e Å⁻³

Absolute structure: Flack (1983);

1887 Friedel pairs

Flack parameter: -0.1 (3)

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O1}-\text{H1}\cdots\text{N1}^i$	0.82 (3)	2.01 (3)	2.825 (2)	174 (3)
$\text{C16}-\text{H16}\cdots\text{O1}^ii$	0.93	2.56	3.272 (3)	133

Symmetry codes: (i) $x - \frac{1}{2}, -y + \frac{1}{2}, -z + 1$; (ii) $x + 1, y, z$.

Data collection: *CrysAlis PRO* (Agilent, 2011); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *OLEX2* (Dolomanov *et al.*, 2009); software used to prepare material for publication: *OLEX2*.

The authors thank Ms Y. Zhu for technical assistance. This research was supported by the National Natural Science Foundation of P. R. China (Nos. 20902017 and 21172055).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RZ2683).

References

- Agilent (2011). *CrysAlis PRO*. Agilent Technologies Ltd, Yarnton, England.
Chen, X., Li, X., Hu, A. & Wang, F. (2008). *Tetrahedron Asymmetry*, **19**, 1–14.
Dolomanov, O. V., Bourhis, L. J., Gildea, R. J., Howard, J. A. K. & Puschmann, H. (2009). *J. Appl. Cryst.* **42**, 339–341.
Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
Mao, P., Cai, Y., Xiao, Y., Yang, L., Xue, Y. & Song, M. (2010). *Phosphorus Sulfur Silicon Relat. Elem.* **185**, 2418–2425.
Olivier-Bourbigou, H., Magna, L. & Morvan, D. (2010). *Appl. Catal. A*, **373**, 1–56.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supporting information

Acta Cryst. (2012). E68, o264 [doi:10.1107/S1600536811054766]

2-(4,5-Diphenyl-2-*p*-tolyl-1*H*-imidazol-1-yl)-3-phenylpropan-1-ol

Yongmei Xiao, Liangru Yang, Kun He, Jinwei Yuan and Pu Mao

S1. Comment

Ionic liquids (ILs) have received considerable interest in the fields of synthesis, analysis and catalysis due to their unique properties (Olivier-Bourbigou *et al.*, 2010). Chiral ionic liquids (CILs) derived from naturally abundant precursors have also attracted much interest (Chen *et al.*, 2008). Our group is interested in the preparation and application of imidazole derived CILs (Mao *et al.*, 2010), and we observed that the condensation reaction between *L*-phenylalaninol (easily available from *L*-phenylalanine), dibenzoyl, 4-methylbenzaldehyde and ammonium acetate afforded the title compound, a multi-aryl substituted imidazole derivative containing an appended chiral functionality. The chiral C22 carbon atom maintains the *S* configuration observed in *L*-phenylalaninol.


The molecular structure of the title compound is shown in Figure 1. As expected, the imidazole core (N1/C7/C8/N2/C24) is essentially planar. The dihedral angles formed by the imidazole ring with the three aryl substituents are 18.52 (8) (C1–C6), 85.56 (7) (C9–C14) and 85.57 (7)° (C25–C30), respectively. In the crystal structure, molecules are linked by O—H···N and C—H···O hydrogen bonds (Table 1) into chains parallel to the *a* axis.

S2. Experimental

To a solution of *L*-phenylalaninol (15.1 g, 0.1 mol) in MeOH (50 ml) in an ice-bath, a molar equivalent of dibenzoyl, 4-methylbenzaldehyde and ammonium acetate were added. The mixture was kept stirring in the ice-bath until all the solids were dissolved before being heated to 60°C for 5 h. The mixture was then cooled to room temperature and the solvent was removed by evaporation. The residue was washed with H₂O to obtain the crude product. Crystallization of the crude product in EtOH afforded colourless crystals of the title compound.

S3. Refinement

The hydroxyl H atom was located in a difference Fourier map and refined freely. All other H atoms were placed in calculated positions with C—H = 0.93–0.98 Å and refined as riding, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ or $1.5U_{\text{eq}}(\text{C})$ for methyl H atoms.

**Figure 1**

The molecular structure of the title compound showing 30% probability displacement ellipsoids. Hydrogen atoms are omitted for clarity.

2-(4,5-Diphenyl-2-*p*-tolyl-1*H*-imidazol-1-yl)-3-phenylpropan-1-ol

Crystal data

$C_{31}H_{28}N_2O$

$M_r = 444.55$

Orthorhombic, $P2_12_12_1$

Hall symbol: P 2ac 2ab

$a = 9.3413$ (7) Å

$b = 13.7402$ (11) Å

$c = 19.6296$ (14) Å

$V = 2519.5$ (3) Å³

$Z = 4$

$F(000) = 944$

$D_x = 1.172$ Mg m⁻³

Cu $K\alpha$ radiation, $\lambda = 1.5418$ Å

Cell parameters from 3569 reflections

$\theta = 3.2$ – 67.0°

$\mu = 0.55$ mm⁻¹

$T = 291$ K

Prismatic, colourless

$0.25 \times 0.20 \times 0.20$ mm

Data collection

Agilent Xcalibur Eos Gemini
diffractometer

Radiation source: Enhance (Cu) X-ray Source
Graphite monochromator

Detector resolution: 16.2312 pixels mm⁻¹

ω scans

Absorption correction: multi-scan
(*CrysAlis PRO*; Agilent, 2011)

$T_{\min} = 0.866$, $T_{\max} = 1.000$

9302 measured reflections

4441 independent reflections

4007 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.027$

$\theta_{\max} = 66.9^\circ$, $\theta_{\min} = 3.9^\circ$

$h = -9 \rightarrow 11$

$k = -16 \rightarrow 13$

$l = -21 \rightarrow 23$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.037$

$wR(F^2) = 0.098$

$S = 1.03$

4441 reflections

313 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H atoms treated by a mixture of independent
and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0481P)^2 + 0.1385P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.12 \text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\min} = -0.13 \text{ e } \text{\AA}^{-3}$

Extinction correction: *SHELXL97* (Sheldrick,
2008), $F_c^* = kF_c[1 + 0.001 \times F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$

Extinction coefficient: 0.0029 (2)

Absolute structure: Flack (1983); 1887 Friedel
pairs

Absolute structure parameter: -0.1 (3)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.03250 (17)	0.28449 (11)	0.37156 (10)	0.0793 (5)
N1	0.38409 (17)	0.31434 (10)	0.52389 (7)	0.0523 (3)
N2	0.32181 (16)	0.32519 (10)	0.41559 (7)	0.0479 (3)
C1	0.3733 (2)	0.45666 (14)	0.63052 (9)	0.0622 (5)
H1A	0.3616	0.3912	0.6413	0.075*
C2	0.3881 (3)	0.52437 (18)	0.68230 (11)	0.0786 (6)
H2	0.3855	0.5040	0.7275	0.094*
C3	0.4063 (4)	0.62017 (19)	0.66770 (12)	0.0932 (8)
H3	0.4181	0.6652	0.7026	0.112*
C4	0.4069 (4)	0.64981 (17)	0.60113 (13)	0.1027 (10)
H4	0.4174	0.7155	0.5909	0.123*
C5	0.3922 (3)	0.58302 (15)	0.54886 (11)	0.0797 (7)
H5	0.3933	0.6043	0.5039	0.096*
C6	0.3757 (2)	0.48512 (13)	0.56267 (9)	0.0547 (4)
C7	0.3616 (2)	0.41103 (12)	0.50849 (8)	0.0497 (4)
C8	0.32230 (19)	0.41909 (11)	0.44146 (8)	0.0474 (4)
C9	0.2811 (2)	0.50422 (12)	0.39900 (8)	0.0518 (4)
C10	0.1384 (3)	0.52990 (14)	0.39201 (10)	0.0646 (5)
H10	0.0678	0.4931	0.4133	0.077*
C11	0.1006 (3)	0.61040 (17)	0.35333 (12)	0.0844 (7)
H11	0.0046	0.6273	0.3489	0.101*
C12	0.2033 (4)	0.66518 (16)	0.32168 (13)	0.0948 (9)

H12	0.1776	0.7197	0.2964	0.114*
C13	0.3428 (4)	0.63930 (17)	0.32753 (13)	0.0941 (9)
H13	0.4124	0.6759	0.3054	0.113*
C14	0.3837 (3)	0.55919 (14)	0.36594 (11)	0.0719 (6)
H14	0.4799	0.5425	0.3694	0.086*
C15	0.6439 (3)	0.33594 (17)	0.33126 (13)	0.0782 (6)
H15	0.6546	0.2910	0.3663	0.094*
C16	0.7497 (3)	0.4045 (2)	0.32048 (17)	0.0972 (8)
H16	0.8298	0.4058	0.3486	0.117*
C17	0.7373 (3)	0.4703 (2)	0.26865 (16)	0.0901 (7)
H17	0.8090	0.5160	0.2611	0.108*
C18	0.6186 (3)	0.46844 (18)	0.22809 (12)	0.0836 (7)
H18	0.6088	0.5135	0.1931	0.100*
C19	0.5128 (3)	0.39962 (16)	0.23896 (10)	0.0694 (5)
H19	0.4333	0.3984	0.2105	0.083*
C20	0.5231 (2)	0.33255 (13)	0.29131 (9)	0.0551 (4)
C21	0.4054 (2)	0.25978 (13)	0.30306 (9)	0.0559 (4)
H21A	0.4449	0.2037	0.3265	0.067*
H21B	0.3697	0.2378	0.2593	0.067*
C22	0.2799 (2)	0.29999 (12)	0.34487 (8)	0.0482 (4)
H22	0.2516	0.3611	0.3230	0.058*
C23	0.1490 (2)	0.23495 (13)	0.34372 (9)	0.0562 (4)
H23A	0.1276	0.2161	0.2972	0.067*
H23B	0.1675	0.1763	0.3698	0.067*
C24	0.35972 (19)	0.26487 (12)	0.46788 (8)	0.0478 (4)
C25	0.3693 (2)	0.15676 (12)	0.46470 (8)	0.0505 (4)
C26	0.2543 (3)	0.10109 (16)	0.48526 (13)	0.0757 (6)
H26	0.1712	0.1313	0.5004	0.091*
C27	0.2621 (3)	0.00056 (17)	0.48343 (15)	0.0865 (7)
H27	0.1833	-0.0358	0.4973	0.104*
C28	0.3829 (3)	-0.04669 (14)	0.46169 (11)	0.0725 (6)
C29	0.4981 (3)	0.00908 (15)	0.44320 (11)	0.0697 (5)
H29	0.5818	-0.0214	0.4290	0.084*
C30	0.4930 (2)	0.10967 (13)	0.44520 (11)	0.0611 (5)
H30	0.5735	0.1457	0.4333	0.073*
C31	0.3906 (5)	-0.15668 (17)	0.45750 (18)	0.1156 (12)
H31A	0.3258	-0.1847	0.4900	0.173*
H31B	0.4864	-0.1777	0.4674	0.173*
H31C	0.3645	-0.1774	0.4125	0.173*
H1	-0.008 (3)	0.2522 (19)	0.4008 (15)	0.090 (9)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0651 (9)	0.0671 (9)	0.1056 (12)	0.0056 (7)	0.0218 (8)	0.0281 (9)
N1	0.0603 (8)	0.0461 (7)	0.0505 (7)	0.0012 (6)	-0.0088 (6)	0.0028 (6)
N2	0.0599 (8)	0.0403 (6)	0.0434 (7)	0.0029 (6)	0.0022 (6)	0.0007 (5)
C1	0.0732 (12)	0.0594 (10)	0.0540 (9)	0.0054 (10)	-0.0049 (9)	-0.0026 (7)

C2	0.1002 (17)	0.0851 (15)	0.0506 (10)	0.0051 (13)	-0.0054 (11)	-0.0118 (10)
C3	0.135 (2)	0.0790 (15)	0.0656 (13)	-0.0143 (16)	0.0045 (14)	-0.0287 (11)
C4	0.173 (3)	0.0568 (12)	0.0780 (15)	-0.0268 (16)	0.0106 (17)	-0.0160 (11)
C5	0.129 (2)	0.0531 (10)	0.0574 (10)	-0.0127 (12)	0.0033 (12)	-0.0049 (8)
C6	0.0599 (10)	0.0525 (9)	0.0517 (9)	0.0002 (8)	-0.0003 (8)	-0.0063 (7)
C7	0.0563 (9)	0.0440 (8)	0.0489 (8)	-0.0007 (7)	0.0002 (7)	-0.0013 (6)
C8	0.0543 (9)	0.0406 (7)	0.0472 (8)	0.0015 (7)	0.0059 (7)	-0.0005 (6)
C9	0.0729 (11)	0.0409 (8)	0.0417 (8)	0.0063 (8)	0.0082 (7)	-0.0005 (6)
C10	0.0804 (13)	0.0544 (10)	0.0588 (10)	0.0156 (10)	0.0056 (9)	0.0002 (8)
C11	0.116 (2)	0.0647 (12)	0.0726 (13)	0.0359 (14)	-0.0086 (13)	-0.0029 (11)
C12	0.162 (3)	0.0499 (11)	0.0730 (14)	0.0280 (16)	0.0052 (16)	0.0124 (10)
C13	0.147 (3)	0.0552 (12)	0.0799 (15)	-0.0041 (15)	0.0280 (17)	0.0176 (10)
C14	0.0925 (16)	0.0530 (10)	0.0701 (12)	-0.0028 (10)	0.0186 (12)	0.0086 (9)
C15	0.0719 (13)	0.0700 (12)	0.0927 (15)	0.0040 (11)	-0.0099 (12)	0.0185 (11)
C16	0.0627 (14)	0.0957 (18)	0.133 (2)	-0.0085 (14)	-0.0198 (15)	0.0150 (17)
C17	0.0751 (15)	0.0857 (16)	0.109 (2)	-0.0166 (13)	0.0163 (14)	0.0086 (15)
C18	0.0989 (18)	0.0788 (14)	0.0731 (13)	-0.0138 (13)	0.0097 (13)	0.0166 (11)
C19	0.0784 (13)	0.0721 (12)	0.0577 (10)	-0.0050 (11)	-0.0024 (10)	0.0057 (9)
C20	0.0588 (10)	0.0537 (9)	0.0528 (10)	0.0077 (8)	0.0093 (8)	-0.0044 (7)
C21	0.0692 (11)	0.0478 (9)	0.0507 (9)	0.0043 (8)	0.0052 (8)	-0.0058 (7)
C22	0.0614 (10)	0.0412 (7)	0.0422 (8)	0.0013 (7)	0.0021 (7)	0.0012 (6)
C23	0.0647 (11)	0.0535 (9)	0.0505 (9)	-0.0047 (8)	-0.0015 (8)	0.0071 (7)
C24	0.0526 (9)	0.0429 (8)	0.0479 (8)	0.0019 (7)	-0.0032 (7)	0.0025 (6)
C25	0.0601 (10)	0.0430 (8)	0.0484 (8)	-0.0011 (8)	-0.0097 (8)	0.0038 (6)
C26	0.0677 (13)	0.0595 (11)	0.0997 (17)	-0.0002 (10)	0.0061 (12)	0.0179 (11)
C27	0.0852 (16)	0.0605 (12)	0.1136 (19)	-0.0212 (12)	-0.0098 (14)	0.0231 (12)
C28	0.1010 (17)	0.0450 (9)	0.0716 (12)	-0.0019 (11)	-0.0277 (12)	0.0060 (8)
C29	0.0806 (14)	0.0497 (10)	0.0788 (13)	0.0136 (10)	-0.0095 (11)	0.0008 (9)
C30	0.0617 (11)	0.0486 (9)	0.0731 (11)	0.0009 (9)	-0.0096 (9)	0.0038 (8)
C31	0.167 (3)	0.0480 (12)	0.131 (3)	-0.0091 (17)	-0.024 (2)	0.0048 (13)

Geometric parameters (Å, °)

O1—C23	1.395 (3)	C15—C16	1.382 (4)
O1—H1	0.82 (3)	C15—C20	1.375 (3)
N1—C7	1.379 (2)	C16—H16	0.9300
N1—C24	1.312 (2)	C16—C17	1.366 (4)
N2—C8	1.387 (2)	C17—H17	0.9300
N2—C22	1.483 (2)	C17—C18	1.365 (4)
N2—C24	1.366 (2)	C18—H18	0.9300
C1—H1A	0.9300	C18—C19	1.384 (3)
C1—C2	1.385 (3)	C19—H19	0.9300
C1—C6	1.388 (3)	C19—C20	1.383 (3)
C2—H2	0.9300	C20—C21	1.503 (3)
C2—C3	1.358 (4)	C21—H21A	0.9700
C3—H3	0.9300	C21—H21B	0.9700
C3—C4	1.369 (4)	C21—C22	1.534 (2)
C4—H4	0.9300	C22—H22	0.9800

C4—C5	1.383 (3)	C22—C23	1.515 (3)
C5—H5	0.9300	C23—H23A	0.9700
C5—C6	1.381 (3)	C23—H23B	0.9700
C6—C7	1.478 (2)	C24—C25	1.489 (2)
C7—C8	1.371 (2)	C25—C26	1.379 (3)
C8—C9	1.487 (2)	C25—C30	1.379 (3)
C9—C10	1.386 (3)	C26—H26	0.9300
C9—C14	1.382 (3)	C26—C27	1.384 (3)
C10—H10	0.9300	C27—H27	0.9300
C10—C11	1.387 (3)	C27—C28	1.370 (4)
C11—H11	0.9300	C28—C29	1.370 (4)
C11—C12	1.368 (4)	C28—C31	1.515 (3)
C12—H12	0.9300	C29—H29	0.9300
C12—C13	1.355 (5)	C29—C30	1.384 (3)
C13—H13	0.9300	C30—H30	0.9300
C13—C14	1.388 (3)	C31—H31A	0.9600
C14—H14	0.9300	C31—H31B	0.9600
C15—H15	0.9300	C31—H31C	0.9600
C23—O1—H1	111.8 (19)	C18—C17—H17	120.3
C24—N1—C7	106.80 (13)	C17—C18—H18	119.9
C8—N2—C22	124.11 (13)	C17—C18—C19	120.2 (2)
C24—N2—C8	106.77 (13)	C19—C18—H18	119.9
C24—N2—C22	129.06 (13)	C18—C19—H19	119.3
C2—C1—H1A	119.6	C20—C19—C18	121.3 (2)
C2—C1—C6	120.89 (19)	C20—C19—H19	119.3
C6—C1—H1A	119.6	C15—C20—C19	117.2 (2)
C1—C2—H2	119.7	C15—C20—C21	122.33 (18)
C3—C2—C1	120.6 (2)	C19—C20—C21	120.43 (18)
C3—C2—H2	119.7	C20—C21—H21A	108.8
C2—C3—H3	120.3	C20—C21—H21B	108.8
C2—C3—C4	119.4 (2)	C20—C21—C22	113.64 (14)
C4—C3—H3	120.3	H21A—C21—H21B	107.7
C3—C4—H4	119.7	C22—C21—H21A	108.8
C3—C4—C5	120.7 (2)	C22—C21—H21B	108.8
C5—C4—H4	119.7	N2—C22—C21	112.53 (15)
C4—C5—H5	119.6	N2—C22—H22	106.3
C6—C5—C4	120.8 (2)	N2—C22—C23	111.38 (13)
C6—C5—H5	119.6	C21—C22—H22	106.3
C1—C6—C7	119.66 (16)	C23—C22—C21	113.37 (14)
C5—C6—C1	117.68 (17)	C23—C22—H22	106.3
C5—C6—C7	122.65 (17)	O1—C23—C22	109.65 (15)
N1—C7—C6	119.49 (14)	O1—C23—H23A	109.7
C8—C7—N1	109.22 (14)	O1—C23—H23B	109.7
C8—C7—C6	131.23 (15)	C22—C23—H23A	109.7
N2—C8—C9	121.73 (14)	C22—C23—H23B	109.7
C7—C8—N2	106.10 (14)	H23A—C23—H23B	108.2
C7—C8—C9	132.14 (15)	N1—C24—N2	111.12 (14)

C10—C9—C8	120.32 (17)	N1—C24—C25	122.76 (14)
C14—C9—C8	120.90 (18)	N2—C24—C25	126.10 (14)
C14—C9—C10	118.77 (18)	C26—C25—C24	119.62 (18)
C9—C10—H10	119.9	C30—C25—C24	122.01 (17)
C9—C10—C11	120.1 (2)	C30—C25—C26	118.28 (17)
C11—C10—H10	119.9	C25—C26—H26	119.8
C10—C11—H11	119.7	C25—C26—C27	120.4 (2)
C12—C11—C10	120.6 (3)	C27—C26—H26	119.8
C12—C11—H11	119.7	C26—C27—H27	119.2
C11—C12—H12	120.3	C28—C27—C26	121.6 (2)
C13—C12—C11	119.4 (2)	C28—C27—H27	119.2
C13—C12—H12	120.3	C27—C28—C29	117.69 (18)
C12—C13—H13	119.4	C27—C28—C31	121.9 (3)
C12—C13—C14	121.3 (3)	C29—C28—C31	120.4 (3)
C14—C13—H13	119.4	C28—C29—H29	119.2
C9—C14—C13	119.8 (3)	C28—C29—C30	121.6 (2)
C9—C14—H14	120.1	C30—C29—H29	119.2
C13—C14—H14	120.1	C25—C30—C29	120.4 (2)
C16—C15—H15	119.2	C25—C30—H30	119.8
C20—C15—H15	119.2	C29—C30—H30	119.8
C20—C15—C16	121.5 (2)	C28—C31—H31A	109.5
C15—C16—H16	119.9	C28—C31—H31B	109.5
C17—C16—C15	120.3 (2)	C28—C31—H31C	109.5
C17—C16—H16	119.9	H31A—C31—H31B	109.5
C16—C17—H17	120.3	H31A—C31—H31C	109.5
C18—C17—C16	119.4 (2)	H31B—C31—H31C	109.5
N1—C7—C8—N2	-0.31 (19)	C11—C12—C13—C14	-1.1 (4)
N1—C7—C8—C9	177.69 (18)	C12—C13—C14—C9	0.1 (4)
N1—C24—C25—C26	82.9 (2)	C14—C9—C10—C11	-1.1 (3)
N1—C24—C25—C30	-93.5 (2)	C15—C16—C17—C18	0.7 (5)
N2—C8—C9—C10	84.7 (2)	C15—C20—C21—C22	97.1 (2)
N2—C8—C9—C14	-95.4 (2)	C16—C15—C20—C19	1.1 (4)
N2—C22—C23—O1	63.07 (19)	C16—C15—C20—C21	-178.8 (2)
N2—C24—C25—C26	-95.2 (2)	C16—C17—C18—C19	-0.8 (4)
N2—C24—C25—C30	88.4 (2)	C17—C18—C19—C20	1.1 (4)
C1—C2—C3—C4	1.4 (5)	C18—C19—C20—C15	-1.2 (3)
C1—C6—C7—N1	-17.1 (3)	C18—C19—C20—C21	178.7 (2)
C1—C6—C7—C8	159.8 (2)	C19—C20—C21—C22	-82.8 (2)
C2—C1—C6—C5	-0.5 (3)	C20—C15—C16—C17	-0.9 (5)
C2—C1—C6—C7	179.2 (2)	C20—C21—C22—N2	-64.97 (19)
C2—C3—C4—C5	-1.3 (6)	C20—C21—C22—C23	167.51 (15)
C3—C4—C5—C6	0.3 (5)	C21—C22—C23—O1	-168.81 (14)
C4—C5—C6—C1	0.5 (4)	C22—N2—C8—C7	177.67 (16)
C4—C5—C6—C7	-179.1 (3)	C22—N2—C8—C9	-0.6 (3)
C5—C6—C7—N1	162.6 (2)	C22—N2—C24—N1	-177.41 (16)
C5—C6—C7—C8	-20.5 (3)	C22—N2—C24—C25	0.9 (3)
C6—C1—C2—C3	-0.5 (4)	C24—N1—C7—C6	177.73 (16)

C6—C7—C8—N2	-177.50 (18)	C24—N1—C7—C8	0.2 (2)
C6—C7—C8—C9	0.5 (3)	C24—N2—C8—C7	0.34 (19)
C7—N1—C24—N2	0.1 (2)	C24—N2—C8—C9	-177.92 (16)
C7—N1—C24—C25	-178.29 (17)	C24—N2—C22—C21	-69.1 (2)
C7—C8—C9—C10	-93.1 (3)	C24—N2—C22—C23	59.5 (2)
C7—C8—C9—C14	86.8 (2)	C24—C25—C26—C27	-179.2 (2)
C8—N2—C22—C21	114.21 (17)	C24—C25—C30—C29	179.66 (18)
C8—N2—C22—C23	-117.21 (17)	C25—C26—C27—C28	0.3 (4)
C8—N2—C24—N1	-0.3 (2)	C26—C25—C30—C29	3.2 (3)
C8—N2—C24—C25	178.03 (17)	C26—C27—C28—C29	1.6 (4)
C8—C9—C10—C11	178.86 (17)	C26—C27—C28—C31	-178.1 (3)
C8—C9—C14—C13	-178.9 (2)	C27—C28—C29—C30	-1.0 (3)
C9—C10—C11—C12	0.1 (3)	C28—C29—C30—C25	-1.4 (3)
C10—C9—C14—C13	1.0 (3)	C30—C25—C26—C27	-2.7 (3)
C10—C11—C12—C13	1.0 (4)	C31—C28—C29—C30	178.7 (2)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
O1—H1...N1 ⁱ	0.82 (3)	2.01 (3)	2.825 (2)	174 (3)
C16—H16...O1 ⁱⁱ	0.93	2.56	3.272 (3)	133

Symmetry codes: (i) $x-1/2, -y+1/2, -z+1$; (ii) $x+1, y, z$.