

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

# A second monoclinic polymorph of (pyridine-2-carboxaldehyde oximato- $\kappa^2N,N'$ )(pyridine-2-carboxaldehyde oxime- $\kappa^2N,N'$ )palladium(II) chloride

Kwang Ha

School of Applied Chemical Engineering, The Research Institute of Catalysis, Chonnam National University, Gwangju 500-757, Republic of Korea

Correspondence e-mail: hakwang@chonnam.ac.kr


Received 10 January 2012; accepted 13 January 2012

Key indicators: single-crystal X-ray study;  $T = 200$  K; mean  $\sigma(C-C) = 0.008$  Å;  $R$  factor = 0.030;  $wR$  factor = 0.075; data-to-parameter ratio = 13.5.

The asymmetric unit of the title compound,  $[Pd(C_6H_5N_2O)(C_6H_6N_2O)]Cl$ , contains one half of a cationic  $Pd^{II}$  complex and a  $Cl^-$  anion, with a crystallographic mirror plane parallel to the  $ac$  plane passing through the Pd and Cl atoms. In the complex, the  $Pd^{II}$  ion is four-coordinated in a distorted square-planar environment by four N atoms derived from the two chelating ligands. The hydroxy H atom lies on the mirror plane and so is equidistant from the O atoms. This indicates that the negative charge is delocalized over the two O atoms. The complex molecules are stacked in columns along the  $c$  axis and are connected by  $C-H \cdots O$  hydrogen bonds, forming a three-dimensional network. The structure reported herein represents a new monoclinic polymorph of the previously reported monoclinic ( $C2/c$ ) form [Torabi *et al.* (2007). *Z. Kristallogr. New Cryst. Struct.* **222**, 197–198].

## Related literature

For the  $C2/c$  polymorph of the title compound, see: Torabi *et al.* (2007). For the crystal structure of the related complex  $[PdCl_2(C_6H_6N_2O)]$ , see: Ha (2011).



## Experimental

## Crystal data

$[Pd(C_6H_5N_2O)(C_6H_6N_2O)]Cl$   
 $M_r = 385.10$   
 Monoclinic,  $C2/m$   
 $a = 14.0865$  (16) Å  
 $b = 12.1439$  (14) Å  
 $c = 8.2723$  (9) Å  
 $\beta = 114.447$  (2)°

$V = 1288.2$  (3) Å<sup>3</sup>  
 $Z = 4$   
 Mo  $K\alpha$  radiation  
 $\mu = 1.65$  mm<sup>-1</sup>  
 $T = 200$  K  
 $0.22 \times 0.15 \times 0.10$  mm

## Data collection

Bruker SMART 1000 CCD diffractometer  
 Absorption correction: multi-scan (*SADABS*; Bruker, 2000)  
 $T_{min} = 0.894$ ,  $T_{max} = 1.000$

3935 measured reflections  
 1312 independent reflections  
 1188 reflections with  $I > 2\sigma(I)$   
 $R_{int} = 0.025$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.030$   
 $wR(F^2) = 0.075$   
 $S = 1.09$   
 1312 reflections  
 97 parameters

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{max} = 0.75$  e Å<sup>-3</sup>  
 $\Delta\rho_{min} = -0.53$  e Å<sup>-3</sup>

Table 1

Selected geometric parameters (Å, °).

Pd1—N1	2.068 (3)	Pd1—N2	1.985 (3)
N2—Pd1—N1	78.91 (14)		

Table 2

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
O1—H1 $o \cdots$ O1 <sup>i</sup>	1.21 (1)	1.21 (1)	2.418 (7)	175 (7)
C2—H2 $ \cdots$ O1 <sup>ii</sup>	0.95	2.49	3.178 (5)	129

Symmetry codes: (i)  $x, -y + 1, z$ ; (ii)  $x - \frac{1}{2}, -y + \frac{1}{2}, z - 1$ .

Data collection: *SMART* (Bruker, 2000); cell refinement: *SAINT* (Bruker, 2000); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

This work was supported by the Priority Research Centers Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (2010-0029626).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK5047).

## References

- Bruker (2000). *SADABS*, *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Ha, K. (2011). *Z. Kristallogr. New Cryst. Struct.* **226**, 525–526.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Torabi, A. A., Soudozi, A. & Welter, R. (2007). *Z. Kristallogr. New Cryst. Struct.* **222**, 197–198.

## supporting information

*Acta Cryst.* (2012). E68, m176–m177 [doi:10.1107/S1600536812001559]

## A second monoclinic polymorph of (pyridine-2-carboxaldehyde oximate- $\kappa^2N,N'$ )(pyridine-2-carboxaldehyde oxime- $\kappa^2N,N'$ )palladium(II) chloride

Kwang Ha

### S1. Comment

The title compound,  $[\text{Pd}(\text{C}_{12}\text{H}_{11}\text{N}_4\text{O}_2)]\text{Cl}$ , was obtained as a by-product from the reaction of  $\text{Na}_2\text{PdCl}_4$  with *syn*-2-pyridinealdoxime in  $\text{H}_2\text{O}$ . The X-ray crystal structure of the compound was previously reported in the monoclinic space group  $C2/c$  (Torabi *et al.*, 2007). In the latter, the molecule does not exhibit molecular symmetry. The structure presented herein is essentially the same as the published structure but represents a new monoclinic polymorph with the space group  $C2/m$ . The yellow main product of the reaction,  $[\text{PdCl}_2(\text{C}_6\text{H}_6\text{N}_2\text{O})]$ , was investigated previously (Ha, 2011).

The asymmetric unit of the title compound contains one half of a cationic  $\text{Pd}^{\text{II}}$  complex and a  $\text{Cl}^-$  anion (Fig. 1). The compound is disposed about a crystallographic mirror plane parallel to the *ac* plane passing through the Pd and Cl atoms. In the complex, the  $\text{Pd}^{\text{II}}$  ion is four-coordinated in a distorted square-planar environment by four N atoms of the two chelating ligands. Formally, one of the ligands is coordinated to the Pd atom in the monoanionic form, but the negative charge is delocalized over the two O atoms of the ligands; the hydroxy H atom is located in the middle of the O atoms forming a nearly planar six-membered ring. The tight  $\text{N1—Pd1—N2}$  chelate angle of  $78.91(14)^\circ$  contributes the distortion of the square planar structure. The *trans*  $\text{N1—Pd1—N2}^i$  (symmetry code *i*:  $x, 1 - y, z$ ) bond angle is  $173.62(12)^\circ$ . The  $\text{Pd1—N1}$ (pyridine) bond length is slightly longer than the  $\text{Pd1—N2}$ (oxime) bond length (Table 1). The ligands are nearly planar, with a maximum deviation of  $0.024(3) \text{ \AA}$  from the least-squares plane, and the dihedral angle between the ligands is  $5.06(8)^\circ$ . The complex molecules are stacked in columns along the *c* axis and are connected by intermolecular  $\text{C—H}\cdots\text{O}$  hydrogen bonds, forming a three-dimensional network (Fig. 2 and Table 2). In the columns, intermolecular  $\pi$ - $\pi$  interactions between the pyridine rings are present, the shortest ring centroid-centroid distance being  $3.787(3) \text{ \AA}$ .


### S2. Experimental

The title compound was obtained as a by-product from the reaction of  $\text{Na}_2\text{PdCl}_4$  (0.2942 g, 1.000 mmol) with *syn*-2-pyridinealdoxime (0.2444 g, 2.001 mmol) in  $\text{H}_2\text{O}$  (20 ml). After stirring of the reaction mixture for 3 h at room temperature, the formed precipitate was separated by filtration, washed with  $\text{H}_2\text{O}$  and acetone, to give the main product as a yellow powder (0.2302 g) (Ha, 2011). The orange by-product in a small amount was obtained from the mixture of filtrate and washing solution. Crystals suitable for X-ray analysis were obtained by slow evaporation from an *N,N*-dimethylformamide (DMF) solution of the by-product at  $60^\circ\text{C}$ .

### S3. Refinement


H atoms were positioned geometrically and allowed to ride on their respective parent atoms [ $\text{C—H} = 0.95 \text{ \AA}$  and  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ ]. The hydroxy H atom was located from Fourier difference maps and refined isotropically. A number of outlying reflections, *i.e.* (6 0 7), (9 7 4), ( $\bar{1}1$  9 7), ( $\bar{1}2$  2 9), ( $\bar{1}2$  4 9), ( $\bar{1}0$  8 8), ( $\bar{1}7$  1 6), ( $\bar{8}$  8 8), ( $\bar{1}0$  0 10), ( $\bar{1}2$  0 9), ( $\bar{6}$  14 1),

( $\bar{1}$  1 5 9) and (0 2 4), were omitted owing to poor agreement.



**Figure 1**

A view of the molecular structure of the title compound, with displacement ellipsoids drawn at the 30% probability level. Unlabelled atoms are related to the reference atoms by the  $(x, 1-y, z)$  symmetry transformation.



**Figure 2**

A view of the crystal packing of the title compound. Intermolecular hydrogen-bond interactions are drawn with dashed lines.

**(pyridine-2-carboxaldehyde oximato- $\kappa^2N,N'$ )(pyridine-2-carboxaldehyde oxime- $\kappa^2N,N'$ )palladium(II) chloride**

*Crystal data*

[Pd(C<sub>6</sub>H<sub>5</sub>N<sub>2</sub>O)(C<sub>6</sub>H<sub>6</sub>N<sub>2</sub>O)]Cl

$M_r = 385.10$

Monoclinic,  $C2/m$

Hall symbol:  $-C\ 2y$

$a = 14.0865\ (16)\ \text{\AA}$

$b = 12.1439\ (14)\ \text{\AA}$

$c = 8.2723\ (9)\ \text{\AA}$

$\beta = 114.447\ (2)^\circ$

$V = 1288.2\ (3)\ \text{\AA}^3$

$Z = 4$

$F(000) = 760$

$D_x = 1.986\ \text{Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 2872 reflections

$\theta = 2.3\text{--}26.0^\circ$

$\mu = 1.65\ \text{mm}^{-1}$

$T = 200\ \text{K}$

Prism, orange

$0.22 \times 0.15 \times 0.10\ \text{mm}$

*Data collection*

Bruker SMART 1000 CCD  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

$\varphi$  and  $\omega$  scans

Absorption correction: multi-scan  
(*SADABS*; Bruker, 2000)

$T_{\min} = 0.894$ ,  $T_{\max} = 1.000$

3935 measured reflections

1312 independent reflections

1188 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.025$   
 $\theta_{\text{max}} = 26.0^\circ$ ,  $\theta_{\text{min}} = 2.3^\circ$

$h = -17 \rightarrow 17$   
 $k = -12 \rightarrow 14$   
 $l = -10 \rightarrow 10$

### Refinement

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.030$   
 $wR(F^2) = 0.075$   
 $S = 1.09$   
 1312 reflections  
 97 parameters  
 0 restraints  
 Primary atom site location: structure-invariant  
 direct methods

Secondary atom site location: difference Fourier  
 map  
 Hydrogen site location: inferred from  
 neighbouring sites  
 H atoms treated by a mixture of independent  
 and constrained refinement  
 $w = 1/[\sigma^2(F_o^2) + (0.035P)^2 + 2.4037P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\text{max}} < 0.001$   
 $\Delta\rho_{\text{max}} = 0.75 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -0.53 \text{ e } \text{\AA}^{-3}$

### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
Pd1	0.09319 (3)	0.5000	0.41451 (4)	0.02259 (15)
O1	0.2571 (2)	0.4005 (3)	0.7251 (4)	0.0546 (9)
H1o	0.261 (5)	0.5000	0.725 (10)	0.08 (2)*
N1	0.0142 (2)	0.3633 (2)	0.2741 (4)	0.0295 (7)
N2	0.1746 (2)	0.3793 (3)	0.5732 (4)	0.0342 (7)
C1	-0.0700 (3)	0.3564 (4)	0.1201 (5)	0.0444 (11)
H1	-0.1006	0.4219	0.0576	0.053*
C2	-0.1134 (4)	0.2558 (6)	0.0502 (8)	0.075 (2)
H2	-0.1730	0.2539	-0.0599	0.090*
C3	-0.0738 (6)	0.1612 (5)	0.1328 (11)	0.092 (3)
H3	-0.1051	0.0926	0.0848	0.110*
C4	0.0139 (5)	0.1668 (4)	0.2904 (9)	0.0721 (18)
H4	0.0452	0.1013	0.3524	0.086*
C5	0.0562 (3)	0.2684 (3)	0.3576 (6)	0.0413 (10)
C6	0.1470 (3)	0.2806 (4)	0.5255 (6)	0.0457 (11)
H6	0.1831	0.2191	0.5946	0.055*
Cl1	0.23809 (11)	0.0000	0.72959 (19)	0.0424 (3)

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Pd1	0.0211 (2)	0.0221 (2)	0.0211 (2)	0.000	0.00530 (15)	0.000
O1	0.0321 (15)	0.080 (2)	0.0384 (16)	0.0115 (16)	0.0015 (13)	0.0273 (17)
N1	0.0320 (15)	0.0266 (17)	0.0349 (17)	-0.0057 (13)	0.0187 (14)	-0.0088 (13)
N2	0.0270 (15)	0.044 (2)	0.0316 (17)	0.0108 (14)	0.0122 (13)	0.0164 (15)
C1	0.036 (2)	0.058 (3)	0.042 (2)	-0.018 (2)	0.0183 (19)	-0.023 (2)
C2	0.066 (3)	0.100 (5)	0.078 (4)	-0.058 (4)	0.048 (3)	-0.066 (4)
C3	0.120 (6)	0.059 (4)	0.141 (6)	-0.064 (4)	0.098 (5)	-0.069 (4)
C4	0.109 (4)	0.025 (2)	0.127 (5)	-0.012 (3)	0.093 (4)	-0.015 (3)
C5	0.051 (2)	0.027 (2)	0.066 (3)	-0.0016 (18)	0.044 (2)	-0.002 (2)
C6	0.048 (2)	0.037 (2)	0.068 (3)	0.020 (2)	0.040 (2)	0.027 (2)
Cl1	0.0381 (7)	0.0499 (9)	0.0418 (8)	0.000	0.0190 (6)	0.000

Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )

Pd1—N1	2.068 (3)	C1—H1	0.9500
Pd1—N1 <sup>i</sup>	2.068 (3)	C2—C3	1.335 (10)
Pd1—N2 <sup>i</sup>	1.985 (3)	C2—H2	0.9500
Pd1—N2	1.985 (3)	C3—C4	1.377 (9)
O1—N2	1.337 (4)	C3—H3	0.9500
O1—H1 <sub>o</sub>	1.210 (5)	C4—C5	1.383 (6)
N1—C1	1.336 (5)	C4—H4	0.9500
N1—C5	1.347 (5)	C5—C6	1.455 (6)
N2—C6	1.271 (5)	C6—H6	0.9500
C1—C2	1.381 (7)		
N2 <sup>i</sup> —Pd1—N2	95.2 (2)	C3—C2—C1	121.9 (6)
N2 <sup>i</sup> —Pd1—N1	173.62 (12)	C3—C2—H2	119.1
N2—Pd1—N1	78.91 (14)	C1—C2—H2	119.1
N2 <sup>i</sup> —Pd1—N1 <sup>i</sup>	78.91 (14)	C2—C3—C4	117.6 (5)
N2—Pd1—N1 <sup>i</sup>	173.62 (13)	C2—C3—H3	121.2
N1—Pd1—N1 <sup>i</sup>	106.87 (17)	C4—C3—H3	121.2
N2—O1—H1 <sub>o</sub>	103 (3)	C3—C4—C5	119.5 (5)
C1—N1—C5	117.6 (4)	C3—C4—H4	120.2
C1—N1—Pd1	130.1 (3)	C5—C4—H4	120.2
C5—N1—Pd1	112.3 (3)	N1—C5—C4	122.2 (5)
C6—N2—O1	120.6 (3)	N1—C5—C6	115.3 (3)
C6—N2—Pd1	118.2 (3)	C4—C5—C6	122.5 (5)
O1—N2—Pd1	121.3 (3)	N2—C6—C5	115.3 (4)
N1—C1—C2	121.3 (5)	N2—C6—H6	122.3
N1—C1—H1	119.4	C5—C6—H6	122.3
C2—C1—H1	119.4		
N2—Pd1—N1—C1	178.8 (3)	C2—C3—C4—C5	1.1 (8)
N1 <sup>i</sup> —Pd1—N1—C1	1.6 (4)	C1—N1—C5—C4	-1.1 (5)
N2—Pd1—N1—C5	0.0 (2)	Pd1—N1—C5—C4	177.9 (3)

N1 <sup>i</sup> —Pd1—N1—C5	-177.22 (17)	C1—N1—C5—C6	-179.4 (3)
N2 <sup>i</sup> —Pd1—N2—C6	178.0 (2)	Pd1—N1—C5—C6	-0.4 (4)
N1—Pd1—N2—C6	0.4 (3)	C3—C4—C5—N1	0.1 (7)
N2 <sup>i</sup> —Pd1—N2—O1	-2.8 (3)	C3—C4—C5—C6	178.3 (4)
N1—Pd1—N2—O1	179.6 (3)	O1—N2—C6—C5	-180.0 (3)
C5—N1—C1—C2	0.9 (5)	Pd1—N2—C6—C5	-0.7 (5)
Pd1—N1—C1—C2	-177.9 (3)	N1—C5—C6—N2	0.7 (5)
N1—C1—C2—C3	0.3 (7)	C4—C5—C6—N2	-177.6 (4)
C1—C2—C3—C4	-1.3 (8)		

Symmetry code: (i)  $x, -y+1, z$ .

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
O1—H1 <sup><i>o</i></sup> ...O1 <sup><i>i</i></sup>	1.21 (1)	1.21 (1)	2.418 (7)	175 (7)
C2—H2...O1 <sup><i>ii</i></sup>	0.95	2.49	3.178 (5)	129

Symmetry codes: (i)  $x, -y+1, z$ ; (ii)  $x-1/2, -y+1/2, z-1$ .