

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

(2E)-2-(4-Bromobenzylidene)-2,3-dihydro-1H-inden-1-one

 Abdullah M. Asiri,^{a,b,‡} Hassan M. Faidallah,^a Khulud F. Al-Nemari,^{a,b} Seik Weng Ng^c and Edward R. T. Tiekink^{c*}

^aChemistry Department, Faculty of Science, King Abdulaziz University, PO Box 80203, Jeddah, Saudi Arabia, ^bThe Center of Excellence for Advanced Materials Research, King Abdulaziz University, Jeddah, PO Box 80203, Saudi Arabia, and ^cDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
Correspondence e-mail: edward.tiekink@gmail.com


Received 13 February 2012; accepted 14 February 2012

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.036; wR factor = 0.068; data-to-parameter ratio = 16.5.

The title inden-1-one derivative, $\text{C}_{16}\text{H}_{11}\text{BrO}$, is planar, the r.m.s. deviation for all 18 non-H atoms being 0.071 Å. The configuration about the $\text{C}=\text{C}$ bond [1.337 (5) Å] is *E*. In the crystal, supramolecular layers in the *ab* plane are formed by $\text{C}-\text{H}\cdots\text{O}$ interactions, involving the bifurcated carbonyl O atom, as well as $\text{C}-\text{H}\cdots\pi$ interactions. The studied crystal was an inversion twin.

Related literature

For the activity of related species for the treatment of Chagas disease, see: Vera-DiVaio *et al.* (2009).



Experimental

Crystal data

$\text{C}_{16}\text{H}_{11}\text{BrO}$	$V = 617.61$ (11) Å ³
$M_r = 299.16$	$Z = 2$
Monoclinic, Pn	Mo $K\alpha$ radiation
$a = 6.1359$ (5) Å	$\mu = 3.31$ mm ⁻¹
$b = 4.7512$ (4) Å	$T = 100$ K
$c = 21.310$ (3) Å	$0.20 \times 0.10 \times 0.05$ mm
$\beta = 96.195$ (9)°	

Data collection

Agilent SuperNova Dual diffractometer with an Atlas detector	5083 measured reflections
Absorption correction: multi-scan (<i>CrysAlis PRO</i> ; Agilent, 2011)	2705 independent reflections
$T_{\min} = 0.798$, $T_{\max} = 1.000$	2455 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.039$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.036$	H-atom parameters constrained
$wR(F^2) = 0.068$	$\Delta\rho_{\text{max}} = 0.70$ e Å ⁻³
$S = 0.99$	$\Delta\rho_{\text{min}} = -0.35$ e Å ⁻³
2705 reflections	Absolute structure: Flack (1983),
164 parameters	1288 Friedel pairs
2 restraints	Flack parameter: 0.343 (10)

Table 1

Hydrogen-bond geometry (Å, °).

Cg1 is the centroid of the C2–C7 ring.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C1}-\text{H1b}\cdots\text{O1}^{\text{i}}$	0.99	2.35	3.220 (5)	146
$\text{C15}-\text{H15}\cdots\text{O1}^{\text{ii}}$	0.95	2.54	3.171 (5)	124
$\text{C1}-\text{H1A}\cdots\text{Cg1}^{\text{iii}}$	0.99	2.61	3.479 (4)	147

 Symmetry codes: (i) $x + 1, y, z$; (ii) $x + 1, y - 1, z$; (iii) $x, y + 1, z$.

Data collection: *CrysAlis PRO* (Agilent, 2011); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *DIAMOND* (Brandenburg, 2006); software used to prepare material for publication: *publCIF* (Westrip, 2010).

The authors are thankful to the Center of Excellence for Advanced Materials Research and the Chemistry Department of King Abdulaziz University for providing research facilities. We also thank the Ministry of Higher Education (Malaysia) for funding structural studies through the High-Impact Research scheme (UM.C/HIR/MOHE/SC/12).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5818).

References

- Agilent (2011). *CrysAlis PRO*. Agilent Technologies, Yarnton, Oxfordshire, England.
- Brandenburg, K. (2006). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Vera-DiVaio, M. A. F., Freitas, A. C. C., Castro, F. H. C., de Albuquerque, S., Cabral, L. M., Rodrigues, C. R., Albuquerque, M. G., Martins, R. C. A., Henriques, M. G. M. O. & Dias, L. R. S. (2009). *Bioorg. Med. Chem.* **17**, 295–302.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

[‡] Additional correspondence author, e-mail: aasiri2@kau.edu.sa.

supporting information

Acta Cryst. (2012). E68, o755 [doi:10.1107/S1600536812006654]

(2*E*)-2-(4-Bromobenzylidene)-2,3-dihydro-1*H*-inden-1-one

Abdullah M. Asiri, Hassan M. Faidallah, Khulud F. Al-Nemari, Seik Weng Ng and Edward R. T. Tiekink

S1. Comment

The motivation for the investigation of the title compound, 2-(4-bromobenzylidene)indan-1-one (I), is its relationship to some active compounds developed for the treatment of Chagas disease (Vera-DiVaio *et al.*, 2009).

The molecule of (I), Fig. 1, is planar with a r.m.s. deviation for all 18 non-hydrogen atoms = 0.071 Å. The maximum deviations are found for the Br1 [-0.195 (1) Å] and C16 [-0.096 (3) Å] atoms. The configuration about the C9=C10 bond [1.337 (5) Å] is *E*.


In the crystal packing, C—H \cdots O interactions involving the bifurcated carbonyl-O atom as well as C—H \cdots π interactions link molecules into layers in the *ab* plane, Fig. 2 and Table 1. The layers stack along the *c* axis with no specific interactions between them, Fig. 2.

S2. Experimental


A solution of the 4-bromobenzaldehyde (1.8 g, 0.01 mol) in ethanol (20 ml) was added to a stirred solution of 1-indanone (1.3 g, 0.01 mol) in ethanolic KOH (20%, 20 ml), and stirring was maintained at room temperature for 6 h. The reaction mixture was then poured onto water (200 ml) and set aside overnight. The precipitated solid product was collected by filtration, washed with water, dried and recrystallized from its ethanol solution as blocks; *M.pt*: 453–455 K.

S3. Refinement


H-atoms were placed in calculated positions [C—H = 0.95 to 0.99 Å, $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$] and were included in the refinement in the riding model approximation. The structure was refined as a racemic twin; the Flack parameter (Flack, 1983) was explicitly refined to 0.343 (10) indicating the fractional contribution of the minor twin component. Owing to poor agreement, the (3 1 $\bar{6}$) and (3 0 $\bar{5}$) reflections were omitted from the final cycles of refinement.

**Figure 1**

The molecular structure of (I) showing the atom-labelling scheme and displacement ellipsoids at the 50% probability level.

**Figure 2**

A view of the supramolecular layer in (I). The C—H...O and C—H... π interactions are shown as orange and purple dashed lines, respectively.

**Figure 3**

A view in projection down the a axis of the unit-cell contents of (I). The C—H...O and C—H... π interactions are shown as orange and purple dashed lines, respectively.

(2*E*)-2-(4-Bromobenzylidene)-2,3-dihydro-1*H*-inden-1-one

Crystal data

$C_{16}H_{11}BrO$

$M_r = 299.16$

Monoclinic, Pn

Hall symbol: $P -2yc$

$a = 6.1359 (5) \text{ \AA}$

$b = 4.7512 (4) \text{ \AA}$

$c = 21.310 (3) \text{ \AA}$

$\beta = 96.195 (9)^\circ$

$V = 617.61 (11) \text{ \AA}^3$

$Z = 2$

$F(000) = 300$

$D_x = 1.609 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 2365 reflections

$\theta = 2.9\text{--}27.5^\circ$

$\mu = 3.31 \text{ mm}^{-1}$

$T = 100 \text{ K}$

Prism, orange

$0.20 \times 0.10 \times 0.05 \text{ mm}$

Data collection

Agilent SuperNova Dual
diffractometer with an Atlas detector
Radiation source: SuperNova (Mo) X-ray
Source

Mirror monochromator

Detector resolution: $10.4041 \text{ pixels mm}^{-1}$

ω scan

Absorption correction: multi-scan
(*CrysAlis PRO*; Agilent, 2011)

$T_{\min} = 0.798$, $T_{\max} = 1.000$

5083 measured reflections

2705 independent reflections

2455 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.039$

$\theta_{\max} = 27.6^\circ$, $\theta_{\min} = 3.4^\circ$

$h = -7 \rightarrow 7$

$k = -6 \rightarrow 6$

$l = -27 \rightarrow 27$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.036$

$wR(F^2) = 0.068$

$S = 0.99$

2705 reflections

164 parameters

2 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0265P)^2]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.70 \text{ e \AA}^{-3}$

$\Delta\rho_{\min} = -0.35 \text{ e \AA}^{-3}$

Absolute structure: Flack (1983), 1288 Friedel pairs

Absolute structure parameter: 0.343 (10)

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.50001 (3)	0.38435 (6)	0.489989 (19)	0.02212 (10)
O1	-0.1668 (4)	1.6578 (5)	0.71975 (12)	0.0216 (6)
C1	0.3820 (7)	1.3644 (7)	0.74899 (19)	0.0166 (9)
H1A	0.4144	1.1744	0.7667	0.020*
H1B	0.4993	1.4196	0.7230	0.020*
C2	0.3597 (7)	1.5761 (7)	0.80064 (19)	0.0161 (9)
C3	0.5184 (7)	1.6540 (7)	0.84972 (18)	0.0194 (8)
H3	0.6596	1.5695	0.8541	0.023*
C4	0.4630 (7)	1.8602 (7)	0.89226 (18)	0.0235 (9)
H4	0.5677	1.9145	0.9262	0.028*
C5	0.2575 (7)	1.9869 (8)	0.88561 (19)	0.0219 (9)
H5	0.2247	2.1283	0.9147	0.026*
C6	0.0998 (7)	1.9098 (8)	0.83714 (17)	0.0198 (8)
H6	-0.0412	1.9947	0.8328	0.024*
C7	0.1545 (6)	1.7034 (7)	0.79481 (17)	0.0166 (8)
C8	0.0200 (7)	1.5875 (8)	0.73932 (19)	0.0162 (9)
C9	0.1582 (6)	1.3722 (7)	0.71053 (19)	0.0137 (8)
C10	0.0759 (6)	1.2291 (7)	0.65939 (17)	0.0155 (8)
H10	-0.0727	1.2694	0.6449	0.019*
C11	0.1822 (6)	1.0181 (7)	0.62235 (17)	0.0147 (8)
C12	0.0596 (6)	0.9077 (7)	0.56840 (17)	0.0184 (8)
H12	-0.0877	0.9670	0.5581	0.022*
C13	0.1497 (6)	0.7144 (8)	0.53015 (17)	0.0197 (9)
H13	0.0652	0.6404	0.4939	0.024*
C14	0.3645 (6)	0.6305 (7)	0.54539 (17)	0.0168 (8)
C15	0.4890 (6)	0.7281 (8)	0.59864 (18)	0.0184 (8)
H15	0.6355	0.6652	0.6088	0.022*
C16	0.3964 (6)	0.9197 (7)	0.63707 (17)	0.0177 (8)
H16	0.4803	0.9856	0.6742	0.021*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.02551 (18)	0.02170 (16)	0.01955 (16)	0.0011 (2)	0.00433 (13)	-0.0025 (2)
O1	0.0155 (14)	0.0269 (14)	0.0218 (14)	0.0010 (11)	-0.0003 (12)	-0.0013 (11)
C1	0.020 (2)	0.017 (2)	0.0131 (19)	0.0002 (16)	0.0018 (18)	0.0031 (15)
C2	0.022 (2)	0.0155 (18)	0.0107 (19)	-0.0047 (16)	-0.0006 (18)	0.0059 (16)
C3	0.0197 (19)	0.0189 (18)	0.0184 (19)	-0.0011 (16)	-0.0036 (16)	0.0032 (16)
C4	0.032 (2)	0.022 (2)	0.0157 (19)	-0.0059 (17)	-0.0009 (17)	0.0035 (16)
C5	0.029 (2)	0.0162 (19)	0.021 (2)	-0.0011 (17)	0.0057 (18)	0.0020 (17)
C6	0.022 (2)	0.021 (2)	0.018 (2)	-0.0028 (17)	0.0079 (16)	0.0013 (16)
C7	0.020 (2)	0.0139 (17)	0.0160 (18)	-0.0033 (15)	0.0024 (16)	0.0027 (15)
C8	0.018 (2)	0.018 (2)	0.0130 (19)	-0.0058 (17)	0.0037 (18)	0.0039 (16)
C9	0.0134 (18)	0.0120 (19)	0.016 (2)	-0.0036 (14)	0.0034 (17)	0.0046 (15)
C10	0.0146 (18)	0.0200 (19)	0.0122 (17)	-0.0016 (15)	0.0025 (15)	0.0032 (16)
C11	0.019 (2)	0.0118 (17)	0.0138 (19)	-0.0043 (15)	0.0035 (16)	0.0034 (15)
C12	0.0180 (19)	0.0155 (18)	0.0215 (19)	-0.0006 (15)	0.0010 (16)	0.0050 (16)
C13	0.020 (2)	0.0188 (19)	0.0187 (19)	-0.0022 (15)	-0.0054 (17)	0.0021 (15)
C14	0.023 (2)	0.0132 (17)	0.0157 (18)	-0.0038 (15)	0.0068 (16)	0.0025 (14)
C15	0.0158 (19)	0.0207 (18)	0.0187 (18)	0.0008 (15)	0.0022 (16)	0.0051 (16)
C16	0.020 (2)	0.0183 (19)	0.0149 (18)	-0.0040 (16)	0.0012 (15)	0.0013 (16)

Geometric parameters (Å, °)

Br1—C14	1.914 (4)	C7—C8	1.474 (5)
O1—C8	1.223 (5)	C8—C9	1.501 (5)
C1—C2	1.508 (5)	C9—C10	1.337 (5)
C1—C9	1.522 (5)	C10—C11	1.471 (5)
C1—H1A	0.9900	C10—H10	0.9500
C1—H1B	0.9900	C11—C16	1.398 (5)
C2—C7	1.390 (5)	C11—C12	1.406 (5)
C2—C3	1.400 (5)	C12—C13	1.382 (5)
C3—C4	1.401 (5)	C12—H12	0.9500
C3—H3	0.9500	C13—C14	1.382 (5)
C4—C5	1.391 (6)	C13—H13	0.9500
C4—H4	0.9500	C14—C15	1.378 (5)
C5—C6	1.386 (5)	C15—C16	1.387 (5)
C5—H5	0.9500	C15—H15	0.9500
C6—C7	1.398 (5)	C16—H16	0.9500
C6—H6	0.9500		
C2—C1—C9	103.2 (3)	C7—C8—C9	106.9 (3)
C2—C1—H1A	111.1	C10—C9—C8	119.9 (4)
C9—C1—H1A	111.1	C10—C9—C1	131.8 (4)
C2—C1—H1B	111.1	C8—C9—C1	108.3 (3)
C9—C1—H1B	111.1	C9—C10—C11	129.3 (3)
H1A—C1—H1B	109.1	C9—C10—H10	115.3
C7—C2—C3	120.1 (4)	C11—C10—H10	115.3

C7—C2—C1	112.2 (3)	C16—C11—C12	117.8 (3)
C3—C2—C1	127.7 (4)	C16—C11—C10	124.6 (3)
C2—C3—C4	118.1 (4)	C12—C11—C10	117.6 (3)
C2—C3—H3	121.0	C13—C12—C11	121.2 (4)
C4—C3—H3	121.0	C13—C12—H12	119.4
C5—C4—C3	121.2 (4)	C11—C12—H12	119.4
C5—C4—H4	119.4	C14—C13—C12	119.0 (3)
C3—C4—H4	119.4	C14—C13—H13	120.5
C6—C5—C4	120.9 (4)	C12—C13—H13	120.5
C6—C5—H5	119.5	C15—C14—C13	121.8 (3)
C4—C5—H5	119.5	C15—C14—Br1	118.3 (3)
C5—C6—C7	118.0 (4)	C13—C14—Br1	119.9 (3)
C5—C6—H6	121.0	C14—C15—C16	118.8 (3)
C7—C6—H6	121.0	C14—C15—H15	120.6
C2—C7—C6	121.7 (3)	C16—C15—H15	120.6
C2—C7—C8	109.4 (3)	C15—C16—C11	121.5 (3)
C6—C7—C8	128.8 (4)	C15—C16—H16	119.3
O1—C8—C7	126.5 (4)	C11—C16—H16	119.3
O1—C8—C9	126.6 (4)		
C9—C1—C2—C7	-1.2 (4)	O1—C8—C9—C1	177.3 (4)
C9—C1—C2—C3	-179.9 (4)	C7—C8—C9—C1	-1.2 (4)
C7—C2—C3—C4	0.4 (5)	C2—C1—C9—C10	-178.9 (4)
C1—C2—C3—C4	179.0 (4)	C2—C1—C9—C8	1.5 (4)
C2—C3—C4—C5	-0.8 (5)	C8—C9—C10—C11	177.7 (4)
C3—C4—C5—C6	1.0 (6)	C1—C9—C10—C11	-1.9 (7)
C4—C5—C6—C7	-0.7 (6)	C9—C10—C11—C16	2.8 (6)
C3—C2—C7—C6	-0.2 (6)	C9—C10—C11—C12	-177.2 (4)
C1—C2—C7—C6	-178.9 (3)	C16—C11—C12—C13	-1.8 (5)
C3—C2—C7—C8	179.3 (3)	C10—C11—C12—C13	178.2 (3)
C1—C2—C7—C8	0.5 (5)	C11—C12—C13—C14	-0.3 (5)
C5—C6—C7—C2	0.3 (5)	C12—C13—C14—C15	1.9 (5)
C5—C6—C7—C8	-179.0 (4)	C12—C13—C14—Br1	-175.5 (3)
C2—C7—C8—O1	-178.1 (4)	C13—C14—C15—C16	-1.3 (5)
C6—C7—C8—O1	1.3 (7)	Br1—C14—C15—C16	176.2 (3)
C2—C7—C8—C9	0.4 (4)	C14—C15—C16—C11	-1.0 (5)
C6—C7—C8—C9	179.9 (4)	C12—C11—C16—C15	2.5 (5)
O1—C8—C9—C10	-2.3 (6)	C10—C11—C16—C15	-177.5 (3)
C7—C8—C9—C10	179.1 (3)		

Hydrogen-bond geometry (\AA , $^\circ$)

Cg1 is the centroid of the C2–C7 ring.

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C1—H1b \cdots O1 ⁱ	0.99	2.35	3.220 (5)	146

C15—H15 \cdots O1 ⁱⁱ	0.95	2.54	3.171 (5)	124
C1—H1A \cdots Cg1 ⁱⁱⁱ	0.99	2.61	3.479 (4)	147

Symmetry codes: (i) $x+1, y, z$; (ii) $x+1, y-1, z$; (iii) $x, y+1, z$.