

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

# N-(4-Chloro-3-nitrophenyl)maleamic acid

 U. Chaithanya,<sup>a</sup> Sabine Foro<sup>b</sup> and B. Thimme Gowda<sup>a\*</sup>
<sup>a</sup>Department of Chemistry, Mangalore University, Mangalagangothri 574 199, Mangalore, India, and <sup>b</sup>Institute of Materials Science, Darmstadt University of Technology, Petersenstrasse 23, D-64287 Darmstadt, Germany

Correspondence e-mail: gowdabt@yahoo.com


Received 21 February 2012; accepted 22 February 2012

 Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(\text{C}-\text{C}) = 0.006$  Å;  $R$  factor = 0.063;  $wR$  factor = 0.180; data-to-parameter ratio = 13.3.

In the molecule of the title compound,  $\text{C}_{10}\text{H}_7\text{ClN}_2\text{O}_5$ , the acyclic  $\text{C}=\text{C}$  double bond is *cis* configured. The  $\text{C}=\text{O}$  and  $\text{O}-\text{H}$  bonds of the acid group are in a relatively rare *anti* position to each other, due to the donation of intramolecular hydrogen bond to the amide by the carboxyl group. The nitro group is significantly twisted [dihedral angle =  $66.9(3)^\circ$ ] out of the plane of the remaining atoms, which are almost coplanar (r.m.s. deviation for non-H atoms except the nitro group =  $0.202$  Å). In the crystal,  $\text{N}-\text{H}\cdots\text{O}$  hydrogen bonds link the molecules into zigzag chains running along the  $b$  axis.

## Related literature

For our studies of the effects of substituents on the structures and other aspects of *N*-(aryl)amides, see: Gowda *et al.* (2000, 2003); Chaithanya *et al.* (2012); *N*-(aryl)methanesulfonamides, see: Gowda *et al.* (2007); *N*-chloroarylamides, see: Jyothi & Gowda (2004) and *N*-bromoarylsulfonamides, see: Usha & Gowda (2006).



## Experimental

## Crystal data

 $\text{C}_{10}\text{H}_7\text{ClN}_2\text{O}_5$   
 $M_r = 270.63$   
 Monoclinic,  $P2_1/c$   
 $a = 9.7187(9)$  Å  
 $b = 13.596(1)$  Å  
 $c = 8.4990(9)$  Å  
 $\beta = 99.64(1)^\circ$ 
 $V = 1107.16(18)$  Å<sup>3</sup>  
 $Z = 4$   
 Mo  $K\alpha$  radiation  
 $\mu = 0.36$  mm<sup>-1</sup>  
 $T = 293$  K  
 $0.42 \times 0.12 \times 0.06$  mm

## Data collection

 Oxford Xcalibur diffractometer  
 with a Sapphire CCD detector  
 Absorption correction: multi-scan  
 (*CrysAlis RED*; Oxford  
 Diffraction, 2009)  
 $T_{\min} = 0.863$ ,  $T_{\max} = 0.979$ 

 4360 measured reflections  
 2243 independent reflections  
 1540 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.025$ 

## Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.063$   
 $wR(F^2) = 0.180$   
 $S = 0.97$   
 2243 reflections  
 169 parameters  
 2 restraints

 H atoms treated by a mixture of  
 independent and constrained  
 refinement  
 $\Delta\rho_{\max} = 0.38$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.25$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O3}-\text{H3O}\cdots\text{O1}$	0.82 (2)	1.67 (2)	2.494 (4)	174 (7)
$\text{N1}-\text{H1N}\cdots\text{O2}^i$	0.86 (2)	2.01 (2)	2.839 (4)	162 (4)

 Symmetry code: (i)  $-x + 1, y - \frac{1}{2}, -z + \frac{3}{2}$ .

Data collection: *CrysAlis CCD* (Oxford Diffraction, 2009); cell refinement: *CrysAlis RED* (Oxford Diffraction, 2009); data reduction: *CrysAlis RED*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

BTG thanks the University Grants Commission, Government of India, New Delhi, for a special grant under a UGC-BSR one-time grant to faculty.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5827).

## References

- Chaithanya, U., Foro, S. & Gowda, B. T. (2012). *Acta Cryst.* E68. In the press.  
 Gowda, B. T., Foro, S. & Fuess, H. (2007). *Acta Cryst.* E63, o2337.  
 Gowda, B. T., Kumar, B. H. A. & Fuess, H. (2000). *Z. Naturforsch. Teil A*, **55**, 721–728.  
 Gowda, B. T., Usha, K. M. & Jayalakshmi, K. L. (2003). *Z. Naturforsch. Teil A*, **58**, 801–806.  
 Jyothi, K. & Gowda, B. T. (2004). *Z. Naturforsch. Teil A*, **59**, 64–68.  
 Oxford Diffraction (2009). *CrysAlis CCD* and *CrysAlis RED*. Oxford Diffraction Ltd, Yarnton, Oxfordshire, England.  
 Sheldrick, G. M. (2008). *Acta Cryst.* A64, 112–122.  
 Spek, A. L. (2009). *Acta Cryst.* D65, 148–155.  
 Usha, K. M. & Gowda, B. T. (2006). *J. Chem. Sci.* **118**, 351–359.

## supporting information

*Acta Cryst.* (2012). E68, o873 [doi:10.1107/S1600536812008021]

## ***N*-(4-Chloro-3-nitrophenyl)maleamic acid**

**U. Chaithanya, Sabine Foro and B. Thimme Gowda**

### **S1. Comment**

As part of our studies on the substituent effects on the structures and other aspects of *N*-(aryl)-amides (Gowda *et al.*, 2000, 2003; Chaithanya *et al.*, 2012), *N*-(aryl)-methanesulfonamides (Gowda *et al.*, 2007); *N*-chloroarylsulfonamides (Jyothi & Gowda, 2004) and *N*-bromoarylsulfonamides (Usha & Gowda, 2006), in the present work, the crystal structure of *N*-(4-Chloro-3-nitrophenyl)maleamic acid has been determined (Fig. 1). The conformations of the N—H and the C=O bonds in the amide segment are *anti* to each other. The N—H bond is also *anti* to the *meta*-nitro group. Further, the conformation of the amide C=O is *anti* to the H atom on the adjacent —CH group, while the carboxyl C=O of the acid segment is *syn* to the adjacent —CH group. Furthermore, the C=O and O—H bond of the acid group are in relatively rare *anti* position to each other, due to the donation of hydrogen bond to the amide by the carboxyl group, similar to that observed in *N*-(3-Chloro-4-methylphenyl)maleamic acid (I) (Chaithanya *et al.*, 2012).

The dihedral angle between the phenyl ring and the amide group in the title compound is 11.52 (27)°, compared to the value of 6.55 (99)° in (I).

In the structure, the pairs of O—H···O and N—H···O intermolecular hydrogen bonds pack the molecules into zigzag chains (Table 1, Fig.2).


### **S2. Experimental**

Maleic anhydride (0.025 mol) in toluene (25 ml) was treated dropwise with 4-chloro-3-nitroaniline (0.025 mol) also in toluene (20 ml) with constant stirring. The resulting mixture was stirred for about 30 min and set aside for an additional 30 min at room temperature for the completion of reaction. The mixture was then treated with dilute hydrochloric acid to remove the unreacted 4-chloro-3-nitroaniline. The resultant solid *N*-(4-Chloro-3-nitrophenyl)maleamic acid was filtered under suction and washed thoroughly with water to remove the unreacted maleic anhydride and maleic acid. It was recrystallized to constant melting point from ethanol. The purity of the compound was checked and characterized by its infrared spectra.


Rod like colorless single crystals of the title compound used in X-ray diffraction studies were grown in an ethanol solution by slow evaporation of the solvent (0.5 g in about 30 ml of ethanol) at room temperature.

### **S3. Refinement**

The H atoms of the NH group and the OH group were located in a difference map and later restrained to the distance N—H = 0.86 (2) Å and O—H = 0.82 (2) Å, respectively. The other H atoms were positioned with idealized geometry using a riding model with the aromatic C—H = 0.93 Å and methylene C—H = 0.97 Å. All H atoms were refined with isotropic displacement parameters set at 1.2 $U_{eq}$ .

**Figure 1**

Molecular structure of the title compound, showing the atom labelling scheme and with displacement ellipsoids drawn at the 50% probability level.

**Figure 2**

Molecular packing of the title compound with hydrogen bonding shown as dashed lines.

### ***N*-(4-Chloro-3-nitrophenyl)maleamic acid**

#### *Crystal data*

$C_{10}H_7ClN_2O_5$

$M_r = 270.63$

Monoclinic,  $P2_1/c$

Hall symbol:  $-P\ 2_1/c$

$a = 9.7187(9)\ \text{\AA}$

$b = 13.596(1)\ \text{\AA}$

$c = 8.4990(9)\ \text{\AA}$

$\beta = 99.64(1)^\circ$

$V = 1107.16(18)\ \text{\AA}^3$

$Z = 4$

$F(000) = 552$

$D_x = 1.624\ \text{Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 1582 reflections

$\theta = 2.6\text{--}27.7^\circ$

$\mu = 0.36\ \text{mm}^{-1}$

$T = 293\ \text{K}$

Rod, colourless

$0.42 \times 0.12 \times 0.06\ \text{mm}$

*Data collection*

Oxford Xcalibur  
diffractometer with a Sapphire CCD detector  
Radiation source: fine-focus sealed tube  
Graphite monochromator  
Rotation method data acquisition using  $\omega$  and  $\varphi$   
scans  
Absorption correction: multi-scan  
(*CrysAlis RED*; Oxford Diffraction, 2009)  
 $T_{\min} = 0.863$ ,  $T_{\max} = 0.979$

4360 measured reflections  
2243 independent reflections  
1540 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.025$   
 $\theta_{\max} = 26.4^\circ$ ,  $\theta_{\min} = 2.6^\circ$   
 $h = -12 \rightarrow 6$   
 $k = -16 \rightarrow 16$   
 $l = -7 \rightarrow 10$

*Refinement*

Refinement on  $F^2$   
Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.063$   
 $wR(F^2) = 0.180$   
 $S = 0.97$   
2243 reflections  
169 parameters  
2 restraints  
Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map  
Hydrogen site location: inferred from  
neighbouring sites  
H atoms treated by a mixture of independent  
and constrained refinement  
 $w = 1/[\sigma^2(F_o^2) + (0.0665P)^2 + 3.5725P]$   
where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} = 0.034$   
 $\Delta\rho_{\max} = 0.38 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.25 \text{ e } \text{\AA}^{-3}$

*Special details*

**Experimental.** *CrysAlis RED* (Oxford Diffraction, 2009) Empirical absorption correction using spherical harmonics, implemented in SCALE3 ABSPACK scaling algorithm.

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	-0.22417 (12)	-0.02257 (9)	0.06740 (14)	0.0526 (4)
O1	0.3160 (3)	0.2207 (2)	0.5078 (4)	0.0543 (10)
O2	0.6054 (3)	0.3654 (2)	0.8733 (4)	0.0527 (9)
O3	0.4475 (4)	0.3576 (2)	0.6601 (4)	0.0582 (10)
H3O	0.402 (6)	0.315 (3)	0.604 (6)	0.087*
O4	-0.0092 (5)	0.2184 (3)	0.0224 (5)	0.0750 (12)
O5	-0.1941 (4)	0.1979 (3)	0.1206 (6)	0.0852 (14)
N1	0.3025 (3)	0.0549 (2)	0.5150 (4)	0.0353 (8)
H1N	0.332 (5)	0.004 (2)	0.569 (5)	0.042*
N2	-0.0772 (4)	0.1760 (3)	0.1083 (4)	0.0420 (9)
C1	0.1800 (4)	0.0411 (3)	0.4016 (5)	0.0320 (9)
C2	0.1145 (4)	0.1155 (3)	0.3048 (5)	0.0337 (9)
H2	0.1534	0.1781	0.3068	0.040*

C3	-0.0098 (4)	0.0945 (3)	0.2053 (5)	0.0331 (9)
C4	-0.0696 (4)	0.0024 (3)	0.1939 (5)	0.0357 (10)
C5	-0.0005 (5)	-0.0720 (3)	0.2858 (5)	0.0441 (11)
H5	-0.0375	-0.1352	0.2787	0.053*
C6	0.1233 (4)	-0.0536 (3)	0.3887 (5)	0.0416 (11)
H6	0.1688	-0.1045	0.4494	0.050*
C7	0.3599 (4)	0.1411 (3)	0.5669 (5)	0.0357 (9)
C8	0.4800 (4)	0.1338 (3)	0.6985 (5)	0.0379 (10)
H8	0.5099	0.0704	0.7281	0.046*
C9	0.5505 (4)	0.2064 (3)	0.7794 (5)	0.0409 (10)
H9	0.6240	0.1848	0.8561	0.049*
C10	0.5357 (4)	0.3159 (3)	0.7725 (5)	0.0384 (10)

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C11	0.0429 (6)	0.0487 (7)	0.0575 (7)	-0.0113 (5)	-0.0165 (5)	-0.0070 (6)
O1	0.0534 (19)	0.0280 (16)	0.067 (2)	0.0018 (14)	-0.0321 (16)	-0.0018 (15)
O2	0.059 (2)	0.0404 (18)	0.0501 (19)	-0.0077 (16)	-0.0147 (16)	-0.0133 (15)
O3	0.066 (2)	0.0288 (16)	0.067 (2)	-0.0032 (15)	-0.0280 (18)	-0.0044 (15)
O4	0.100 (3)	0.053 (2)	0.075 (3)	0.007 (2)	0.020 (2)	0.026 (2)
O5	0.051 (2)	0.071 (3)	0.126 (4)	0.018 (2)	-0.008 (2)	0.027 (3)
N1	0.0350 (18)	0.0245 (17)	0.0408 (19)	-0.0004 (14)	-0.0095 (15)	0.0022 (14)
N2	0.041 (2)	0.0329 (19)	0.045 (2)	-0.0018 (17)	-0.0123 (17)	0.0000 (17)
C1	0.0296 (19)	0.028 (2)	0.035 (2)	-0.0004 (16)	-0.0044 (16)	-0.0027 (16)
C2	0.035 (2)	0.027 (2)	0.035 (2)	-0.0060 (16)	-0.0063 (17)	-0.0014 (16)
C3	0.036 (2)	0.028 (2)	0.033 (2)	0.0013 (17)	-0.0020 (17)	-0.0014 (17)
C4	0.034 (2)	0.032 (2)	0.038 (2)	-0.0053 (16)	-0.0063 (17)	-0.0078 (17)
C5	0.050 (2)	0.027 (2)	0.050 (3)	-0.0077 (19)	-0.007 (2)	-0.0007 (19)
C6	0.044 (2)	0.026 (2)	0.048 (2)	0.0010 (18)	-0.012 (2)	0.0027 (18)
C7	0.032 (2)	0.028 (2)	0.043 (2)	0.0011 (17)	-0.0041 (18)	-0.0006 (17)
C8	0.040 (2)	0.028 (2)	0.041 (2)	0.0037 (18)	-0.0095 (18)	-0.0013 (18)
C9	0.040 (2)	0.039 (2)	0.039 (2)	0.0034 (19)	-0.0092 (19)	0.0011 (18)
C10	0.036 (2)	0.035 (2)	0.041 (2)	-0.0022 (18)	-0.0043 (19)	-0.0050 (18)

*Geometric parameters (Å, °)*

C11—C4	1.729 (4)	C2—C3	1.383 (5)
O1—C7	1.239 (5)	C2—H2	0.9300
O2—C10	1.205 (5)	C3—C4	1.377 (5)
O3—C10	1.302 (5)	C4—C5	1.382 (6)
O3—H3O	0.82 (2)	C5—C6	1.386 (6)
O4—N2	1.209 (5)	C5—H5	0.9300
O5—N2	1.196 (5)	C6—H6	0.9300
N1—C7	1.340 (5)	C7—C8	1.479 (5)
N1—C1	1.413 (5)	C8—C9	1.326 (6)
N1—H1N	0.857 (19)	C8—H8	0.9300
N2—C3	1.469 (5)	C9—C10	1.497 (6)

C1—C2	1.389 (5)	C9—H9	0.9300
C1—C6	1.397 (6)		
C10—O3—H3O	110 (4)	C4—C5—C6	120.8 (4)
C7—N1—C1	126.7 (3)	C4—C5—H5	119.6
C7—N1—H1N	117 (3)	C6—C5—H5	119.6
C1—N1—H1N	115 (3)	C5—C6—C1	120.4 (4)
O5—N2—O4	123.9 (4)	C5—C6—H6	119.8
O5—N2—C3	118.7 (4)	C1—C6—H6	119.8
O4—N2—C3	117.5 (4)	O1—C7—N1	122.3 (3)
C2—C1—C6	119.2 (3)	O1—C7—C8	122.7 (4)
C2—C1—N1	123.8 (3)	N1—C7—C8	115.0 (3)
C6—C1—N1	116.9 (3)	C9—C8—C7	128.0 (4)
C3—C2—C1	118.6 (4)	C9—C8—H8	116.0
C3—C2—H2	120.7	C7—C8—H8	116.0
C1—C2—H2	120.7	C8—C9—C10	133.0 (4)
C4—C3—C2	123.1 (4)	C8—C9—H9	113.5
C4—C3—N2	120.2 (3)	C10—C9—H9	113.5
C2—C3—N2	116.7 (3)	O2—C10—O3	120.1 (4)
C3—C4—C5	117.8 (4)	O2—C10—C9	119.3 (4)
C3—C4—Cl1	122.4 (3)	O3—C10—C9	120.6 (3)
C5—C4—Cl1	119.8 (3)		
C7—N1—C1—C2	12.4 (7)	N2—C3—C4—Cl1	0.0 (6)
C7—N1—C1—C6	-167.4 (4)	C3—C4—C5—C6	1.4 (7)
C6—C1—C2—C3	3.9 (6)	Cl1—C4—C5—C6	-179.7 (4)
N1—C1—C2—C3	-176.0 (4)	C4—C5—C6—C1	0.3 (7)
C1—C2—C3—C4	-2.1 (7)	C2—C1—C6—C5	-3.0 (7)
C1—C2—C3—N2	178.4 (4)	N1—C1—C6—C5	176.8 (4)
O5—N2—C3—C4	57.7 (6)	C1—N1—C7—O1	-6.7 (7)
O4—N2—C3—C4	-123.7 (5)	C1—N1—C7—C8	173.9 (4)
O5—N2—C3—C2	-122.8 (5)	O1—C7—C8—C9	5.4 (8)
O4—N2—C3—C2	55.7 (5)	N1—C7—C8—C9	-175.2 (5)
C2—C3—C4—C5	-0.5 (7)	C7—C8—C9—C10	1.3 (9)
N2—C3—C4—C5	178.9 (4)	C8—C9—C10—O2	171.7 (5)
C2—C3—C4—Cl1	-179.4 (3)	C8—C9—C10—O3	-7.5 (8)

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
O3—H3O...O1	0.82 (2)	1.67 (2)	2.494 (4)	174 (7)
N1—H1N...O2 <sup>i</sup>	0.86 (2)	2.01 (2)	2.839 (4)	162 (4)

Symmetry code: (i)  $-x+1, y-1/2, -z+3/2$ .