

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

4,6-Dinitropyrogallol

Christian Neis, Günter J. Merten and Kaspar Hegetschweiler*

Fachrichtung Chemie, Universität des Saarlandes, Postfach 151150, D-66041 Saarbrücken, Germany

Correspondence e-mail: hegetschweiler@mx.uni-saarland.de


Received 25 January 2012; accepted 3 February 2012

 Key indicators: single-crystal X-ray study; $T = 200$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.031; wR factor = 0.089; data-to-parameter ratio = 10.2.

In the title molecule, $\text{C}_6\text{H}_4\text{N}_2\text{O}_7$, the two nitro groups are tilted with respect to the aromatic ring by 11.2 (1) and 10.9 (1)°. All three hydroxy groups are involved in the formation of bifurcated intra- and intermolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds. The crystal packing exhibits short $\text{O}\cdots\text{O}$ distances of 2.823 (2) Å between two O atoms of the nitro groups.

Related literature

The synthesis of the title compound has recently been reported by Merten *et al.* (2012). The importance of hydrogen bonding, π stacking and donor-acceptor interactions in solid-state structures and in solution has been reviewed by Schneider (2009). The crystal structure of dinitrophenoroglucinol, a constitutional isomer of the title compound, has been described by Schweitzer *et al.* (2008). H atoms were treated as recommended by Müller *et al.* (2006).



Experimental

Crystal data

$\text{C}_6\text{H}_4\text{N}_2\text{O}_7$	$V = 756.5$ (3) Å ³
$M_r = 216.11$	$Z = 4$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 6.7612$ (14) Å	$\mu = 0.18$ mm ⁻¹
$b = 10.878$ (2) Å	$T = 200$ K
$c = 10.297$ (2) Å	$0.60 \times 0.22 \times 0.20$ mm
$\beta = 92.75$ (3)°	

Data collection

Stoe IPDS image plate diffractometer	1488 independent reflections
5824 measured reflections	1326 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.059$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.031$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.089$	$\Delta\rho_{\text{max}} = 0.31$ e Å ⁻³
$S = 1.06$	$\Delta\rho_{\text{min}} = -0.23$ e Å ⁻³
1488 reflections	
146 parameters	
3 restraints	

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O2}-\text{H2O}\cdots\text{O3}$	0.85 (1)	2.14 (2)	2.6467 (14)	118 (2)
$\text{O2}-\text{H2O}\cdots\text{O4}^i$	0.85 (1)	2.27 (2)	2.9531 (14)	138 (2)
$\text{O1}-\text{H1O}\cdots\text{O6}$	0.86 (2)	1.91 (2)	2.6359 (15)	141 (2)
$\text{O1}-\text{H1O}\cdots\text{O6}^{ii}$	0.86 (2)	2.46 (2)	3.1361 (15)	137 (2)
$\text{O3}-\text{H3O}\cdots\text{O4}$	0.83 (2)	1.97 (2)	2.6357 (14)	137 (2)
$\text{O3}-\text{H3O}\cdots\text{O7}^{iii}$	0.83 (2)	2.19 (2)	2.7658 (15)	127 (2)

Symmetry codes: (i) $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$; (ii) $-x - 1, -y + 1, -z + 1$; (iii) $x + 1, -y + \frac{1}{2}, z - \frac{1}{2}$.

Data collection: *IPDS Software* (Stoe & Cie, 1997); cell refinement: *IPDS Software*; data reduction: *IPDS Software*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *DIAMOND* (Brandenburg, 2011); software used to prepare material for publication: *SHELXL97*.

The authors thank Dr Volker Huch (Universität des Saarlandes) for the collection of the data set.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CV5241).

References

- Brandenburg, K. (2011). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Merten, G. J., Neis, C., Stucky, S., Huch, V., Rentschler, E., Natter, H., Hempelmann, R., Stöwe, K. & Hegetschweiler, K. (2012). *Eur. J. Inorg. Chem.* pp. 31–35.
- Müller, P., Herbst-Irmer, R., Spek, A. L., Schneider, T. R. & Sawaya, M. R. (2006). *Crystal Structure Refinement – A Crystallographer's Guide to SHELXL*. Oxford University Press.
- Schneider, H.-J. (2009). *Angew. Chem. Int. Ed.* **48**, 3924–3977.
- Schweitzer, T., Taylor, D., Choppin, G., Neis, C. & Hegetschweiler, K. (2008). *Z. Kristallogr. New Cryst. Struct.* **223**, 369–370.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Stoe & Cie (1997). *IPDS Software*. Stoe & Cie, Darmstadt, Germany.

supporting information

Acta Cryst. (2012). E68, o695 [doi:10.1107/S1600536812004771]

4,6-Dinitropyrogallol

Christian Neis, Günter J. Merten and Kaspar Hegetschweiler

S1. Comment


As expected, the aromatic C₆-ring is planar (the positions of the six C atoms deviate from the mean plane by 0.0025 (8)–0.0196 (8) Å). The C₆-ring and the two nitro groups are also approximately coplanar, allowing significant π -delocalization. In the extended hydrogen bonding network all three phenolic hydroxy groups act as hydrogen donors and the nitro groups as hydrogen acceptors. The O atoms O1 and O6 of two neighbouring molecules form a planar, four-membered O1 \cdots O6 \cdots O1' \cdots O6' ring with two bifurcated hydrogen bonds. Remarkably, the O6 \cdots O6' distance of 2.823 (2) Å (*i.e.* one of the diagonal of the corresponding parallelogram) is somewhat shorter than the sum of the van der Waals radii, although no direct hydrogen bond between these two atoms is operative. Some intermolecular C \cdots C and C \cdots O distances (C1 \cdots C1': 3.279 Å, C5 \cdots O2": 3.046 Å) are slightly shorter than the sum of the van der Waals radii and may be interpreted in terms of π -stacking or weak donor–acceptor interactions.

S2. Experimental


The title compound has been prepared by nitration of pyrogallol-triacetate. ¹H NMR ([D₆]acetone): δ (p.p.m.) = 8.51. ¹³C NMR ([D₆]acetone): δ (p.p.m.) = 114.4, 128.7, 136.6, 148.8. Elemental analysis calculated for C₆H₄N₂O₇ (%): C 33.35, H 1.87, N 12.96; found (%): C 33.19, H 2.24, N 12.83. Single crystals were obtained directly from the reaction mixture at 278 K.

S3. Refinement

All non-hydrogen atoms were refined using anisotropic displacement parameters. H atoms were treated as recommended by Müller *et al.* (2006). A riding model was used for the C-bonded H5. The positional parameters of the O-bonded H1O, H2O and H3O were refined using isotropic displacement parameters which were set to 1.5 U_{eq} of the pivot atom. In addition, a restraint of 0.84 Å was used for the O—H distances.

**Figure 1**

Molecular structure of the title compound. Displacement ellipsoids are drawn at the 50% probability level.

**Figure 2**

Section of the hydrogen bonding network, showing bifurcated hydrogen bonds for all three hydroxy groups (displacement ellipsoids are drawn at the 50% probability level).

4,6-Dinitrobenzene-1,2,3-triol

Crystal data

C₆H₄N₂O₇ $M_r = 216.11$ Monoclinic, $P2_1/c$ $a = 6.7612$ (14) Å $b = 10.878$ (2) Å $c = 10.297$ (2) Å $\beta = 92.75$ (3)° $V = 756.5$ (3) Å³ $Z = 4$ $F(000) = 440$ $D_x = 1.898$ Mg m⁻³Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å $\mu = 0.18$ mm⁻¹ $T = 200$ K

Prism, light brown

 $0.60 \times 0.22 \times 0.20$ mm

Data collection

Stoe IPDS image plate
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 φ scans

5824 measured reflections

1488 independent reflections

1326 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.059$ $\theta_{\text{max}} = 26.0^\circ$, $\theta_{\text{min}} = 2.7^\circ$ $h = -8 \rightarrow 8$ $k = -13 \rightarrow 13$ $l = -12 \rightarrow 12$

Refinement

Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.031$ $wR(F^2) = 0.089$ $S = 1.06$

1488 reflections

146 parameters

3 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sitesH atoms treated by a mixture of independent
and constrained refinement $w = 1/[\sigma^2(F_o^2) + (0.0484P)^2 + 0.1574P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\text{max}} < 0.001$ $\Delta\rho_{\text{max}} = 0.31$ e Å⁻³ $\Delta\rho_{\text{min}} = -0.23$ e Å⁻³Extinction correction: *SHELXL97* (Sheldrick,
2008), $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$

Extinction coefficient: 0.093 (9)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C5	0.03683 (18)	0.22251 (11)	0.46470 (11)	0.0173 (3)
H5	0.0181	0.1542	0.5203	0.021*
C2	0.09313 (18)	0.42549 (12)	0.30137 (11)	0.0176 (3)
C6	-0.10033 (17)	0.31771 (12)	0.45731 (11)	0.0166 (3)
C1	-0.07860 (17)	0.42018 (11)	0.37270 (11)	0.0168 (3)

C4	0.20152 (17)	0.22875 (11)	0.38968 (11)	0.0167 (3)
C3	0.23403 (17)	0.33087 (12)	0.30757 (11)	0.0176 (3)
O2	0.11938 (14)	0.52556 (9)	0.22378 (9)	0.0253 (3)
H2O	0.235 (2)	0.5179 (17)	0.1964 (17)	0.038*
O1	-0.20607 (13)	0.51466 (9)	0.35774 (9)	0.0228 (3)
H1O	-0.310 (2)	0.4983 (17)	0.3996 (17)	0.034*
O3	0.38713 (14)	0.34789 (10)	0.23055 (10)	0.0265 (3)
H3O	0.467 (3)	0.2905 (15)	0.2362 (18)	0.040*
N4	0.34037 (15)	0.12578 (10)	0.39485 (10)	0.0184 (3)
N6	-0.26572 (15)	0.31109 (10)	0.54231 (10)	0.0195 (3)
O4	0.50228 (13)	0.14010 (9)	0.34200 (9)	0.0250 (3)
O5	0.29368 (14)	0.03088 (9)	0.44938 (9)	0.0259 (3)
O6	-0.40392 (14)	0.38686 (10)	0.52605 (9)	0.0288 (3)
O7	-0.26491 (15)	0.23213 (9)	0.62784 (10)	0.0289 (3)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C5	0.0202 (6)	0.0161 (6)	0.0156 (5)	-0.0017 (5)	0.0016 (4)	-0.0004 (5)
C2	0.0208 (6)	0.0168 (6)	0.0152 (6)	-0.0010 (5)	0.0011 (4)	0.0022 (4)
C6	0.0159 (6)	0.0189 (6)	0.0154 (6)	-0.0021 (5)	0.0029 (4)	-0.0024 (4)
C1	0.0179 (6)	0.0171 (6)	0.0151 (6)	0.0020 (5)	-0.0010 (4)	-0.0027 (4)
C4	0.0189 (6)	0.0150 (6)	0.0162 (6)	0.0017 (4)	0.0007 (5)	-0.0014 (4)
C3	0.0165 (6)	0.0208 (6)	0.0158 (6)	-0.0009 (5)	0.0025 (4)	-0.0013 (5)
O2	0.0246 (5)	0.0238 (5)	0.0283 (5)	0.0039 (4)	0.0084 (4)	0.0113 (4)
O1	0.0209 (5)	0.0231 (5)	0.0247 (5)	0.0065 (4)	0.0048 (4)	0.0036 (4)
O3	0.0227 (5)	0.0280 (6)	0.0300 (5)	0.0065 (4)	0.0135 (4)	0.0090 (4)
N4	0.0199 (5)	0.0191 (6)	0.0163 (5)	0.0019 (4)	0.0011 (4)	-0.0012 (4)
N6	0.0189 (5)	0.0196 (6)	0.0204 (5)	-0.0016 (4)	0.0041 (4)	-0.0027 (4)
O4	0.0208 (5)	0.0265 (5)	0.0284 (5)	0.0056 (4)	0.0084 (4)	0.0003 (4)
O5	0.0306 (5)	0.0186 (5)	0.0288 (5)	0.0030 (4)	0.0040 (4)	0.0062 (4)
O6	0.0214 (5)	0.0345 (6)	0.0314 (5)	0.0085 (4)	0.0091 (4)	0.0023 (4)
O7	0.0319 (5)	0.0252 (5)	0.0310 (5)	0.0009 (4)	0.0155 (4)	0.0071 (4)

Geometric parameters (Å, °)

C5—C4	1.3870 (17)	C4—N4	1.4609 (16)
C5—C6	1.3898 (17)	C3—O3	1.3467 (15)
C5—H5	0.9500	O2—H2O	0.846 (14)
C2—O2	1.3669 (15)	O1—H1O	0.859 (15)
C2—C3	1.4019 (18)	O3—H3O	0.827 (15)
C2—C1	1.4046 (17)	N4—O5	1.2237 (15)
C6—C1	1.4267 (17)	N4—O4	1.2554 (14)
C6—N6	1.4545 (15)	N6—O7	1.2301 (15)
C1—O1	1.3454 (15)	N6—O6	1.2512 (15)
C4—C3	1.4195 (18)		
C4—C5—C6	118.97 (11)	C5—C4—N4	118.28 (11)

C4—C5—H5	120.5	O3—C3—C2	114.34 (11)
C6—C5—H5	120.5	O3—C3—C4	127.14 (12)
O2—C2—C3	120.29 (11)	C2—C3—C4	118.51 (11)
O2—C2—C1	118.18 (11)	C2—O2—H2O	105.3 (13)
C3—C2—C1	121.53 (11)	C1—O1—H1O	108.5 (13)
C5—C6—C1	121.85 (11)	C3—O3—H3O	111.9 (13)
C5—C6—N6	117.42 (11)	O5—N4—O4	123.58 (11)
C1—C6—N6	120.71 (11)	O5—N4—C4	118.95 (11)
O1—C1—C2	116.73 (11)	O4—N4—C4	117.47 (10)
O1—C1—C6	125.64 (11)	O7—N6—O6	122.24 (11)
C2—C1—C6	117.61 (11)	O7—N6—C6	119.22 (11)
C5—C4—C3	121.43 (11)	O6—N6—C6	118.54 (10)
C4—C5—C6—C1	1.40 (18)	O2—C2—C3—C4	179.79 (11)
C4—C5—C6—N6	-176.90 (10)	C1—C2—C3—C4	-0.88 (18)
O2—C2—C1—O1	0.65 (16)	C5—C4—C3—O3	179.92 (11)
C3—C2—C1—O1	-178.70 (11)	N4—C4—C3—O3	-1.37 (19)
O2—C2—C1—C6	-177.52 (10)	C5—C4—C3—C2	-1.28 (18)
C3—C2—C1—C6	3.13 (17)	N4—C4—C3—C2	177.43 (10)
C5—C6—C1—O1	178.59 (11)	C5—C4—N4—O5	10.41 (16)
N6—C6—C1—O1	-3.17 (18)	C3—C4—N4—O5	-168.34 (11)
C5—C6—C1—C2	-3.43 (17)	C5—C4—N4—O4	-169.81 (10)
N6—C6—C1—C2	174.81 (10)	C3—C4—N4—O4	11.44 (16)
C6—C5—C4—C3	1.01 (18)	C5—C6—N6—O7	9.35 (16)
C6—C5—C4—N4	-177.72 (10)	C1—C6—N6—O7	-168.96 (11)
O2—C2—C3—O3	-1.27 (17)	C5—C6—N6—O6	-171.03 (10)
C1—C2—C3—O3	178.07 (10)	C1—C6—N6—O6	10.65 (16)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O2—H2O \cdots O3	0.85 (1)	2.14 (2)	2.6467 (14)	118 (2)
O2—H2O \cdots O4 ⁱ	0.85 (1)	2.27 (2)	2.9531 (14)	138 (2)
O1—H1O \cdots O6	0.86 (2)	1.91 (2)	2.6359 (15)	141 (2)
O1—H1O \cdots O6 ⁱⁱ	0.86 (2)	2.46 (2)	3.1361 (15)	137 (2)
O3—H3O \cdots O4	0.83 (2)	1.97 (2)	2.6357 (14)	137 (2)
O3—H3O \cdots O7 ⁱⁱⁱ	0.83 (2)	2.19 (2)	2.7658 (15)	127 (2)

Symmetry codes: (i) $-x+1, y+1/2, -z+1/2$; (ii) $-x-1, -y+1, -z+1$; (iii) $x+1, -y+1/2, z-1/2$.