

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

(E)-2,3-Bis[(E)-benzylideneamino]but-2-enedinitrile

Matthew P. Akerman* and Sarah J. Maher

School of Chemistry and Physics, University of KwaZulu-Natal, Private Bag X01, Scottsville, Pietermaritzburg 3209, South Africa
Correspondence e-mail: akermanm@ukzn.ac.za


Received 18 January 2012; accepted 7 February 2012

Key indicators: single-crystal X-ray study; $T = 295$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.045; wR factor = 0.117; data-to-parameter ratio = 15.0.

The asymmetric unit of the title compound, $\text{C}_{18}\text{H}_{12}\text{N}_4$, consists of a half-molecule, where the two halves of the molecule are related by inversion symmetry. The molecule is effectively planar, with the largest deviation from the 22-atom mean plane, measuring 0.024 (2) Å, exhibited by the *ortho*-C atom of the phenyl ring. The crystal structure exhibits π -stacking, with an interplanar spacing of 3.431 (3) Å.

Related literature

For applications of the title molecule as a semiconductor, see: Tanaka *et al.* (2009). For applications of the title compound and its various derivatives as a dye, see: Neumer (1977); Begland (1976). For the crystal structures of three di(azomethine) dyes with various substituents at the *para* position of the benzene ring of the title compound, see: Matsumoto *et al.* (2004). For a study of the nonlinear optics applications of both the title compound and the mono-condensation product, see: Das *et al.* (2001). For a review of the chemistry and reactions of the diaminomaleonitrile, see: Al-Azmi *et al.* (2003).



Experimental

Crystal data

$\text{C}_{18}\text{H}_{12}\text{N}_4$	$\gamma = 102.97$ (5)°
$M_r = 284.32$	$V = 380.6$ (4) Å ³
Triclinic, $P\bar{1}$	$Z = 1$
$a = 6.389$ (4) Å	Mo $K\alpha$ radiation
$b = 7.608$ (5) Å	$\mu = 0.08$ mm ⁻¹
$c = 8.311$ (5) Å	$T = 295$ K
$\alpha = 103.96$ (5)°	$0.50 \times 0.20 \times 0.10$ mm
$\beta = 91.67$ (5)°	

Data collection

Oxford Diffraction Xcalibur 2 CCD diffractometer	2824 measured reflections
Absorption correction: multi-scan (Blessing, 1995)	1498 independent reflections
$T_{\min} = 0.963$, $T_{\max} = 0.992$	964 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.025$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.045$	100 parameters
$wR(F^2) = 0.117$	H-atom parameters constrained
$S = 0.89$	$\Delta\rho_{\text{max}} = 0.11$ e Å ⁻³
1498 reflections	$\Delta\rho_{\text{min}} = -0.21$ e Å ⁻³

Data collection: *CrysAlis CCD* (Oxford Diffraction, 2008); cell refinement: *CrysAlis CCD*; data reduction: *CrysAlis RED* (Oxford Diffraction, 2008); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *publCIF* (Westrip, 2010).

We would like to thank the University of KwaZulu-Natal for providing the funding and research facilities.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: EZ2280).

References

- Al-Azmi, A., Elassar, A.-Z. A. & Booth, B. L. (2003). *Tetrahedron*, **59**, 2749–2763.
- Begland, R. W. (1976). US Patent Application 3962220A19760608.
- Blessing, R. H. (1995). *Acta Cryst.* **A51**, 33–38.
- Das, P. K., Krishnan, A., Pal, S. K., Nandakumar, P. & Samuelson, A. G. (2001). *Chem. Phys.* **265**, 313–322.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Matsumoto, S., Shirai, K., Kobayashi, K., Wada, T. & Shiro, M. (2004). *Z. Kristallogr.* **219**, 239–243.
- Neumer, J. F. (1977). US Patent Application 4002616A19770111.
- Oxford Diffraction (2008). *CrysAlis CCD* and *CrysAlis RED*. Oxford Diffraction Ltd, Yarnton, Oxfordshire, England.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Tanaka, T., Matsumoto, S., Aoyama, T. & Wada, T. (2009). *Jpn Kokai Tokkyo Koho*, JP Patent 2009283523 A 20091203.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2012). E68, o688 [doi:10.1107/S1600536812005363]

(E)-2,3-Bis[(E)-benzylideneamino]but-2-enedinitrile

Matthew P. Akerman and Sarah J. Maher

S1. Comment

Di(azomethine) compounds derived from diaminomaleonitrile (DAMN) have been extensively studied for their application as dyes, semiconductors and in non-linear optics (Tanaka *et al.*, 2009; Neumer, 1977; Begland, 1976; Matsumoto *et al.*, 2004 and Das *et al.*, 2001). The title compound is interesting in that although it is orientated in a *trans* configuration about the ethylene group of the di(azomethine) linkage the DAMN synthon is exclusively *cis* about this bond. It is proposed that at elevated temperatures during the chemical synthesis an equilibrium is established between the *cis* and *trans* isomers of DAMN in solution. Although it has been shown that the *cis* isomer of DAMN is lower in energy than the *trans* isomer (Al-Azmi *et al.*, 2003) it is likely that non-bonded repulsion between the hydrogen atoms in the *ortho* positions of (1) would result in the *trans* isomer of (1) having a lower energy than the *cis* isomer. The double condensation reaction required to form (1) is therefore more likely to take place with the *trans* isomer of DAMN in solution; the *trans* isomer of (1) would thus be the major product.


Compound (1) crystallized in the triclinic space group $P\bar{1}$ with a half molecule in the asymmetric unit. The two halves of the molecule are related by inversion symmetry with an inversion centre at the midpoint of the C=C double bond of the di(azomethine) linkage unit. The molecule is effectively planar with the largest deviations from the 22-atom mean plane exhibited by the *ortho* C-atoms C3 and C5, measuring 0.022 (2) and 0.024 (2) Å, respectively. The structure shows that adjacent molecules are parallel, but are in a staggered configuration. There appear to be π - π interactions between adjacent molecules with an interplanar spacing of 3.431 (3) Å. The structure shows no genuine hydrogen bonding as all interactions are longer than the sum of their van der Waals radii. The lack of meaningful hydrogen bonds is likely due to a lack of good H-bond donors, as both the cyanide groups and the imine nitrogen atoms are potentially good H-bond acceptors.

S2. Experimental

Benzaldehyde (0.500 g, 4.7 mmol), *cis*-diaminomaleonitrile (0.255 g, 2.4 mmol) and a catalytic amount of piperidine were dissolved in dry toluene (100 ml) and the solution brought to reflux for four hours. Water was continuously removed from the reaction *via* a Dean and Stark apparatus. The solvent was removed by rotary evaporation under reduced pressure and the resulting solid dissolved in a minimum of dichloromethane. The desired product was obtained by column chromatography on silica gel, using dichloromethane as the eluent. Single crystals were grown by slow evaporation of the eluent. Yield (0.413 g, 62%). $^1\text{H NMR}$ (500 MHz, CD_3CN): 7.52 (t, 4H, *m*-phenyl), 7.60 (t, 2H, *p*-phenyl), 8.02 (d, 4H, *o*-phenyl), 8.75 (s, 2H, imine).

S3. Refinement

The positions of all hydrogen atoms were calculated using the standard riding model of *SHELXL97* with C—H(aromatic) distances of 0.93 Å and $U_{\text{iso}} = 1.2 U_{\text{eq}}$.

**Figure 1**

Labelled thermal ellipsoid plot of (1) (50% probability surfaces). Hydrogen atoms have been rendered as spheres of arbitrary radius.

(E)-2,3-Bis[(E)-benzylideneamino]but-2-enedinitrile*Crystal data* $C_{18}H_{12}N_4$ $M_r = 284.32$ Triclinic, $P\bar{1}$

Hall symbol: -P 1

 $a = 6.389$ (4) Å $b = 7.608$ (5) Å $c = 8.311$ (5) Å $\alpha = 103.96$ (5)° $\beta = 91.67$ (5)° $\gamma = 102.97$ (5)° $V = 380.6$ (4) Å³ $Z = 1$ $F(000) = 148$ $D_x = 1.241$ Mg m⁻³Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 2824 reflections

 $\theta = 3.3$ – 26° $\mu = 0.08$ mm⁻¹ $T = 295$ K

Needle, yellow

 $0.50 \times 0.20 \times 0.10$ mm*Data collection*Oxford Diffraction Xcalibur 2 CCD
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 ω scans at fixed θ anglesAbsorption correction: multi-scan
(Blessing, 1995) $T_{\min} = 0.963$, $T_{\max} = 0.992$

2824 measured reflections

1498 independent reflections

964 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.025$ $\theta_{\max} = 26.1^\circ$, $\theta_{\min} = 3.3^\circ$ $h = -6 \rightarrow 7$ $k = -9 \rightarrow 9$ $l = -9 \rightarrow 10$ *Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.045$ $wR(F^2) = 0.117$ $S = 0.89$

1498 reflections

100 parameters

0 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0717P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} < 0.001$ $\Delta\rho_{\max} = 0.11$ e Å⁻³ $\Delta\rho_{\min} = -0.21$ e Å⁻³

Special details

Experimental. ¹H NMR (500 MHz, CD₃CN): 7.52 (t, 4H, m-phenyl), 7.60 (t, 2H, p-phenyl), 8.02 (d, 4H, o-phenyl), 8.75 (s, 2H, imine).

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	-0.8068 (3)	0.8047 (3)	0.9287 (2)	0.0708 (5)
H1	-0.9171	0.8241	0.9965	0.085*
C2	-0.7329 (3)	0.9254 (3)	0.8351 (2)	0.0664 (5)
H2	-0.7932	1.0265	0.8383	0.080*
C3	-0.5686 (2)	0.8984 (2)	0.7353 (2)	0.0547 (4)
H3	-0.5176	0.9818	0.6720	0.066*
C4	-0.7201 (3)	0.6539 (3)	0.9241 (2)	0.0670 (5)
H4	-0.7715	0.5720	0.9887	0.080*
C5	-0.5573 (2)	0.6243 (2)	0.82398 (19)	0.0544 (4)
H5	-0.4998	0.5215	0.8202	0.065*
C6	-0.4787 (2)	0.74650 (19)	0.72903 (16)	0.0436 (4)
C7	-0.3065 (2)	0.72118 (19)	0.62138 (17)	0.0451 (4)
H7	-0.2608	0.8070	0.5592	0.054*
C8	0.0179 (2)	0.7027 (2)	0.40742 (18)	0.0469 (4)
C9	-0.0504 (2)	0.57010 (17)	0.50538 (16)	0.0399 (3)
N1	0.0604 (2)	0.81235 (19)	0.33473 (18)	0.0687 (5)
N2	-0.21601 (17)	0.58581 (15)	0.60912 (13)	0.0412 (3)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0504 (10)	0.1120 (15)	0.0524 (10)	0.0354 (10)	0.0149 (8)	0.0092 (10)
C2	0.0622 (11)	0.0768 (11)	0.0636 (11)	0.0404 (9)	0.0068 (9)	0.0018 (9)
C3	0.0541 (9)	0.0575 (9)	0.0552 (9)	0.0230 (8)	0.0095 (7)	0.0101 (7)
C4	0.0553 (10)	0.0953 (13)	0.0575 (10)	0.0227 (10)	0.0167 (8)	0.0273 (10)
C5	0.0481 (9)	0.0645 (9)	0.0553 (9)	0.0207 (8)	0.0091 (7)	0.0168 (8)
C6	0.0369 (7)	0.0502 (8)	0.0406 (8)	0.0138 (6)	0.0035 (6)	0.0026 (6)
C7	0.0439 (8)	0.0442 (8)	0.0489 (8)	0.0131 (7)	0.0105 (7)	0.0115 (7)
C8	0.0472 (8)	0.0461 (8)	0.0510 (8)	0.0166 (6)	0.0142 (7)	0.0128 (7)
C9	0.0363 (7)	0.0415 (7)	0.0415 (7)	0.0085 (6)	0.0073 (6)	0.0101 (6)
N1	0.0862 (11)	0.0602 (8)	0.0735 (10)	0.0251 (8)	0.0335 (8)	0.0327 (8)
N2	0.0375 (6)	0.0438 (7)	0.0440 (7)	0.0133 (5)	0.0101 (5)	0.0102 (5)

Geometric parameters (Å, °)

C1—C2	1.358 (2)	C5—C6	1.381 (2)
C1—C4	1.375 (2)	C5—H5	0.9300
C1—H1	0.9300	C6—C7	1.4551 (19)
C2—C3	1.377 (2)	C7—N2	1.2751 (17)
C2—H2	0.9300	C7—H7	0.9300
C3—C6	1.392 (2)	C8—N1	1.1340 (17)
C3—H3	0.9300	C8—C9	1.4456 (19)
C4—C5	1.375 (2)	C9—C9 ⁱ	1.352 (2)
C4—H4	0.9300	C9—N2	1.3918 (17)
C2—C1—C4	120.52 (15)	C4—C5—H5	119.9
C2—C1—H1	119.7	C6—C5—H5	119.9
C4—C1—H1	119.7	C5—C6—C3	118.88 (13)
C1—C2—C3	120.11 (15)	C5—C6—C7	122.30 (13)
C1—C2—H2	119.9	C3—C6—C7	118.82 (14)
C3—C2—H2	119.9	N2—C7—C6	122.49 (13)
C2—C3—C6	120.20 (16)	N2—C7—H7	118.8
C2—C3—H3	119.9	C6—C7—H7	118.8
C6—C3—H3	119.9	N1—C8—C9	175.27 (15)
C1—C4—C5	120.00 (17)	C9 ⁱ —C9—N2	120.38 (15)
C1—C4—H4	120.0	C9 ⁱ —C9—C8	118.58 (14)
C5—C4—H4	120.0	N2—C9—C8	121.04 (11)
C4—C5—C6	120.29 (15)	C7—N2—C9	119.86 (12)
C4—C1—C2—C3	-0.5 (3)	C2—C3—C6—C7	179.53 (14)
C1—C2—C3—C6	0.5 (3)	C5—C6—C7—N2	-0.7 (2)
C2—C1—C4—C5	-0.1 (3)	C3—C6—C7—N2	179.89 (12)
C1—C4—C5—C6	0.7 (3)	C6—C7—N2—C9	-179.29 (12)
C4—C5—C6—C3	-0.6 (2)	C9 ⁱ —C9—N2—C7	178.32 (15)
C4—C5—C6—C7	179.93 (14)	C8—C9—N2—C7	-1.5 (2)
C2—C3—C6—C5	0.1 (2)		

Symmetry code: (i) $-x, -y+1, -z+1$.