

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

3-(1*H*-Imidazol-1-yl)-1-phenylpropan-1-ol

Hoong-Kun Fun,^{a*} Ching Kheng Quah,^{a§} Mohamed I. Attia,^b Hatem A. Abdel-Aziz^b and Khalid A. Al-Rashood^b

^aX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, and ^bDepartment of Pharmaceutical Chemistry, College of Pharmacy, King Saud University, Riyadh 11451, Saudi Arabia

Correspondence e-mail: hkfun@usm.my


Received 31 January 2012; accepted 1 February 2012

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.001$ Å; R factor = 0.043; wR factor = 0.125; data-to-parameter ratio = 27.0.

In the title compound, $\text{C}_{12}\text{H}_{14}\text{N}_2\text{O}$, the imidazole ring forms a dihedral angle of 66.73 (5)° with the phenyl ring. In the crystal, molecules are linked *via* $\text{O}-\text{H}\cdots\text{N}$ and $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds into sheets lying parallel to (100). The crystal structure is further consolidated by $\text{C}-\text{H}\cdots\pi$ interactions.

Related literature

For general background to and the pharmacological activities of the title compound, see: Latge (1999); Steenberg & Casadevall (2000); Pacetti & Gelone (2003); Spellberg *et al.* (2006). For the preparation of the title compound, see: Aboul-Enein *et al.* (2011). For the stability of the temperature controller used for the data collection, see: Cosier & Glazer (1986).



Experimental

Crystal data

| | |
|--|-----------------------------------|
| $\text{C}_{12}\text{H}_{14}\text{N}_2\text{O}$ | $V = 1043.17$ (10) Å ³ |
| $M_r = 202.25$ | $Z = 4$ |
| Monoclinic, $P2_1/c$ | Mo $K\alpha$ radiation |
| $a = 9.0352$ (5) Å | $\mu = 0.08$ mm ⁻¹ |
| $b = 11.8521$ (7) Å | $T = 100$ K |
| $c = 10.3462$ (6) Å | $0.34 \times 0.26 \times 0.19$ mm |
| $\beta = 109.688$ (1)° | |

Data collection

| | |
|--|--|
| Bruker SMART APEXII DUO | 14426 measured reflections |
| CCD diffractometer | 3777 independent reflections |
| Absorption correction: multi-scan (SADABS; Bruker, 2009) | 3245 reflections with $I > 2\sigma(I)$ |
| $T_{\min} = 0.972$, $T_{\max} = 0.985$ | $R_{\text{int}} = 0.022$ |

Refinement

| | |
|---------------------------------|--|
| $R[F^2 > 2\sigma(F^2)] = 0.043$ | H atoms treated by a mixture of independent and constrained refinement |
| $wR(F^2) = 0.125$ | $\Delta\rho_{\text{max}} = 0.67$ e Å ⁻³ |
| $S = 1.05$ | $\Delta\rho_{\text{min}} = -0.26$ e Å ⁻³ |
| 3777 reflections | |
| 140 parameters | |

Table 1

Hydrogen-bond geometry (Å, °).

Cg1 is the centroid of the $\text{C1}-\text{C6}$ phenyl ring.

| $D-\text{H}\cdots A$ | $D-\text{H}$ | $\text{H}\cdots A$ | $D\cdots A$ | $D-\text{H}\cdots A$ |
|--|--------------|--------------------|-------------|----------------------|
| $\text{O1}-\text{H1O1}\cdots\text{N2}^i$ | 0.964 (19) | 1.89 (2) | 2.8432 (12) | 171.7 (17) |
| $\text{C1}-\text{H1B}\cdots\text{O1}^{ii}$ | 0.95 | 2.48 | 3.4188 (12) | 172 |
| $\text{C10}-\text{H10A}\cdots\text{Cg1}^{iii}$ | 0.95 | 2.68 | 3.4778 (12) | 142 |
| $\text{C12}-\text{H12A}\cdots\text{Cg1}^{iv}$ | 0.95 | 2.69 | 3.5373 (11) | 149 |

Symmetry codes: (i) $-x + 1, y - \frac{1}{2}, -z + \frac{1}{2}$; (ii) $-x + 1, -y, -z + 1$; (iii) $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$; (iv) $-x, y + \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: APEX2 (Bruker, 2009); cell refinement: SAINT (Bruker, 2009); data reduction: SAINT; program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL and PLATON (Spek, 2009).

The authors thank Universiti Sains Malaysia (USM) for a Research University Grant (No. 1001/PFIZIK/811160) and the Deanship of Scientific Research and the Research Center of the College of Pharmacy, King Saud University, for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB6620).

References

- Aboul-Enein, M. N., El-Azzouny, A. A., Attia, M. I., Saleh, O. A. & Kansoh, A. L. (2011). *Arch. Pharm.* **344**, 794–801.
- Bruker (2009). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cosier, J. & Glazer, A. M. (1986). *J. Appl. Cryst.* **19**, 105–107.
- Latge, J. P. (1999). *Clin. Microbiol. Rev.* **12**, 310–350.
- Pacetti, S. A. & Gelone, S. P. (2003). *Ann Pharmacother.* **37**, 90–98.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Spellberg, B. J., Filler, S. G. & Edwards, J. E. (2006). *Clin. Infect. Dis.* **42**, 244–251.
- Steenberg, J. N. & Casadevall, A. (2000). *J. Clin. Microbiol.* **38**, 1974–1976.

* Thomson Reuters ResearcherID: A-3561-2009.

§ Thomson Reuters ResearcherID: A-5525-2009.

supporting information

Acta Cryst. (2012). E68, o628 [doi:10.1107/S1600536812004254]

3-(1*H*-Imidazol-1-yl)-1-phenylpropan-1-ol

Hoong-Kun Fun, Ching Kheng Quah, Mohamed I. Attia, Hatem A. Abdel-Aziz and Khalid A. Al-Rashood

S1. Comment

Over the past three decades, the incidence of both community-acquired and nosocomial invasive fungal infections has increased substantially. Clinically, candidosis, aspergillosis and cryptococcosis have been identified as three major opportunistic pathogens in the etiology of fungal infections in immune compromised patients (Latge, 1999; Steenberg & Casadevall, 2000). *Candida albicans* accounts for the majority of invasive and superficial *Candida* infections (Pacetti & Gelone, 2003; Spellberg *et al.*, 2006). Accordingly, the need for new antifungal drugs has prompted intensive research worldwide. The azole antifungal drugs constitute one of the major classes which are characterized by having the azole pharmacophoric moiety embedded in their structures. The title molecule exhibited anti-*Candida* activity (MIC = 10 $\mu\text{g/ml}$) and can serve as a prototypic molecule for subsequent molecular modifications.

In the title compound, Fig. 1, the imidazole ring (N1/N2/C10–C12, maximum deviation of 0.002 (1) Å at atoms N1 and C12) forms a dihedral angle of 66.73 (5)° with the phenyl ring (C1–C6).

In the crystal structure, Fig. 2, molecules are linked *via* intermolecular O1—H1O1 \cdots N2 and C1—H1B \cdots O1 hydrogen bonds (Table 1) into two-dimensional networks parallel to (100). The crystal structure is further consolidated by C10—H10A \cdots Cg1ⁱⁱⁱ and C12—H12A \cdots Cg1^{iv} (Table 1) interactions, where Cg1 is the centroid of C1–C6 phenyl ring.

S2. Experimental

NaBH₄ (4.9 g, 0.13 mol) was added portion-wise to an ice cooled, stirred solution of 3-(1*H*-imidazol-1-yl)-1-phenylpropan-1-one (8.6 g, 0.043 mol) (Aboul-Enein *et al.*, 2011) in methanol (100 ml). The mixture was stirred overnight at ambient temperature followed by evaporation of methanol under vacuum. The residue was dissolved in ethyl acetate (150 ml) and washed with water (3 \times 50 ml). The organic layer was separated, dried (Na₂SO₄) and evaporated under reduced pressure. The residue was recrystallized from ethanol to give the title compound as colourless blocks. M.p. = 380–382 K.

S3. Refinement

Atom H1O1 was located in a difference Fourier map and refined freely with O1—H1O1 = 0.97 (2) Å. The remaining H atoms were positioned geometrically and refined using a riding model with C—H = 0.95 or 0.99 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.




Figure 1

The molecular structure of the title compound showing 50% probability displacement ellipsoids for non-H atoms.




Figure 2

The crystal structure of the title compound, viewed along the *c* axis. H atoms not involved in hydrogen bonds (dashed lines) have been omitted for clarity.

3-(1*H*-Imidazol-1-yl)-1-phenylpropan-1-ol

Crystal data

$C_{12}H_{14}N_2O$

$M_r = 202.25$

Monoclinic, $P2_1/c$

Hall symbol: $-P 2_1/c$

$a = 9.0352 (5) \text{ \AA}$

$b = 11.8521 (7) \text{ \AA}$

$c = 10.3462 (6) \text{ \AA}$

$\beta = 109.688 (1)^\circ$

$V = 1043.17 (10) \text{ \AA}^3$

$Z = 4$

$F(000) = 432$

$D_x = 1.288 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 6547 reflections

$\theta = 3.0\text{--}32.6^\circ$

$\mu = 0.08 \text{ mm}^{-1}$

$T = 100 \text{ K}$

Block, colourless

$0.34 \times 0.26 \times 0.19 \text{ mm}$

Data collection

Bruker SMART APEXII DUO CCD
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 φ and ω scans
Absorption correction: multi-scan
(*SADABS*; Bruker, 2009)
 $T_{\min} = 0.972$, $T_{\max} = 0.985$

14426 measured reflections
3777 independent reflections
3245 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.022$
 $\theta_{\text{max}} = 32.6^\circ$, $\theta_{\text{min}} = 2.7^\circ$
 $h = -13 \rightarrow 12$
 $k = -17 \rightarrow 17$
 $l = -14 \rightarrow 15$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.043$
 $wR(F^2) = 0.125$
 $S = 1.05$
3777 reflections
140 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H atoms treated by a mixture of independent
and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0644P)^2 + 0.3608P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.001$
 $\Delta\rho_{\text{max}} = 0.67 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.26 \text{ e } \text{\AA}^{-3}$

Special details

Experimental. The crystal was placed in the cold stream of an Oxford Cryosystems Cobra open-flow nitrogen cryostat (Cosier & Glazer, 1986) operating at 100.0 (1) K.

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

| | <i>x</i> | <i>y</i> | <i>z</i> | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|-----|--------------|--------------|--------------|----------------------------------|
| O1 | 0.48286 (8) | 0.08562 (6) | 0.36457 (8) | 0.02067 (16) |
| N1 | 0.25481 (8) | 0.31589 (7) | 0.35682 (8) | 0.01420 (15) |
| N2 | 0.32358 (10) | 0.45590 (7) | 0.24613 (9) | 0.02066 (17) |
| C1 | 0.31643 (10) | -0.13411 (8) | 0.29554 (9) | 0.01560 (16) |
| H1B | 0.3689 | -0.1285 | 0.3918 | 0.019* |
| C2 | 0.27877 (11) | -0.23971 (8) | 0.23471 (10) | 0.01808 (18) |
| H2B | 0.3073 | -0.3059 | 0.2892 | 0.022* |
| C3 | 0.19909 (12) | -0.24865 (8) | 0.09354 (10) | 0.01975 (18) |
| H3A | 0.1735 | -0.3208 | 0.0519 | 0.024* |
| C4 | 0.15757 (11) | -0.15146 (9) | 0.01470 (9) | 0.01935 (18) |
| H4A | 0.1013 | -0.1570 | -0.0809 | 0.023* |
| C5 | 0.19809 (11) | -0.04577 (8) | 0.07534 (9) | 0.01624 (17) |
| H5A | 0.1713 | 0.0203 | 0.0203 | 0.019* |
| C6 | 0.27760 (10) | -0.03610 (8) | 0.21592 (9) | 0.01380 (16) |

| | | | | |
|------|--------------|-------------|--------------|--------------|
| C7 | 0.32242 (10) | 0.07925 (8) | 0.28116 (9) | 0.01549 (16) |
| H7A | 0.3017 | 0.1370 | 0.2066 | 0.019* |
| C8 | 0.22380 (10) | 0.10796 (8) | 0.37099 (9) | 0.01587 (16) |
| H8A | 0.1118 | 0.1116 | 0.3124 | 0.019* |
| H8B | 0.2350 | 0.0464 | 0.4383 | 0.019* |
| C9 | 0.26905 (11) | 0.21929 (8) | 0.44836 (9) | 0.01656 (17) |
| H9A | 0.2007 | 0.2317 | 0.5044 | 0.020* |
| H9B | 0.3789 | 0.2142 | 0.5117 | 0.020* |
| C10 | 0.37482 (11) | 0.37408 (8) | 0.33713 (10) | 0.01860 (18) |
| H10A | 0.4828 | 0.3579 | 0.3833 | 0.022* |
| C11 | 0.16177 (11) | 0.44949 (8) | 0.20509 (10) | 0.01811 (18) |
| H11A | 0.0914 | 0.4980 | 0.1397 | 0.022* |
| C12 | 0.11734 (10) | 0.36343 (8) | 0.27205 (9) | 0.01602 (17) |
| H12A | 0.0131 | 0.3411 | 0.2621 | 0.019* |
| H10I | 0.543 (2) | 0.0448 (17) | 0.3186 (19) | 0.048 (5)* |

Atomic displacement parameters (Å²)

| | U^{11} | U^{22} | U^{33} | U^{12} | U^{13} | U^{23} |
|-----|------------|------------|------------|-------------|------------|-------------|
| O1 | 0.0152 (3) | 0.0220 (4) | 0.0251 (3) | -0.0023 (2) | 0.0071 (3) | -0.0072 (3) |
| N1 | 0.0142 (3) | 0.0116 (3) | 0.0184 (3) | 0.0000 (2) | 0.0076 (3) | -0.0010 (2) |
| N2 | 0.0182 (3) | 0.0179 (4) | 0.0290 (4) | 0.0007 (3) | 0.0120 (3) | 0.0036 (3) |
| C1 | 0.0146 (3) | 0.0167 (4) | 0.0159 (3) | 0.0024 (3) | 0.0057 (3) | 0.0012 (3) |
| C2 | 0.0204 (4) | 0.0137 (4) | 0.0232 (4) | 0.0028 (3) | 0.0114 (3) | 0.0018 (3) |
| C3 | 0.0227 (4) | 0.0154 (4) | 0.0244 (4) | -0.0015 (3) | 0.0121 (3) | -0.0055 (3) |
| C4 | 0.0207 (4) | 0.0212 (4) | 0.0160 (4) | -0.0009 (3) | 0.0060 (3) | -0.0035 (3) |
| C5 | 0.0174 (4) | 0.0158 (4) | 0.0159 (4) | 0.0010 (3) | 0.0062 (3) | 0.0007 (3) |
| C6 | 0.0125 (3) | 0.0141 (4) | 0.0160 (3) | -0.0003 (3) | 0.0063 (3) | -0.0015 (3) |
| C7 | 0.0155 (3) | 0.0142 (4) | 0.0180 (4) | -0.0009 (3) | 0.0072 (3) | -0.0011 (3) |
| C8 | 0.0176 (4) | 0.0127 (4) | 0.0205 (4) | -0.0004 (3) | 0.0106 (3) | -0.0001 (3) |
| C9 | 0.0207 (4) | 0.0135 (4) | 0.0176 (4) | 0.0012 (3) | 0.0091 (3) | 0.0003 (3) |
| C10 | 0.0141 (4) | 0.0174 (4) | 0.0262 (4) | -0.0003 (3) | 0.0092 (3) | 0.0014 (3) |
| C11 | 0.0176 (4) | 0.0168 (4) | 0.0219 (4) | 0.0032 (3) | 0.0092 (3) | 0.0020 (3) |
| C12 | 0.0136 (3) | 0.0160 (4) | 0.0195 (4) | 0.0003 (3) | 0.0069 (3) | -0.0012 (3) |

Geometric parameters (Å, °)

| | | | |
|---------|-------------|--------|-------------|
| O1—C7 | 1.4173 (11) | C4—H4A | 0.9500 |
| O1—H10I | 0.97 (2) | C5—C6 | 1.3931 (12) |
| N1—C10 | 1.3571 (11) | C5—H5A | 0.9500 |
| N1—C12 | 1.3767 (11) | C6—C7 | 1.5180 (12) |
| N1—C9 | 1.4636 (12) | C7—C8 | 1.5269 (12) |
| N2—C10 | 1.3218 (13) | C7—H7A | 1.0000 |
| N2—C11 | 1.3802 (12) | C8—C9 | 1.5260 (13) |
| C1—C2 | 1.3907 (13) | C8—H8A | 0.9900 |
| C1—C6 | 1.3986 (12) | C8—H8B | 0.9900 |
| C1—H1B | 0.9500 | C9—H9A | 0.9900 |
| C2—C3 | 1.3979 (14) | C9—H9B | 0.9900 |

| | | | |
|-------------|-------------|----------------|--------------|
| C2—H2B | 0.9500 | C10—H10A | 0.9500 |
| C3—C4 | 1.3881 (14) | C11—C12 | 1.3671 (13) |
| C3—H3A | 0.9500 | C11—H11A | 0.9500 |
| C4—C5 | 1.3936 (13) | C12—H12A | 0.9500 |
| | | | |
| C7—O1—H1O1 | 107.8 (11) | C6—C7—C8 | 110.39 (7) |
| C10—N1—C12 | 106.94 (8) | O1—C7—H7A | 108.7 |
| C10—N1—C9 | 126.45 (8) | C6—C7—H7A | 108.7 |
| C12—N1—C9 | 126.60 (7) | C8—C7—H7A | 108.7 |
| C10—N2—C11 | 105.03 (8) | C9—C8—C7 | 113.82 (7) |
| C2—C1—C6 | 120.42 (8) | C9—C8—H8A | 108.8 |
| C2—C1—H1B | 119.8 | C7—C8—H8A | 108.8 |
| C6—C1—H1B | 119.8 | C9—C8—H8B | 108.8 |
| C1—C2—C3 | 120.16 (9) | C7—C8—H8B | 108.8 |
| C1—C2—H2B | 119.9 | H8A—C8—H8B | 107.7 |
| C3—C2—H2B | 119.9 | N1—C9—C8 | 112.79 (7) |
| C4—C3—C2 | 119.55 (9) | N1—C9—H9A | 109.0 |
| C4—C3—H3A | 120.2 | C8—C9—H9A | 109.0 |
| C2—C3—H3A | 120.2 | N1—C9—H9B | 109.0 |
| C3—C4—C5 | 120.22 (8) | C8—C9—H9B | 109.0 |
| C3—C4—H4A | 119.9 | H9A—C9—H9B | 107.8 |
| C5—C4—H4A | 119.9 | N2—C10—N1 | 111.96 (8) |
| C6—C5—C4 | 120.59 (8) | N2—C10—H10A | 124.0 |
| C6—C5—H5A | 119.7 | N1—C10—H10A | 124.0 |
| C4—C5—H5A | 119.7 | C12—C11—N2 | 110.28 (8) |
| C5—C6—C1 | 119.04 (8) | C12—C11—H11A | 124.9 |
| C5—C6—C7 | 120.32 (8) | N2—C11—H11A | 124.9 |
| C1—C6—C7 | 120.64 (8) | C11—C12—N1 | 105.80 (8) |
| O1—C7—C6 | 112.55 (7) | C11—C12—H12A | 127.1 |
| O1—C7—C8 | 107.72 (7) | N1—C12—H12A | 127.1 |
| | | | |
| C6—C1—C2—C3 | -1.16 (13) | O1—C7—C8—C9 | 52.81 (10) |
| C1—C2—C3—C4 | -0.15 (14) | C6—C7—C8—C9 | 176.07 (7) |
| C2—C3—C4—C5 | 1.41 (14) | C10—N1—C9—C8 | -107.01 (10) |
| C3—C4—C5—C6 | -1.37 (14) | C12—N1—C9—C8 | 71.41 (11) |
| C4—C5—C6—C1 | 0.06 (13) | C7—C8—C9—N1 | 59.04 (10) |
| C4—C5—C6—C7 | 179.56 (8) | C11—N2—C10—N1 | -0.15 (11) |
| C2—C1—C6—C5 | 1.20 (12) | C12—N1—C10—N2 | 0.31 (11) |
| C2—C1—C6—C7 | -178.30 (8) | C9—N1—C10—N2 | 178.99 (8) |
| C5—C6—C7—O1 | -129.62 (8) | C10—N2—C11—C12 | -0.07 (11) |
| C1—C6—C7—O1 | 49.88 (10) | N2—C11—C12—N1 | 0.25 (11) |
| C5—C6—C7—C8 | 109.98 (9) | C10—N1—C12—C11 | -0.33 (10) |
| C1—C6—C7—C8 | -70.52 (10) | C9—N1—C12—C11 | -179.01 (8) |

Hydrogen-bond geometry (Å, °)

Cg1 is the centroid of the C1–C6 phenyl ring.

| <i>D</i> —H··· <i>A</i> | <i>D</i> —H | H··· <i>A</i> | <i>D</i> ··· <i>A</i> | <i>D</i> —H··· <i>A</i> |
|-------------------------------|-------------|---------------|-----------------------|-------------------------|
| O1—H1O1···N2 ⁱ | 0.964 (19) | 1.89 (2) | 2.8432 (12) | 171.7 (17) |
| C1—H1B···O1 ⁱⁱ | 0.95 | 2.48 | 3.4188 (12) | 172 |
| C10—H10A···Cg1 ⁱⁱⁱ | 0.95 | 2.68 | 3.4778 (12) | 142 |
| C12—H12A···Cg1 ^{iv} | 0.95 | 2.69 | 3.5373 (11) | 149 |

Symmetry codes: (i) $-x+1, y-1/2, -z+1/2$; (ii) $-x+1, -y, -z+1$; (iii) $-x+1, y+1/2, -z+1/2$; (iv) $-x, y+1/2, -z+1/2$.