

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

(Z)-4-(2-Hydroxybenzylidene)-1-methyl-2-phenyl-1H-imidazol-5(4H)-one

Ming-Jen Chang, Hsing-Yang Tsai, Tzu-Chien Fang, Ming-Hui Luo and Kew-Yu Chen*

Department of Chemical Engineering, Feng Chia University, 40724 Taichung, Taiwan

Correspondence e-mail: kyuchen@fcu.edu.tw


Received 10 February 2012; accepted 22 February 2012

Key indicators: single-crystal X-ray study; $T = 150$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.037; wR factor = 0.101; data-to-parameter ratio = 12.3.

In the title compound, $\text{C}_{17}\text{H}_{14}\text{N}_2\text{O}_2$, the asymmetric unit comprises two molecules that are conformationally similar [the dihedral angles between the phenyl rings in each are $46.35(2)$ and $48.04(3)^\circ$], with the conformation stabilized by intramolecular $\text{O}-\text{H}\cdots\text{N}$ hydrogen bonds, which generate $S(7)$ rings. In the crystal, inversion-related molecules are linked by pairs of weak $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds, forming dimers with an $R_2^2(16)$ graph-set motif. Weak inter-ring $\pi-\pi$ stacking is observed in the structure, the shortest centroid-to-centroid distance being $3.7480(13)$ Å.

Related literature

For the spectroscopy and preparation of the title compound, see: Chuang *et al.* (2011). For the applications of proton-transfer dyes, see: Chen & Pang (2010); Gryko *et al.* (2010); Han *et al.* (2010); Helal *et al.* (2010); Ikeda *et al.* (2010); Ito *et al.* (2011); Lim *et al.* (2011); Lins *et al.* (2010); Maupin *et al.* (2011); Santos *et al.* (2011); Tang *et al.* (2011). For a related structure, see: Chen *et al.* (2007). For graph-set theory of hydrogen bonds, see: Bernstein *et al.* (1995).



Experimental

Crystal data

$\text{C}_{17}\text{H}_{14}\text{N}_2\text{O}_2$	$\gamma = 70.500(2)^\circ$
$M_r = 278.30$	$V = 1334.57(9) \text{ \AA}^3$
Triclinic, $P\bar{1}$	$Z = 4$
$a = 9.7843(4) \text{ \AA}$	Mo $K\alpha$ radiation
$b = 9.8972(3) \text{ \AA}$	$\mu = 0.09 \text{ mm}^{-1}$
$c = 15.4313(6) \text{ \AA}$	$T = 150 \text{ K}$
$\alpha = 72.086(2)^\circ$	$0.32 \times 0.28 \times 0.14 \text{ mm}$
$\beta = 79.301(2)^\circ$	

Data collection

Bruker SMART CCD diffractometer	20392 measured reflections
Absorption correction: multi-scan (SADABS; Bruker, 2001)	4699 independent reflections
$T_{\min} = 0.889$, $T_{\max} = 0.984$	2273 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.054$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.037$	382 parameters
$wR(F^2) = 0.101$	H-atom parameters constrained
$S = 0.80$	$\Delta\rho_{\text{max}} = 0.35 \text{ e \AA}^{-3}$
4699 reflections	$\Delta\rho_{\text{min}} = -0.36 \text{ e \AA}^{-3}$

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O}2-\text{H}2\cdots\text{N}2$	0.82	1.80	2.615(2)	176
$\text{O}4-\text{H}4\cdots\text{N}4$	0.82	1.79	2.612(3)	175
$\text{C}10-\text{H}10\cdots\text{O}1^i$	0.93	2.58	3.403(3)	148
$\text{C}30-\text{H}30\cdots\text{O}3^{\text{ii}}$	0.93	2.68	3.421(3)	137

Symmetry codes: (i) $-x + 1, -y, -z$; (ii) $-x + 1, -y + 1, -z$.

Data collection: SMART (Bruker, 2001); cell refinement: SAINT (Bruker, 2001); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 for Windows (Farrugia, 1997); software used to prepare material for publication: WinGX (Farrugia, 1999).

This work was supported by the National Science Council (grant No. NSC 99-2113-M-035-001-MY2) and Feng Chia University, Taiwan.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: ZS2183).

References

- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2001). SMART, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chen, K.-Y., Cheng, Y.-M., Lai, C.-H., Hsu, C.-C., Ho, M.-L., Lee, G.-H. & Chou, P.-T. (2007). *J. Am. Chem. Soc.* **129**, 4534–4535.
- Chen, W.-H. & Pang, Y. (2010). *Tetrahedron Lett.* **51**, 1914–1918.
- Chuang, W.-T., Hsieh, C.-C., Lai, C.-H., Lai, C.-H., Shih, C.-W., Chen, K.-Y., Hung, W.-Y., Hsu, Y.-H. & Chou, P.-T. (2011). *J. Org. Chem.* **76**, 8189–8202.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Gryko, D. T., Piechowska, J. & Galezowski, M. (2010). *J. Org. Chem.* **75**, 1297–1300.
- Han, D. Y., Kim, J. M., Kim, J., Jung, H. S., Lee, Y. H., Zhang, J. F. & Kim, J. S. (2010). *Tetrahedron Lett.* **51**, 1947–1951.

- Helal, A., Lee, S. H., Kim, S. H. & Kim, H.-S. (2010). *Tetrahedron Lett.* **51**, 3531–3535.
- Ikeda, S., Toganoh, M., Easwaramoorthi, S., Lim, J. M., Kim, D. & Furuta, H. (2010). *J. Org. Chem.* **75**, 8637–8649.
- Ito, Y., Amimoto, K. & Kawato, T. (2011). *Dyes Pigm.* **89**, 319–323.
- Lim, C.-K., Seo, J., Kim, S., Kwon, I. C., Ahn, C.-H. & Park, S. Y. (2011). *Dyes Pigm.* **90**, 284–289.
- Lins, G. O. W., Campo, L. F., Rodembusch, F. S. & Stefani, V. (2010). *Dyes Pigm.* **84**, 114–120.
- Maupin, C. M., Castillo, N., Taraphder, S., Tu, C., McKenna, R., Silverman, D. N. & Voth, G. A. (2011). *J. Am. Chem. Soc.* **133**, 6223–6234.
- Santos, R. C., Silva Faleiro, N. V., Campo, L. F., Scroferneker, M. L., Corbellini, V. A., Rodembusch, F. S. & Stefani, V. (2011). *Tetrahedron Lett.* **52**, 3048–3053.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Tang, K.-C., Chang, M.-J., Lin, T.-Y., Pan, H.-A., Fang, T.-C., Chen, K.-Y., Hung, W.-Y., Hsu, Y.-H. & Chou, P.-T. (2011). *J. Am. Chem. Soc.* **133**, 17738–17745.

supporting information

Acta Cryst. (2012). E68, o902–o903 [doi:10.1107/S1600536812007921]

(Z)-4-(2-Hydroxybenzylidene)-1-methyl-2-phenyl-1H-imidazol-5(4H)-one**Ming-Jen Chang, Hsing-Yang Tsai, Tzu-Chien Fang, Ming-Hui Luo and Kew-Yu Chen****S1. Comment**

The excited-state intramolecular proton transfer (ESIPT) reaction of the title compound has been investigated recently (Chuang *et al.*, 2011), which involves transfer of a hydroxy proton to the imine nitrogen through an intramolecular seven-membered ring hydrogen-bonding system. The proton-transfer dyes have found many important applications.

Prototypical examples are probes for solvation dynamics (Chen & Pang, 2010; Lins *et al.*) and biological environments (Lim *et al.*, 2011; Maupin *et al.*, 2011), near-infrared fluorescent dyes (Ikeda *et al.*, 2010), chemosensors (Han *et al.*, 2010; Helal *et al.*, 2010), photochromic materials (Ito *et al.*, 2011), fluorescence microscopy imaging (Santos *et al.*, 2011), and recent applications in the field of organic light-emitting devices (Gryko *et al.*, 2010; Tang *et al.*, 2011).

The molecular structure of the title compound, C₁₇H₁₄N₂O₂ is shown in Fig. 1. The asymmetric unit comprises two symmetry-independent molecules (*A* and *B*) which are conformationally similar [dihedral angles between the phenyl rings in each are 46.35 (2) and 48.04 (3)°]. The conformation is stabilized in each by intramolecular O—H⋯N hydrogen bonds which generate *S*(7) rings (Chen *et al.*, 2007). Present also are intramolecular C—H⋯O interactions between the methyl group and the ketone O-atom, generating *S*(5) rings (Table 1). In the crystal (Fig. 2), inversion-related molecules are linked by pairs of weak hydrogen bonds (Table 1), forming cyclic dimers with an *R*₂²(16) graph-set motif (Bernstein *et al.*, 1995). Weak π – π stacking is also observed in the crystal structure, the shortest centroid–centroid distance being 3.7480 (13) Å [symmetry code: *x*, *y* - 1, *z*].

S2. Experimental

The title compound was synthesized according to the literature procedure (Chuang *et al.*, 2011). Yellow needle-shaped crystals suitable for the crystallographic studies reported here were isolated over a period of six weeks by slow evaporation from a chloroform solution.

S3. Refinement

The C-bound H atoms were positioned geometrically (C—H = 0.93–0.96 Å) and allowed to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. The O-bound H atoms were positioned geometrically (O—H = 0.82 Å) and allowed to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$.




Figure 1

The molecular structure of the title compound, showing 50% probability displacement ellipsoids.




Figure 2

A section of the crystal packing of the title compound, viewed along the *b* axis. Green and blue dashed lines denote the intermolecular C10—H10 \cdots O1ⁱ and C30—H30 \cdots O3ⁱⁱ hydrogen bonds, respectively. For symmetry codes (i) and (ii), see Table 1.

(*Z*)-4-(2-Hydroxybenzylidene)-1-methyl-2-phenyl-1*H*-imidazol-5(4*H*)-one

Crystal data

C₁₇H₁₄N₂O₂

M_r = 278.30

Triclinic, *P*1

Hall symbol: -P 1

a = 9.7843 (4) Å

b = 9.8972 (3) Å

c = 15.4313 (6) Å

α = 72.086 (2)°

β = 79.301 (2)°

γ = 70.500 (2)°

V = 1334.57 (9) Å³

Z = 4

F(000) = 584

D_x = 1.385 Mg m⁻³

Mo *K*α radiation, λ = 0.71073 Å

Cell parameters from 3314 reflections

θ = 2.5–25.0°

μ = 0.09 mm⁻¹

$T = 150$ K $0.32 \times 0.28 \times 0.14$ mm
 Prism, yellow

Data collection

Bruker SMART CCD diffractometer	20392 measured reflections
Radiation source: fine-focus sealed tube	4699 independent reflections
Graphite monochromator	2273 reflections with $I > 2\sigma(I)$
φ and ω scans	$R_{\text{int}} = 0.054$
Absorption correction: multi-scan (SADABS; Bruker, 2001)	$\theta_{\text{max}} = 25.0^\circ$, $\theta_{\text{min}} = 1.4^\circ$
$T_{\text{min}} = 0.889$, $T_{\text{max}} = 0.984$	$h = -11 \rightarrow 11$
	$k = -11 \rightarrow 10$
	$l = -18 \rightarrow 18$

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.037$	$w = 1/[\sigma^2(F_o^2) + (0.049P)^2]$
$wR(F^2) = 0.101$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 0.80$	$(\Delta/\sigma)_{\text{max}} < 0.001$
4699 reflections	$\Delta\rho_{\text{max}} = 0.35 \text{ e } \text{\AA}^{-3}$
382 parameters	$\Delta\rho_{\text{min}} = -0.36 \text{ e } \text{\AA}^{-3}$
0 restraints	Extinction correction: <i>SHELXL97</i> (Sheldrick, 2008), $F_c^* = kFc[1 + 0.001x \text{Fc}^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
Primary atom site location: structure-invariant direct methods	Extinction coefficient: 0.037 (2)
Secondary atom site location: difference Fourier map	

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.13687 (16)	-0.12970 (16)	0.02374 (11)	0.0292 (4)
O2	0.24941 (16)	0.01412 (16)	0.32193 (9)	0.0331 (5)
H2	0.1962	-0.0125	0.2996	0.050*
O3	0.15388 (16)	0.37862 (16)	0.01672 (11)	0.0299 (4)
O4	0.23819 (17)	0.52664 (17)	0.32188 (10)	0.0410 (5)
H4	0.1961	0.4877	0.2999	0.061*
N1	-0.00550 (19)	-0.16641 (18)	0.16225 (12)	0.0229 (5)
N2	0.09021 (19)	-0.07230 (18)	0.24367 (12)	0.0219 (5)
N3	0.00423 (19)	0.34215 (18)	0.15218 (12)	0.0228 (5)
N4	0.10847 (19)	0.41218 (18)	0.24187 (12)	0.0227 (5)
C1	-0.0086 (2)	-0.1355 (2)	0.24468 (15)	0.0223 (6)
C2	0.1046 (2)	-0.1190 (2)	0.10228 (16)	0.0227 (6)

C3	0.1655 (2)	-0.0577 (2)	0.15660 (15)	0.0206 (6)
C4	-0.1125 (2)	-0.2116 (2)	0.13239 (15)	0.0306 (6)
H4A	-0.1132	-0.1750	0.0671	0.046*
H4B	-0.2073	-0.1713	0.1609	0.046*
H4C	-0.0875	-0.3181	0.1495	0.046*
C5	0.2778 (2)	-0.0006 (2)	0.12414 (15)	0.0215 (6)
H5	0.3074	0.0033	0.0628	0.026*
C6	0.3625 (2)	0.0560 (2)	0.16442 (15)	0.0193 (6)
C7	0.3494 (2)	0.0583 (2)	0.25650 (16)	0.0232 (6)
C8	0.4470 (3)	0.1070 (2)	0.28520 (16)	0.0307 (7)
H8	0.4385	0.1081	0.3460	0.037*
C9	0.5556 (3)	0.1535 (2)	0.22577 (16)	0.0312 (7)
H9	0.6200	0.1846	0.2469	0.037*
C10	0.5697 (2)	0.1544 (2)	0.13480 (16)	0.0288 (6)
H10	0.6424	0.1868	0.0941	0.035*
C11	0.4740 (2)	0.1065 (2)	0.10597 (15)	0.0251 (6)
H11	0.4834	0.1074	0.0447	0.030*
C13	-0.1095 (2)	-0.1721 (2)	0.32475 (15)	0.0213 (6)
C14	-0.1679 (2)	-0.0741 (2)	0.37879 (15)	0.0259 (6)
H14	-0.1459	0.0154	0.3627	0.031*
C15	-0.2590 (2)	-0.1091 (2)	0.45685 (16)	0.0316 (7)
H15	-0.2995	-0.0424	0.4925	0.038*
C16	-0.2899 (3)	-0.2423 (3)	0.48168 (16)	0.0335 (7)
H16	-0.3511	-0.2654	0.5342	0.040*
C17	-0.2304 (3)	-0.3420 (3)	0.42909 (17)	0.0403 (7)
H17	-0.2504	-0.4327	0.4465	0.048*
C18	-0.1414 (3)	-0.3063 (2)	0.35063 (16)	0.0342 (7)
H18	-0.1023	-0.3729	0.3147	0.041*
C21	0.0052 (2)	0.3587 (2)	0.23784 (15)	0.0213 (6)
C22	0.1193 (2)	0.3853 (2)	0.09563 (16)	0.0224 (6)
C23	0.1837 (2)	0.4354 (2)	0.15464 (15)	0.0218 (6)
C24	-0.0927 (2)	0.2851 (2)	0.12174 (15)	0.0300 (6)
H24A	-0.0467	0.2512	0.0685	0.045*
H24B	-0.1819	0.3626	0.1071	0.045*
H24C	-0.1133	0.2037	0.1696	0.045*
C25	0.2933 (2)	0.4973 (2)	0.12575 (15)	0.0241 (6)
H25	0.3284	0.5009	0.0650	0.029*
C26	0.3682 (2)	0.5595 (2)	0.16923 (16)	0.0219 (6)
C27	0.3378 (2)	0.5747 (2)	0.25873 (16)	0.0238 (6)
C28	0.4158 (3)	0.6439 (2)	0.28910 (16)	0.0288 (6)
H28	0.3938	0.6553	0.3482	0.035*
C29	0.5248 (3)	0.6955 (2)	0.23309 (17)	0.0314 (7)
H29	0.5759	0.7410	0.2544	0.038*
C30	0.5579 (2)	0.6795 (2)	0.14511 (17)	0.0319 (7)
H30	0.6319	0.7136	0.1071	0.038*
C31	0.4811 (2)	0.6132 (2)	0.11422 (16)	0.0289 (6)
H31	0.5042	0.6032	0.0548	0.035*
C33	-0.0964 (2)	0.3198 (2)	0.31732 (15)	0.0221 (6)

C34	-0.0444 (3)	0.2553 (2)	0.40301 (16)	0.0295 (6)
H34	0.0537	0.2357	0.4088	0.035*
C35	-0.1373 (3)	0.2200 (2)	0.47980 (16)	0.0334 (7)
H35	-0.1012	0.1757	0.5368	0.040*
C36	-0.2833 (3)	0.2502 (2)	0.47222 (16)	0.0290 (6)
H36	-0.3458	0.2261	0.5239	0.035*
C37	-0.3360 (3)	0.3161 (2)	0.38798 (16)	0.0282 (6)
H37	-0.4348	0.3376	0.3830	0.034*
C38	-0.2438 (2)	0.3510 (2)	0.31025 (15)	0.0254 (6)
H38	-0.2807	0.3952	0.2535	0.030*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0265 (10)	0.0404 (10)	0.0249 (11)	-0.0125 (8)	0.0015 (8)	-0.0138 (9)
O2	0.0353 (11)	0.0523 (11)	0.0223 (10)	-0.0289 (9)	0.0055 (9)	-0.0123 (8)
O3	0.0312 (11)	0.0353 (10)	0.0240 (11)	-0.0109 (9)	0.0004 (9)	-0.0096 (9)
O4	0.0483 (12)	0.0557 (12)	0.0318 (11)	-0.0307 (10)	0.0043 (10)	-0.0175 (9)
N1	0.0195 (12)	0.0289 (12)	0.0253 (13)	-0.0123 (10)	0.0000 (10)	-0.0099 (10)
N2	0.0184 (12)	0.0245 (11)	0.0230 (13)	-0.0092 (10)	0.0017 (10)	-0.0055 (9)
N3	0.0228 (13)	0.0275 (12)	0.0217 (12)	-0.0112 (10)	-0.0014 (10)	-0.0079 (10)
N4	0.0206 (12)	0.0259 (11)	0.0228 (13)	-0.0098 (10)	0.0011 (10)	-0.0068 (9)
C1	0.0199 (15)	0.0205 (14)	0.0256 (16)	-0.0033 (12)	-0.0017 (12)	-0.0078 (12)
C2	0.0187 (15)	0.0232 (14)	0.0241 (16)	-0.0041 (12)	-0.0015 (13)	-0.0061 (12)
C3	0.0170 (14)	0.0212 (13)	0.0217 (15)	-0.0042 (11)	-0.0019 (12)	-0.0044 (11)
C4	0.0265 (16)	0.0392 (16)	0.0340 (17)	-0.0156 (13)	-0.0043 (13)	-0.0136 (13)
C5	0.0208 (15)	0.0233 (14)	0.0171 (14)	-0.0032 (12)	-0.0008 (11)	-0.0049 (11)
C6	0.0164 (14)	0.0186 (13)	0.0208 (15)	-0.0044 (11)	-0.0010 (12)	-0.0036 (11)
C7	0.0217 (15)	0.0229 (14)	0.0244 (16)	-0.0086 (12)	0.0017 (12)	-0.0054 (12)
C8	0.0332 (16)	0.0412 (16)	0.0242 (16)	-0.0164 (14)	-0.0004 (13)	-0.0133 (13)
C9	0.0283 (16)	0.0406 (16)	0.0325 (17)	-0.0189 (13)	-0.0029 (13)	-0.0113 (13)
C10	0.0197 (15)	0.0357 (15)	0.0334 (17)	-0.0145 (13)	0.0025 (13)	-0.0084 (13)
C11	0.0227 (15)	0.0282 (14)	0.0239 (15)	-0.0070 (12)	0.0010 (12)	-0.0084 (12)
C13	0.0176 (14)	0.0243 (14)	0.0231 (15)	-0.0084 (12)	-0.0009 (12)	-0.0058 (12)
C14	0.0264 (15)	0.0258 (14)	0.0269 (15)	-0.0106 (12)	-0.0003 (12)	-0.0068 (12)
C15	0.0316 (16)	0.0329 (16)	0.0312 (17)	-0.0101 (13)	0.0045 (13)	-0.0135 (13)
C16	0.0296 (17)	0.0405 (16)	0.0319 (17)	-0.0190 (14)	0.0078 (13)	-0.0086 (14)
C17	0.0440 (18)	0.0402 (17)	0.0427 (18)	-0.0269 (15)	0.0126 (15)	-0.0134 (14)
C18	0.0376 (17)	0.0318 (16)	0.0386 (17)	-0.0162 (13)	0.0071 (14)	-0.0166 (13)
C21	0.0220 (15)	0.0184 (14)	0.0222 (16)	-0.0052 (12)	-0.0017 (12)	-0.0048 (12)
C22	0.0222 (15)	0.0193 (14)	0.0215 (16)	-0.0027 (11)	-0.0011 (13)	-0.0038 (12)
C23	0.0192 (14)	0.0233 (14)	0.0210 (15)	-0.0058 (12)	0.0015 (12)	-0.0057 (12)
C24	0.0318 (16)	0.0322 (15)	0.0303 (16)	-0.0111 (13)	-0.0054 (13)	-0.0109 (12)
C25	0.0229 (15)	0.0230 (14)	0.0227 (15)	-0.0039 (12)	0.0026 (12)	-0.0068 (12)
C26	0.0202 (15)	0.0194 (14)	0.0256 (16)	-0.0047 (12)	-0.0013 (12)	-0.0073 (12)
C27	0.0217 (15)	0.0236 (14)	0.0243 (16)	-0.0090 (12)	0.0012 (12)	-0.0032 (12)
C28	0.0324 (16)	0.0286 (14)	0.0257 (16)	-0.0059 (13)	-0.0073 (13)	-0.0081 (12)
C29	0.0250 (16)	0.0274 (15)	0.0443 (19)	-0.0077 (13)	-0.0107 (14)	-0.0087 (14)

C30	0.0205 (15)	0.0297 (15)	0.0430 (19)	-0.0080 (12)	0.0009 (13)	-0.0080 (14)
C31	0.0249 (16)	0.0252 (14)	0.0363 (17)	-0.0089 (12)	-0.0019 (13)	-0.0066 (13)
C33	0.0229 (15)	0.0215 (14)	0.0234 (16)	-0.0087 (12)	-0.0009 (12)	-0.0068 (12)
C34	0.0245 (15)	0.0384 (16)	0.0283 (16)	-0.0130 (13)	-0.0031 (13)	-0.0080 (13)
C35	0.0361 (17)	0.0436 (17)	0.0226 (16)	-0.0178 (14)	-0.0025 (13)	-0.0050 (13)
C36	0.0299 (17)	0.0327 (15)	0.0253 (16)	-0.0139 (13)	0.0061 (13)	-0.0091 (13)
C37	0.0207 (15)	0.0283 (15)	0.0354 (17)	-0.0094 (12)	0.0015 (13)	-0.0079 (13)
C38	0.0248 (15)	0.0257 (14)	0.0248 (15)	-0.0080 (12)	-0.0035 (12)	-0.0043 (12)

Geometric parameters (Å, °)

O1—C2	1.222 (2)	C14—H14	0.9300
O2—C7	1.347 (2)	C15—C16	1.373 (3)
O2—H2	0.8200	C15—H15	0.9300
O3—C22	1.218 (2)	C16—C17	1.380 (3)
O4—C27	1.341 (2)	C16—H16	0.9300
O4—H4	0.8200	C17—C18	1.378 (3)
N1—C2	1.392 (3)	C17—H17	0.9300
N1—C1	1.390 (3)	C18—H18	0.9300
N1—C4	1.458 (2)	C21—C33	1.467 (3)
N2—C1	1.311 (2)	C22—C23	1.474 (3)
N2—C3	1.397 (3)	C23—C25	1.348 (3)
N3—C21	1.383 (3)	C24—H24A	0.9600
N3—C22	1.392 (3)	C24—H24B	0.9600
N3—C24	1.458 (2)	C24—H24C	0.9600
N4—C21	1.306 (2)	C25—C26	1.450 (3)
N4—C23	1.400 (3)	C25—H25	0.9300
C1—C13	1.464 (3)	C26—C27	1.402 (3)
C2—C3	1.471 (3)	C26—C31	1.410 (3)
C3—C5	1.347 (3)	C27—C28	1.396 (3)
C4—H4A	0.9600	C28—C29	1.376 (3)
C4—H4B	0.9600	C28—H28	0.9300
C4—H4C	0.9600	C29—C30	1.381 (3)
C5—C6	1.449 (3)	C29—H29	0.9300
C5—H5	0.9300	C30—C31	1.371 (3)
C6—C11	1.406 (3)	C30—H30	0.9300
C6—C7	1.409 (3)	C31—H31	0.9300
C7—C8	1.392 (3)	C33—C34	1.388 (3)
C8—C9	1.374 (3)	C33—C38	1.389 (3)
C8—H8	0.9300	C34—C35	1.382 (3)
C9—C10	1.382 (3)	C34—H34	0.9300
C9—H9	0.9300	C35—C36	1.378 (3)
C10—C11	1.372 (3)	C35—H35	0.9300
C10—H10	0.9300	C36—C37	1.373 (3)
C11—H11	0.9300	C36—H36	0.9300
C13—C14	1.383 (3)	C37—C38	1.387 (3)
C13—C18	1.388 (3)	C37—H37	0.9300
C14—C15	1.385 (3)	C38—H38	0.9300

C7—O2—H2	109.5	C18—C17—H17	120.2
C27—O4—H4	109.5	C16—C17—H17	120.2
C2—N1—C1	108.23 (18)	C17—C18—C13	120.6 (2)
C2—N1—C4	122.40 (19)	C17—C18—H18	119.7
C1—N1—C4	128.31 (19)	C13—C18—H18	119.7
C1—N2—C3	106.55 (18)	N4—C21—N3	112.87 (19)
C21—N3—C22	108.32 (18)	N4—C21—C33	122.4 (2)
C21—N3—C24	128.78 (18)	N3—C21—C33	124.7 (2)
C22—N3—C24	122.86 (19)	O3—C22—N3	125.6 (2)
C21—N4—C23	106.86 (19)	O3—C22—C23	130.9 (2)
N2—C1—N1	112.80 (19)	N3—C22—C23	103.46 (19)
N2—C1—C13	123.5 (2)	C25—C23—N4	127.5 (2)
N1—C1—C13	123.68 (19)	C25—C23—C22	124.1 (2)
O1—C2—N1	125.2 (2)	N4—C23—C22	108.41 (18)
O1—C2—C3	131.3 (2)	N3—C24—H24A	109.5
N1—C2—C3	103.43 (19)	N3—C24—H24B	109.5
C5—C3—N2	128.0 (2)	H24A—C24—H24B	109.5
C5—C3—C2	123.0 (2)	N3—C24—H24C	109.5
N2—C3—C2	108.99 (18)	H24A—C24—H24C	109.5
N1—C4—H4A	109.5	H24B—C24—H24C	109.5
N1—C4—H4B	109.5	C23—C25—C26	133.6 (2)
H4A—C4—H4B	109.5	C23—C25—H25	113.2
N1—C4—H4C	109.5	C26—C25—H25	113.2
H4A—C4—H4C	109.5	C27—C26—C31	117.1 (2)
H4B—C4—H4C	109.5	C27—C26—C25	126.9 (2)
C3—C5—C6	133.7 (2)	C31—C26—C25	115.9 (2)
C3—C5—H5	113.2	O4—C27—C28	114.2 (2)
C6—C5—H5	113.2	O4—C27—C26	125.9 (2)
C11—C6—C7	117.1 (2)	C28—C27—C26	120.0 (2)
C11—C6—C5	116.0 (2)	C29—C28—C27	121.1 (2)
C7—C6—C5	126.9 (2)	C29—C28—H28	119.4
O2—C7—C8	115.5 (2)	C27—C28—H28	119.4
O2—C7—C6	125.2 (2)	C28—C29—C30	119.8 (2)
C8—C7—C6	119.3 (2)	C28—C29—H29	120.1
C9—C8—C7	121.5 (2)	C30—C29—H29	120.1
C9—C8—H8	119.2	C31—C30—C29	119.5 (2)
C7—C8—H8	119.2	C31—C30—H30	120.2
C8—C9—C10	120.4 (2)	C29—C30—H30	120.2
C8—C9—H9	119.8	C30—C31—C26	122.4 (2)
C10—C9—H9	119.8	C30—C31—H31	118.8
C11—C10—C9	118.4 (2)	C26—C31—H31	118.8
C11—C10—H10	120.8	C34—C33—C38	118.9 (2)
C9—C10—H10	120.8	C34—C33—C21	118.9 (2)
C10—C11—C6	123.3 (2)	C38—C33—C21	122.2 (2)
C10—C11—H11	118.4	C35—C34—C33	120.6 (2)
C6—C11—H11	118.4	C35—C34—H34	119.7
C14—C13—C18	119.2 (2)	C33—C34—H34	119.7

C14—C13—C1	119.10 (19)	C36—C35—C34	120.2 (2)
C18—C13—C1	121.6 (2)	C36—C35—H35	119.9
C13—C14—C15	120.1 (2)	C34—C35—H35	119.9
C13—C14—H14	119.9	C37—C36—C35	119.6 (2)
C15—C14—H14	119.9	C37—C36—H36	120.2
C16—C15—C14	120.1 (2)	C35—C36—H36	120.2
C16—C15—H15	119.9	C36—C37—C38	120.8 (2)
C14—C15—H15	119.9	C36—C37—H37	119.6
C15—C16—C17	120.3 (2)	C38—C37—H37	119.6
C15—C16—H16	119.8	C33—C38—C37	119.9 (2)
C17—C16—H16	119.8	C33—C38—H38	120.1
C18—C17—C16	119.6 (2)	C37—C38—H38	120.1
C2—N1—C1—N2	-0.1 (2)	C3—C5—C6—C7	3.3 (4)
C2—N1—C1—C13	-179.03 (19)	C3—C5—C6—C11	179.8 (2)
C4—N1—C1—N2	-168.39 (19)	C5—C6—C7—O2	-3.5 (3)
C1—N1—C2—O1	179.9 (2)	C5—C6—C7—C8	175.3 (2)
C1—N1—C2—C3	0.0 (2)	C11—C6—C7—O2	-179.95 (18)
C4—N1—C2—O1	-11.0 (3)	C11—C6—C7—C8	-1.1 (3)
C4—N1—C2—C3	169.06 (17)	C5—C6—C11—C10	-175.71 (19)
C3—N2—C1—N1	0.3 (2)	C7—C6—C11—C10	1.1 (3)
C3—N2—C1—C13	179.15 (19)	O2—C7—C8—C9	179.22 (19)
C1—N2—C3—C2	-0.3 (2)	C6—C7—C8—C9	0.3 (3)
C1—N2—C3—C5	-179.1 (2)	C7—C8—C9—C10	0.7 (3)
N1—C1—C13—C14	-142.3 (2)	C8—C9—C10—C11	-0.7 (3)
N1—C1—C13—C18	41.4 (3)	C9—C10—C11—C6	-0.2 (3)
N2—C1—C13—C14	38.9 (3)	C1—C13—C14—C15	-177.4 (2)
N2—C1—C13—C18	-137.4 (2)	C18—C13—C14—C15	-1.1 (3)
O1—C2—C3—N2	-179.8 (2)	C1—C13—C18—C17	176.4 (2)
O1—C2—C3—C5	-0.8 (4)	C14—C13—C18—C17	0.1 (4)
N1—C2—C3—N2	0.2 (2)	C13—C14—C15—C16	1.0 (3)
N1—C2—C3—C5	179.09 (19)	C14—C15—C16—C17	0.0 (4)
N2—C3—C5—C6	3.6 (4)	C15—C16—C17—C18	-1.0 (4)
C2—C3—C5—C6	-175.1 (2)	C16—C17—C18—C13	0.9 (4)

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O2—H2 \cdots N2	0.82	1.80	2.615 (2)	176
O4—H4 \cdots N4	0.82	1.79	2.612 (3)	175
C4—H4 <i>A</i> \cdots O1	0.96	2.56	2.896 (3)	100
C10—H10 \cdots O1 ⁱ	0.93	2.58	3.403 (3)	148
C30—H30 \cdots O3 ⁱⁱ	0.93	2.68	3.421 (3)	137
C24—H24 <i>A</i> \cdots O3	0.96	2.56	2.902 (3)	101

Symmetry codes: (i) $-x+1, -y, -z$; (ii) $-x+1, -y+1, -z$.