

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

7-Chloro-4-[(*E*)-2-(3,4,5-trimethoxybenzylidene)hydrazin-1-yl]quinoline

Marcelle de Lima Ferreira,^a Marcus V. N. de Souza,^a
Solange M. S. V. Wardell,^b Edward R. T. Tiekink^{c*} and
James L. Wardell^{d‡}

^aInstituto de Tecnologia em Fármacos—Farmanguinhos, FioCruz—Fundação Oswaldo Cruz, R. Sizenando Nabuco, 100, Manguinhos, 21041-250 Rio de Janeiro, RJ, Brazil, ^bCHEMSOL, 1 Harcourt Road, Aberdeen AB15 5NY, Scotland, ^cDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia, and ^dCentro de Desenvolvimento Tecnológico em Saúde (CDTS), Fundação Oswaldo Cruz (FIOCRUZ), Casa Amarela, Campus de Manguinhos, Av. Brasil 4365, 21040-900 Rio de Janeiro, RJ, Brazil

Correspondence e-mail: edward.tiekink@gmail.com


Received 23 March 2012; accepted 23 March 2012

Key indicators: single-crystal X-ray study; $T = 120$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.036; wR factor = 0.101; data-to-parameter ratio = 16.2.

In the title compound, $\text{C}_{19}\text{H}_{18}\text{ClN}_3\text{O}_3$, the r.m.s. deviation through the 23 non-H and non-methoxy atoms is 0.088 Å, indicating a planar molecule with the exception of the methoxy groups. One methoxy group, surrounded on either side by the other methoxy groups, is almost normal to the benzene ring to which it is connected [$\text{C}-\text{O}-\text{C}_{\text{ar}}-\text{C}_{\text{ar}}$ torsion angle = $81.64(15)^\circ$]. In the crystal, $\text{N}-\text{H}\cdots\text{O}$, $\text{C}-\text{H}\cdots\text{O}$ and $\pi-\pi$ interactions [between quinoline residues; centroid-centroid distance = $3.4375(8)$ Å] link molecules into a three-dimensional architecture.

Related literature

For the biological activity, including anti-tubercular and anti-tumour activity, of compounds containing the quinolinyl nucleus, see: de Souza *et al.* (2009); Candea *et al.* (2009); Montenegro *et al.* (2012). For related structures, see: Howie *et al.* (2010); de Souza *et al.* (2010, 2012).



Experimental

Crystal data

 $\text{C}_{19}\text{H}_{18}\text{ClN}_3\text{O}_3$ $M_r = 371.81$ Orthorhombic, *Pbca* $a = 7.6338(2)$ Å $b = 15.5335(4)$ Å $c = 28.7960(7)$ Å $V = 3414.62(15)$ Å³ $Z = 8$ Mo $K\alpha$ radiation $\mu = 0.25$ mm⁻¹ $T = 120$ K $0.45 \times 0.40 \times 0.30$ mm

Data collection

Bruker–Nonius Roper CCD camera

on a κ -goniostat diffractometer

Absorption correction: multi-scan

(SADABS; Sheldrick, 2007)

 $T_{\text{min}} = 0.652$, $T_{\text{max}} = 0.746$

22673 measured reflections

3899 independent reflections

3353 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.040$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.036$ $wR(F^2) = 0.101$ $S = 1.02$

3899 reflections

241 parameters

1 restraint

H atoms treated by a mixture of independent and constrained refinement

 $\Delta\rho_{\text{max}} = 0.30$ e Å⁻³ $\Delta\rho_{\text{min}} = -0.30$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N}2-\text{H}2\cdots\text{O}3^{\text{i}}$	0.87 (1)	2.53 (2)	3.0349 (15)	118 (1)
$\text{C}19-\text{H}19\text{B}\cdots\text{N}1^{\text{ii}}$	0.98	2.48	3.3602 (18)	149

Symmetry codes: (i) $x + \frac{1}{2}, -y + \frac{1}{2}, -z$; (ii) $x, -y + \frac{1}{2}, z - \frac{1}{2}$.

Data collection: *COLLECT* (Hooft, 1998); cell refinement: *DENZO* (Otwinowski & Minor, 1997) and *COLLECT*; data reduction: *DENZO* and *COLLECT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *DIAMOND* (Brandenburg, 2006); software used to prepare material for publication: *pubCIF* (Westrip, 2010).

The use of the EPSRC X-ray crystallographic service at the University of Southampton, England, and the valuable assistance of the staff there is gratefully acknowledged. JLW acknowledges support from CAPES (Brazil). Support from the Ministry of Higher Education, Malaysia, High-Impact

‡ Additional correspondence author, e-mail: j.wardell@abdn.ac.uk.

Research scheme (UM.C/HIR/MOHE/SC/12) is gratefully acknowledged.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5858).

References

- Brandenburg, K. (2006). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Candea, A. L. P., Ferreira, M. de L., Pais, K. C., Cardoso, L. N. de F., Kaiser, C. R., Henriques, M., das, G. M. de O., Lourenco, M. C. S., Bezerra, F. A. F. M. & de Souza, M. V. N. (2009). *Bioorg. Med. Chem. Lett.* **19**, 6272–6274.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Hooft, R. W. W. (1998). *COLLECT*. Nonius BV, Delft, The Netherlands.
- Howie, R. A., de Souza, M. V. N., Ferreira, M. de L., Kaiser, C. R., Wardell, J. L. & Wardell, S. M. S. V. (2010). *Z. Kristallogr.* **225**, 440–447.
- Montenegro, R. C., Lotufo, L. V., de Moraes, M. O., do O Pessoa, C., Rodrigues, F. A. R., Bispo, M. L. F., Freire, B. A., Kaiser, C. R. & de Souza, M. V. N. (2012). *Lett. Drug Des. Discov.* **9**, 251–256.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Sheldrick, G. M. (2007). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Souza, M. V. N. de, de Lima Ferreira, M., Wardell, S. M. S. V., Tiekink, E. R. T. & Wardell, J. L. (2012). *Acta Cryst.* **E68**, o1244–o1245.
- Souza, M. V. N. de, Howie, R. A., Tiekink, E. R. T., Wardell, J. L., Wardell, S. M. S. V. & Kaiser, C. R. (2010). *Acta Cryst.* **E66**, o698–o699.
- Souza, M. V. N. de, Pais, K. C., Kaiser, C. R., Peralta, M. A., Ferreira, M. de L. & Lourenco, M. C. S. (2009). *Bioorg. Med. Chem.* **17**, 1474–1480.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2012). E68, o1214–o1215 [https://doi.org/10.1107/S1600536812012755]

7-Chloro-4-[(*E*)-2-(3,4,5-trimethoxybenzylidene)hydrazin-1-yl]quinoline

Marcelle de Lima Ferreira, Marcus V. N. de Souza, Solange M. S. V. Wardell, Edward R. T. Tiekink and James L. Wardell

S1. Comment

The title compound, (I), was investigated as part of on-going crystallographic investigations of arylaldehyde 7-chloro-quinoline-4-hydrazone species (Howie *et al.*, 2010; de Souza *et al.*, 2010; de Souza *et al.*, 2012). The structural studies complement biological studies which show these hydrazones to possess a wide range of pharmacological activities such as anti-tubercular (Candea *et al.*, 2009) and anti-tumour (Montenegro *et al.*, 2012) activities, which are ascribed to the presence of the quinoline nucleus (de Souza *et al.*, 2009).

In (I), Fig. 1, with the exception of two of the methoxy groups, the molecule is planar. The r.m.s. deviation through the 23 non-hydrogen and non-methoxy atoms is 0.0879 Å. The maximum deviations from this plane are 0.1219 (11) Å for the N2 atom and -0.1498 (11) for the C14 atom. The terminal carbon atoms, C17–C19, of the methoxy groups lie -0.0840 (17), 0.7910 (16) and -0.3504 (19) Å, respectively, out of the least-squares plane, indicating that the central methoxy group is almost orthogonal to the benzene ring to which it is connected with the C18—O2—C14—C13 torsion angle being 81.64 (15)°. The conformation about the N3=C10 bond [1.2829 (17) Å] is *E*.

In the crystal packing, weak N—H···O hydrogen bonds along with C—H···O interactions, Table 1, and π — π interactions between symmetry related quinolinyl residues [centroid···centroid distance = 3.4375 (8) Å for symmetry operation $-1/2 + x, y, 1/2 - z$] link molecules into a three-dimensional architecture. Globally, molecules stack along the *c* axis with alternating layers of quinolinyl and trimethoxybenzene residues, Fig. 2.

S2. Experimental

The compound was prepared from 7-chloro-4-quinolinylhydrazone and 3,4,5-trimethoxybenzaldehyde (Montenegro *et al.*, 2012) and was recrystallized from an EtOCH₂CH₂OH solution of the compound.

S3. Refinement

The C-bound H atoms were geometrically placed (C—H = 0.95–0.98 Å) and refined as riding with $U_{\text{iso}}(\text{H}) = 1.2–1.5U_{\text{eq}}(\text{C})$. The N-bound H-atom was located in a difference Fourier map and refined with N—H = 0.88±0.01 Å, and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{N})$. Owing to poor agreement, the (2 2 0), (2 3 0), (0 4 1), (2 2 1), (1 0 4), (2 1 2) and (1 0 2) reflections were omitted from the final cycles of refinement.




Figure 1

The molecular structure showing the atom-labelling scheme and displacement ellipsoids at the 50% probability level.




Figure 2

A view in projection down the a axis of the unit-cell contents of (I). The N—H \cdots O, C—H \cdots N and $\pi\cdots\pi$ interactions are shown as orange, blue and purple dashed lines, respectively.

7-Chloro-4-[(*E*)-2-(3,4,5-trimethoxybenzylidene)hydrazin-1-yl]quinoline

Crystal data

$C_{19}H_{18}ClN_3O_3$

$M_r = 371.81$

Orthorhombic, $Pbca$

Hall symbol: $-P\ 2ac\ 2ab$

$a = 7.6338\ (2)\ \text{\AA}$

$b = 15.5335\ (4)\ \text{\AA}$

$c = 28.7960\ (7)\ \text{\AA}$

$V = 3414.62\ (15)\ \text{\AA}^3$

$Z = 8$

$F(000) = 1552$

$D_x = 1.447\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 24183 reflections

$\theta = 2.9\text{--}27.5^\circ$

$\mu = 0.25\ \text{mm}^{-1}$

$T = 120\ \text{K}$

Prism, colourless

$0.45 \times 0.40 \times 0.30\ \text{mm}$

Data collection

Bruker–Nonius Roper CCD camera on a κ -goniostat diffractometer

Radiation source: Bruker–Nonius FR591 rotating anode

Graphite monochromator

Detector resolution: $9.091\ \text{pixels mm}^{-1}$

φ and ω scans

Absorption correction: multi-scan (*SADABS*; Sheldrick, 2007)

$T_{\min} = 0.652$, $T_{\max} = 0.746$

22673 measured reflections

3899 independent reflections

3353 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.040$

$\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 3.0^\circ$

$h = -7 \rightarrow 9$
 $k = -19 \rightarrow 20$

$l = -26 \rightarrow 37$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.036$
 $wR(F^2) = 0.101$
 $S = 1.02$
 3899 reflections
 241 parameters
 1 restraint
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H atoms treated by a mixture of independent
 and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0576P)^2 + 1.3595P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.30 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.30 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	0.32570 (5)	0.04598 (2)	0.339654 (12)	0.02558 (12)
O1	-0.29202 (13)	0.51292 (6)	-0.01170 (3)	0.0189 (2)
O2	-0.26948 (13)	0.42177 (6)	-0.09143 (3)	0.0185 (2)
O3	-0.08960 (13)	0.27516 (6)	-0.09550 (3)	0.0185 (2)
N1	0.03404 (15)	0.32333 (7)	0.28807 (4)	0.0185 (2)
N2	0.06914 (15)	0.25871 (7)	0.14639 (4)	0.0166 (2)
H2n	0.123 (2)	0.2115 (8)	0.1386 (5)	0.020*
N3	-0.00092 (14)	0.30964 (7)	0.11233 (4)	0.0160 (2)
C1	-0.02236 (18)	0.37484 (8)	0.25478 (5)	0.0183 (3)
H1	-0.0738	0.4278	0.2641	0.022*
C2	-0.01331 (18)	0.35863 (9)	0.20702 (5)	0.0168 (3)
H2A	-0.0579	0.3991	0.1853	0.020*
C3	0.06177 (17)	0.28249 (8)	0.19210 (4)	0.0144 (3)
C4	0.12967 (17)	0.22434 (8)	0.22641 (4)	0.0141 (3)
C5	0.21216 (17)	0.14533 (9)	0.21567 (5)	0.0167 (3)
H5	0.2271	0.1295	0.1840	0.020*
C6	0.27103 (18)	0.09103 (9)	0.24970 (5)	0.0177 (3)
H6	0.3268	0.0383	0.2419	0.021*
C7	0.24732 (18)	0.11490 (9)	0.29641 (5)	0.0174 (3)
C8	0.16982 (18)	0.19071 (9)	0.30860 (5)	0.0179 (3)
H8	0.1564	0.2052	0.3405	0.021*
C9	0.10938 (17)	0.24777 (9)	0.27378 (4)	0.0152 (3)

C10	0.01112 (17)	0.27774 (9)	0.07136 (5)	0.0157 (3)
H10	0.0656	0.2231	0.0679	0.019*
C11	-0.05455 (17)	0.32096 (9)	0.02980 (4)	0.0152 (3)
C12	-0.14098 (17)	0.40041 (8)	0.03151 (4)	0.0155 (3)
H12	-0.1551	0.4297	0.0602	0.019*
C13	-0.20597 (17)	0.43605 (8)	-0.00934 (5)	0.0153 (3)
C14	-0.18806 (17)	0.39204 (9)	-0.05162 (4)	0.0157 (3)
C15	-0.09928 (17)	0.31355 (8)	-0.05291 (4)	0.0152 (3)
C16	-0.03088 (18)	0.27845 (8)	-0.01241 (4)	0.0160 (3)
H16	0.0318	0.2257	-0.0135	0.019*
C17	-0.31426 (19)	0.55864 (9)	0.03114 (5)	0.0219 (3)
H17A	-0.3887	0.5249	0.0521	0.033*
H17B	-0.3696	0.6144	0.0250	0.033*
H17C	-0.1997	0.5678	0.0456	0.033*
C18	-0.1751 (2)	0.48981 (9)	-0.11456 (5)	0.0225 (3)
H18A	-0.1514	0.5364	-0.0925	0.034*
H18B	-0.2456	0.5120	-0.1404	0.034*
H18C	-0.0641	0.4673	-0.1266	0.034*
C19	-0.0276 (2)	0.18760 (9)	-0.09654 (5)	0.0228 (3)
H19A	0.0954	0.1860	-0.0869	0.034*
H19B	-0.0382	0.1648	-0.1282	0.034*
H19C	-0.0979	0.1524	-0.0753	0.034*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.0281 (2)	0.0275 (2)	0.02106 (19)	0.00361 (14)	-0.00295 (14)	0.00990 (14)
O1	0.0219 (5)	0.0175 (5)	0.0174 (5)	0.0038 (4)	-0.0027 (4)	0.0002 (4)
O2	0.0205 (5)	0.0199 (5)	0.0151 (5)	-0.0036 (4)	-0.0054 (4)	0.0060 (4)
O3	0.0240 (5)	0.0200 (5)	0.0115 (4)	0.0007 (4)	-0.0005 (4)	-0.0013 (4)
N1	0.0204 (6)	0.0187 (6)	0.0164 (5)	-0.0006 (5)	0.0001 (4)	-0.0014 (5)
N2	0.0216 (6)	0.0163 (5)	0.0120 (5)	0.0049 (5)	-0.0013 (4)	0.0017 (4)
N3	0.0168 (6)	0.0177 (5)	0.0135 (5)	-0.0003 (4)	-0.0015 (4)	0.0039 (4)
C1	0.0193 (7)	0.0155 (6)	0.0201 (7)	0.0008 (5)	0.0019 (5)	-0.0019 (5)
C2	0.0187 (7)	0.0154 (6)	0.0164 (6)	-0.0006 (5)	-0.0004 (5)	0.0022 (5)
C3	0.0133 (6)	0.0161 (6)	0.0139 (6)	-0.0025 (5)	0.0002 (5)	0.0008 (5)
C4	0.0133 (6)	0.0157 (6)	0.0133 (6)	-0.0033 (5)	-0.0001 (5)	0.0013 (5)
C5	0.0168 (6)	0.0185 (6)	0.0148 (6)	-0.0011 (5)	-0.0010 (5)	-0.0010 (5)
C6	0.0172 (7)	0.0151 (6)	0.0209 (7)	-0.0001 (5)	-0.0018 (5)	-0.0004 (5)
C7	0.0161 (6)	0.0191 (6)	0.0169 (6)	-0.0031 (5)	-0.0030 (5)	0.0057 (5)
C8	0.0186 (7)	0.0222 (7)	0.0128 (6)	-0.0024 (5)	0.0001 (5)	0.0005 (5)
C9	0.0139 (6)	0.0166 (6)	0.0151 (6)	-0.0028 (5)	0.0002 (5)	-0.0001 (5)
C10	0.0166 (6)	0.0154 (6)	0.0152 (6)	0.0000 (5)	-0.0006 (5)	0.0009 (5)
C11	0.0134 (6)	0.0184 (6)	0.0139 (6)	-0.0028 (5)	-0.0008 (5)	0.0019 (5)
C12	0.0162 (6)	0.0174 (6)	0.0129 (6)	-0.0025 (5)	-0.0012 (5)	-0.0005 (5)
C13	0.0137 (6)	0.0137 (6)	0.0184 (7)	-0.0018 (5)	-0.0009 (5)	0.0012 (5)
C14	0.0157 (6)	0.0182 (6)	0.0132 (6)	-0.0030 (5)	-0.0034 (5)	0.0035 (5)
C15	0.0165 (6)	0.0169 (6)	0.0123 (6)	-0.0040 (5)	0.0004 (5)	-0.0002 (5)

C16	0.0165 (6)	0.0158 (6)	0.0159 (6)	-0.0009 (5)	-0.0001 (5)	0.0019 (5)
C17	0.0241 (7)	0.0196 (7)	0.0218 (7)	0.0045 (6)	-0.0039 (6)	-0.0038 (6)
C18	0.0281 (8)	0.0202 (7)	0.0190 (7)	-0.0047 (6)	-0.0019 (6)	0.0062 (6)
C19	0.0316 (8)	0.0207 (7)	0.0162 (6)	0.0024 (6)	0.0010 (6)	-0.0032 (6)

Geometric parameters (Å, °)

C11—C7	1.7478 (13)	C6—H6	0.9500
O1—C13	1.3646 (16)	C7—C8	1.364 (2)
O1—C17	1.4335 (17)	C8—C9	1.4156 (18)
O2—C14	1.3836 (15)	C8—H8	0.9500
O2—C18	1.4422 (16)	C10—C11	1.4609 (18)
O3—C15	1.3659 (15)	C10—H10	0.9500
O3—C19	1.4403 (17)	C11—C16	1.3949 (18)
N1—C1	1.3208 (18)	C11—C12	1.4003 (18)
N1—C9	1.3704 (17)	C12—C13	1.3915 (18)
N2—C3	1.3684 (16)	C12—H12	0.9500
N2—N3	1.3687 (15)	C13—C14	1.4030 (18)
N2—H2n	0.871 (9)	C14—C15	1.3954 (19)
N3—C10	1.2829 (17)	C15—C16	1.3893 (18)
C1—C2	1.4001 (19)	C16—H16	0.9500
C1—H1	0.9500	C17—H17A	0.9800
C2—C3	1.3826 (19)	C17—H17B	0.9800
C2—H2A	0.9500	C17—H17C	0.9800
C3—C4	1.4355 (18)	C18—H18A	0.9800
C4—C5	1.4136 (19)	C18—H18B	0.9800
C4—C9	1.4202 (17)	C18—H18C	0.9800
C5—C6	1.3689 (19)	C19—H19A	0.9800
C5—H5	0.9500	C19—H19B	0.9800
C6—C7	1.4068 (19)	C19—H19C	0.9800
C13—O1—C17	116.59 (10)	C11—C10—H10	118.3
C14—O2—C18	113.76 (10)	C16—C11—C12	120.60 (12)
C15—O3—C19	116.66 (10)	C16—C11—C10	116.86 (12)
C1—N1—C9	115.96 (12)	C12—C11—C10	122.53 (12)
C3—N2—N3	121.14 (11)	C13—C12—C11	119.26 (12)
C3—N2—H2n	119.7 (11)	C13—C12—H12	120.4
N3—N2—H2n	119.1 (11)	C11—C12—H12	120.4
C10—N3—N2	114.06 (11)	O1—C13—C12	124.19 (12)
N1—C1—C2	126.01 (13)	O1—C13—C14	115.47 (11)
N1—C1—H1	117.0	C12—C13—C14	120.33 (12)
C2—C1—H1	117.0	O2—C14—C15	119.20 (12)
C3—C2—C1	118.65 (12)	O2—C14—C13	120.84 (12)
C3—C2—H2A	120.7	C15—C14—C13	119.75 (12)
C1—C2—H2A	120.7	O3—C15—C16	124.20 (12)
N2—C3—C2	123.15 (12)	O3—C15—C14	115.58 (11)
N2—C3—C4	118.51 (12)	C16—C15—C14	120.21 (12)
C2—C3—C4	118.30 (12)	C15—C16—C11	119.79 (12)

C5—C4—C9	118.76 (12)	C15—C16—H16	120.1
C5—C4—C3	123.82 (12)	C11—C16—H16	120.1
C9—C4—C3	117.41 (12)	O1—C17—H17A	109.5
C6—C5—C4	121.64 (12)	O1—C17—H17B	109.5
C6—C5—H5	119.2	H17A—C17—H17B	109.5
C4—C5—H5	119.2	O1—C17—H17C	109.5
C5—C6—C7	118.67 (12)	H17A—C17—H17C	109.5
C5—C6—H6	120.7	H17B—C17—H17C	109.5
C7—C6—H6	120.7	O2—C18—H18A	109.5
C8—C7—C6	121.97 (12)	O2—C18—H18B	109.5
C8—C7—Cl1	119.60 (10)	H18A—C18—H18B	109.5
C6—C7—Cl1	118.40 (11)	O2—C18—H18C	109.5
C7—C8—C9	119.98 (12)	H18A—C18—H18C	109.5
C7—C8—H8	120.0	H18B—C18—H18C	109.5
C9—C8—H8	120.0	O3—C19—H19A	109.5
N1—C9—C8	117.41 (12)	O3—C19—H19B	109.5
N1—C9—C4	123.62 (12)	H19A—C19—H19B	109.5
C8—C9—C4	118.97 (12)	O3—C19—H19C	109.5
N3—C10—C11	123.46 (12)	H19A—C19—H19C	109.5
N3—C10—H10	118.3	H19B—C19—H19C	109.5
C3—N2—N3—C10	-177.97 (12)	N2—N3—C10—C11	-179.92 (12)
C9—N1—C1—C2	0.7 (2)	N3—C10—C11—C16	179.59 (12)
N1—C1—C2—C3	-0.4 (2)	N3—C10—C11—C12	-1.8 (2)
N3—N2—C3—C2	-0.9 (2)	C16—C11—C12—C13	1.09 (19)
N3—N2—C3—C4	176.90 (11)	C10—C11—C12—C13	-177.49 (12)
C1—C2—C3—N2	176.69 (12)	C17—O1—C13—C12	0.29 (18)
C1—C2—C3—C4	-1.07 (19)	C17—O1—C13—C14	179.15 (12)
N2—C3—C4—C5	3.29 (19)	C11—C12—C13—O1	-179.96 (12)
C2—C3—C4—C5	-178.84 (12)	C11—C12—C13—C14	1.23 (19)
N2—C3—C4—C9	-175.68 (11)	C18—O2—C14—C15	-103.67 (14)
C2—C3—C4—C9	2.19 (18)	C18—O2—C14—C13	81.64 (15)
C9—C4—C5—C6	0.58 (19)	O1—C13—C14—O2	-6.46 (18)
C3—C4—C5—C6	-178.38 (13)	C12—C13—C14—O2	172.44 (12)
C4—C5—C6—C7	0.3 (2)	O1—C13—C14—C15	178.87 (11)
C5—C6—C7—C8	-0.9 (2)	C12—C13—C14—C15	-2.22 (19)
C5—C6—C7—Cl1	-179.08 (10)	C19—O3—C15—C16	9.53 (19)
C6—C7—C8—C9	0.4 (2)	C19—O3—C15—C14	-169.05 (12)
Cl1—C7—C8—C9	178.63 (10)	O2—C14—C15—O3	4.77 (18)
C1—N1—C9—C8	-179.44 (12)	C13—C14—C15—O3	179.53 (11)
C1—N1—C9—C4	0.65 (19)	O2—C14—C15—C16	-173.87 (12)
C7—C8—C9—N1	-179.42 (12)	C13—C14—C15—C16	0.88 (19)
C7—C8—C9—C4	0.50 (19)	O3—C15—C16—C11	-177.10 (12)
C5—C4—C9—N1	178.92 (12)	C14—C15—C16—C11	1.4 (2)
C3—C4—C9—N1	-2.05 (19)	C12—C11—C16—C15	-2.4 (2)
C5—C4—C9—C8	-0.99 (18)	C10—C11—C16—C15	176.23 (12)
C3—C4—C9—C8	178.03 (12)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N2—H2 <i>n</i> \cdots O3 ⁱ	0.87 (1)	2.53 (2)	3.0349 (15)	118 (1)
C19—H19 <i>B</i> \cdots N1 ⁱⁱ	0.98	2.48	3.3602 (18)	149

Symmetry codes: (i) $x+1/2, -y+1/2, -z$; (ii) $x, -y+1/2, z-1/2$.