

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

(E)-4-[(4-Diethylamino-2-hydroxybenzylidene)amino]benzoic acidHadi Kargar,^{a*} Zahra Sharafi,^b Reza Kia^c and Muhammad Nawaz Tahir^{d*}

^aDepartment of Chemistry, Payame Noor University, PO Box 19395-3697 Tehran, I. R. of IRAN, ^bDepartment of Chemistry, Marvdasht Branch, Islamic Azad University, Marvdasht, Iran, ^cDepartment of Chemistry, Science and Research Branch, Islamic Azad University, Tehran, Iran, and ^dDepartment of Physics, University of Sargodha, Punjab, Pakistan
Correspondence e-mail: h.kargar@pnu.ac.ir, dmntahir_uos@yahoo.com


Received 4 March 2012; accepted 6 March 2012

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; disorder in main residue; R factor = 0.058; wR factor = 0.133; data-to-parameter ratio = 13.1.

In the title compound, $\text{C}_{18}\text{H}_{20}\text{N}_2\text{O}_3$, a potential bidentate N,O -donor Schiff base ligand, the benzene rings are inclined at an angle of 12.25 (19)°. The molecule has an E conformation about the $\text{C}=\text{N}$ bond. One of the ethyl groups is disordered over two positions, with a refined site-occupancy ratio of 0.55 (1):0.45 (1). An intramolecular $\text{O}-\text{H}\cdots\text{N}$ hydrogen bond makes an $S(6)$ ring motif. In the crystal, pairs of $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds link molecules, forming inversion dimers with $R_2^2(8)$ ring motifs.

Related literature

For background to Schiff base ligands and their metal complexes, see: Kargar *et al.* (2011, 2012); Kia *et al.* (2010). For standard bond lengths, see: Allen *et al.* (1987). For hydrogen-bond motifs, see: Bernstein *et al.* (1995).



Experimental

Crystal data

$\text{C}_{18}\text{H}_{20}\text{N}_2\text{O}_3$
 $M_r = 312.36$
Monoclinic, $P2_1/c$
 $a = 12.4216$ (8) Å
 $b = 8.1511$ (6) Å
 $c = 16.0820$ (11) Å
 $\beta = 93.001$ (3)°

$V = 1626.06$ (19) Å³
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.09$ mm⁻¹
 $T = 296$ K
 $0.25 \times 0.12 \times 0.10$ mm

Data collection

Bruker SMART APEXII CCD
area-detector diffractometer
Absorption correction: multi-scan
(SADABS; Bruker, 2005)
 $T_{\min} = 0.978$, $T_{\max} = 0.991$

11192 measured reflections
2872 independent reflections
918 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.097$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.058$
 $wR(F^2) = 0.133$
 $S = 0.94$
2872 reflections

219 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.14$ e Å⁻³
 $\Delta\rho_{\min} = -0.13$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{O3}-\text{H3}\cdots\text{N1}$	0.82	1.88	2.609 (3)	147
$\text{O2}-\text{H2}\cdots\text{O1}^i$	0.82	1.80	2.613 (2)	170

Symmetry code: (i) $-x + 2, -y + 1, -z + 2$.

Data collection: APEX2 (Bruker, 2005); cell refinement: SAINT (Bruker, 2005); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL and PLATON (Spek, 2009).

HK thanks PNU for financial support. MNT thanks GC University of Sargodha, Pakistan, for the research facility.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SU2388).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
Bernstein, J., Davis, R. E., Shimon, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
Bruker (2005). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
Kargar, H., Kia, R., Abbasian, S. & Tahir, M. N. (2012). *Acta Cryst.* **E68**, m182.
Kargar, H., Kia, R., Pahlavani, E. & Tahir, M. N. (2011). *Acta Cryst.* **E67**, o614.
Kia, R., Kargar, H., Tahir, M. N. & Kianoosh, F. (2010). *Acta Cryst.* **E66**, o2296.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

supporting information

Acta Cryst. (2012). E68, o1036 [https://doi.org/10.1107/S160053681200997X]

(E)-4-[(4-Diethylamino-2-hydroxybenzylidene)amino]benzoic acid**Hadi Kargar, Zahra Sharafi, Reza Kia and Muhammad Nawaz Tahir****S1. Comment**

In continuation of our work on the crystal structure analysis of Schiff base ligands (Kargar *et al.*, 2011, 2012; Kia *et al.*, 2010), we determined the crystal structure of the title compound.

The asymmetric unit of the title compound, Fig. 1, comprises a potential bidentate N,O-donor Schiff base ligand. The bond lengths (Allen *et al.*, 1987) and angles are within the normal ranges. The intramolecular O3—H3···N1 hydrogen bond (Table 1) makes an S(6) ring motif (Bernstein *et al.*, 1995). The dihedral angle between the benzene rings is 12.25 (19)°, and the molecule has an E conformation about the C8=N1 bond. One of the ethyl groups was disordered over two position with a refined site occupancy ratio of 0.55 (1)/0.45 (1).


In the crystal, pairs of O—H···O hydrogen bonds (Table 1) link molecules into inversion dimers with an $R^2_2(8)$ ring motif (Fig. 2).

S2. Experimental


The title compound was synthesized by adding 4-diethylaminosalicylaldehyde (2 mmol) to a solution of 4-carboxyaniline (2 mmol) in ethanol (30 ml). The mixture was refluxed with stirring for 30 min. The resultant solution was filtered. Pale-yellow single crystals of the title compound, suitable for X-ray structure analysis, were obtained by recrystallization from ethanol, by slow evaporation of the solvents at room temperature over several days.

S3. Refinement

The O-bound hydrogen atoms were located in a difference Fourier map and constrained to ride on the parent atoms with $U_{\text{iso}}(\text{H}) = 1.5 U_{\text{eq}}(\text{O})$. The rest of the hydrogen atoms were included in calculated positions and treated as riding atoms: C—H = 0.93, 0.96 and 0.97 Å for CH, CH₃ and CH₂ H atoms, respectively, with $U_{\text{iso}}(\text{H}) = k \times U_{\text{eq}}(\text{C})$, where $k = 1.5$ for CH₃ H atoms, and = 1.2 for other H atoms. A rotating group model was applied to the methyl group. One of the ethyl groups was disordered over two position with a refined site occupancy ratio of 0.55 (1)/0.45 (1). Since the crystal was very small and not optimal and did not diffract significantly, the ratio of observed to unique reflections is only 32%.

**Figure 1**

The ORTEP plot of the title compound, showing 40% probability displacement ellipsoids and the atomic numbering. The open bonds show the minor component of the disordered ethyl group. The dashed lines show the O-H...N hydrogen bond (see Table 1 for details).

**Figure 2**

The crystal packing diagram of the title compound viewed along the *b*-axis, showing the inversion dimers with an $R^2_2(8)$ ring motif. Only the hydrogen atoms involved the hydrogen bonding are shown - see Table 1 for details.

(E)-4-[(4-Diethylamino-2-hydroxybenzylidene)amino]benzoic acid*Crystal data*C₁₈H₂₀N₂O₃ $M_r = 312.36$ Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

 $a = 12.4216$ (8) Å $b = 8.1511$ (6) Å $c = 16.0820$ (11) Å $\beta = 93.001$ (3)° $V = 1626.06$ (19) Å³ $Z = 4$ $F(000) = 664$ $D_x = 1.276$ Mg m⁻³Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 3270 reflections

 $\theta = 2.8$ – 28.8 ° $\mu = 0.09$ mm⁻¹ $T = 296$ K

Block, pale-yellow

 $0.25 \times 0.12 \times 0.10$ mm*Data collection*Bruker SMART APEXII CCD area-detector
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 φ and ω scansAbsorption correction: multi-scan
(*SADABS*; Bruker, 2005) $T_{\min} = 0.978$, $T_{\max} = 0.991$

11192 measured reflections

2872 independent reflections

918 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.097$ $\theta_{\max} = 25.0$ °, $\theta_{\min} = 2.5$ ° $h = -14 \rightarrow 12$ $k = -9 \rightarrow 9$ $l = -19 \rightarrow 19$ *Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.058$ $wR(F^2) = 0.133$ $S = 0.94$

2872 reflections

219 parameters

0 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0393P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} < 0.001$ $\Delta\rho_{\max} = 0.14$ e Å⁻³ $\Delta\rho_{\min} = -0.13$ e Å⁻³*Special details*

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
C1	0.9492 (3)	0.4842 (5)	0.8861 (2)	0.0584 (12)	
C2	0.9076 (3)	0.4702 (5)	0.7993 (2)	0.0508 (11)	
C3	0.8265 (3)	0.3620 (5)	0.7757 (2)	0.0684 (13)	
H18	0.7990	0.2917	0.8150	0.082*	

C4	0.7858 (3)	0.3574 (5)	0.6946 (2)	0.0703 (13)	
H17	0.7299	0.2852	0.6801	0.084*	
C5	0.8256 (3)	0.4564 (5)	0.6347 (2)	0.0541 (12)	
C6	0.9084 (3)	0.5642 (5)	0.6578 (2)	0.0657 (13)	
H10	0.9371	0.6327	0.6184	0.079*	
C7	0.9481 (3)	0.5692 (5)	0.7398 (2)	0.0633 (12)	
H11	1.0036	0.6416	0.7547	0.076*	
C8	0.8135 (3)	0.5152 (5)	0.4913 (2)	0.0627 (13)	
H7	0.8806	0.5655	0.4982	0.075*	
C9	0.7599 (3)	0.5179 (5)	0.4096 (2)	0.0507 (11)	
C10	0.8031 (3)	0.5970 (5)	0.3437 (2)	0.0644 (12)	
H5	0.8700	0.6476	0.3520	0.077*	
C11	0.7519 (3)	0.6043 (5)	0.2663 (2)	0.0627 (12)	
H4	0.7842	0.6589	0.2234	0.075*	
C12	0.6503 (4)	0.5293 (5)	0.2515 (2)	0.0655 (13)	
C13	0.6062 (3)	0.4454 (5)	0.3171 (2)	0.0696 (14)	
H15	0.5408	0.3909	0.3083	0.084*	
C14	0.6582 (4)	0.4422 (5)	0.3949 (2)	0.0614 (12)	
C15	0.6378 (3)	0.6430 (6)	0.1074 (2)	0.0862 (15)	
H2A	0.6707	0.7408	0.1319	0.103*	
H2B	0.5769	0.6770	0.0713	0.103*	
C16	0.7176 (4)	0.5578 (5)	0.0570 (3)	0.1024 (17)	
H1A	0.7796	0.5285	0.0920	0.154*	
H1B	0.7391	0.6293	0.0134	0.154*	
H1C	0.6856	0.4604	0.0329	0.154*	
C17	0.4751 (11)	0.5013 (14)	0.1607 (7)	0.089 (5)	0.549 (11)
H17A	0.4373	0.5097	0.2118	0.107*	0.549 (11)
H17B	0.4407	0.5708	0.1182	0.107*	0.549 (11)
C18	0.4858 (16)	0.3342 (18)	0.1334 (11)	0.111 (6)	0.549 (11)
H18A	0.4156	0.2891	0.1200	0.167*	0.549 (11)
H18B	0.5212	0.2706	0.1771	0.167*	0.549 (11)
H18C	0.5277	0.3313	0.0850	0.167*	0.549 (11)
C17A	0.5142 (16)	0.405 (2)	0.1518 (10)	0.058 (5)	0.451 (11)
H17C	0.5183	0.3745	0.0937	0.070*	0.451 (11)
H17D	0.5286	0.3077	0.1855	0.070*	0.451 (11)
C18A	0.40179 (14)	0.4703 (4)	0.16718 (11)	0.075 (4)	0.451 (11)
H18D	0.3490	0.3878	0.1523	0.112*	0.451 (11)
H18E	0.3880	0.5665	0.1339	0.112*	0.451 (11)
H18F	0.3977	0.4977	0.2250	0.112*	0.451 (11)
N1	0.77425 (12)	0.44793 (19)	0.55449 (9)	0.0639 (11)	
N2	0.59881 (15)	0.5375 (3)	0.17449 (16)	0.0925 (14)	
O1	1.00875 (12)	0.59948 (19)	0.90912 (8)	0.0827 (10)	
O2	0.91877 (12)	0.37246 (19)	0.93606 (10)	0.0889 (10)	
H2	0.9476	0.3876	0.9825	0.133*	
O3	0.6110 (2)	0.3604 (4)	0.45575 (15)	0.0883 (10)	
H3	0.6443	0.3772	0.5003	0.132*	

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.061 (3)	0.068 (3)	0.045 (3)	0.010 (3)	-0.004 (2)	0.005 (2)
C2	0.049 (3)	0.055 (3)	0.047 (3)	0.003 (2)	-0.007 (2)	-0.001 (2)
C3	0.078 (3)	0.078 (4)	0.049 (3)	-0.012 (3)	-0.002 (2)	0.009 (2)
C4	0.078 (3)	0.078 (4)	0.053 (3)	-0.026 (3)	-0.010 (3)	0.005 (3)
C5	0.065 (3)	0.053 (3)	0.044 (2)	0.003 (2)	-0.005 (2)	-0.004 (2)
C6	0.078 (4)	0.072 (3)	0.046 (3)	-0.016 (3)	-0.001 (2)	0.005 (2)
C7	0.068 (3)	0.066 (3)	0.055 (3)	-0.016 (2)	-0.009 (2)	0.000 (2)
C8	0.069 (3)	0.058 (3)	0.060 (3)	0.004 (2)	-0.007 (3)	-0.013 (2)
C9	0.056 (3)	0.051 (3)	0.044 (3)	0.000 (2)	-0.004 (2)	-0.003 (2)
C10	0.064 (3)	0.076 (3)	0.054 (3)	-0.005 (3)	0.004 (2)	0.003 (3)
C11	0.068 (3)	0.076 (3)	0.043 (3)	-0.010 (3)	-0.005 (2)	0.008 (2)
C12	0.064 (3)	0.081 (4)	0.050 (3)	-0.011 (3)	-0.008 (3)	0.006 (3)
C13	0.073 (3)	0.085 (4)	0.050 (3)	-0.024 (3)	-0.011 (3)	0.007 (3)
C14	0.077 (4)	0.058 (3)	0.049 (3)	-0.011 (3)	0.006 (3)	0.009 (2)
C15	0.076 (4)	0.121 (4)	0.060 (3)	0.000 (3)	-0.011 (3)	0.016 (3)
C16	0.111 (5)	0.116 (5)	0.079 (3)	0.000 (3)	-0.002 (3)	-0.005 (3)
C17	0.112 (15)	0.077 (10)	0.078 (7)	0.023 (9)	0.002 (7)	0.012 (7)
C18	0.138 (15)	0.088 (14)	0.109 (13)	-0.014 (10)	0.007 (9)	-0.011 (9)
C17A	0.076 (11)	0.055 (14)	0.040 (7)	0.008 (10)	-0.024 (7)	-0.009 (9)
C18A	0.034 (7)	0.100 (11)	0.090 (8)	0.013 (6)	0.000 (6)	0.013 (7)
N1	0.080 (3)	0.066 (3)	0.045 (2)	-0.0010 (19)	-0.007 (2)	0.0019 (19)
N2	0.071 (3)	0.155 (4)	0.050 (2)	-0.037 (3)	-0.015 (2)	0.022 (3)
O1	0.097 (2)	0.089 (2)	0.0590 (19)	-0.0284 (19)	-0.0243 (16)	0.0038 (17)
O2	0.124 (3)	0.090 (2)	0.0495 (18)	-0.023 (2)	-0.0197 (17)	0.0063 (17)
O3	0.105 (3)	0.105 (2)	0.0537 (18)	-0.042 (2)	-0.0093 (17)	0.0086 (19)

Geometric parameters (Å, °)

C1—O1	1.240 (4)	C13—H15	0.9300
C1—O2	1.284 (4)	C14—O3	1.344 (4)
C1—C2	1.468 (4)	C15—N2	1.481 (4)
C2—C7	1.368 (4)	C15—C16	1.485 (5)
C2—C3	1.377 (4)	C15—H2A	0.9700
C3—C4	1.374 (4)	C15—H2B	0.9700
C3—H18	0.9300	C16—H1A	0.9600
C4—C5	1.369 (4)	C16—H1B	0.9600
C4—H17	0.9300	C16—H1C	0.9600
C5—C6	1.389 (4)	C17—C18	1.440 (19)
C5—N1	1.410 (4)	C17—N2	1.569 (14)
C6—C7	1.383 (5)	C17—H17A	0.9700
C6—H10	0.9300	C17—H17B	0.9700
C7—H11	0.9300	C18—H18A	0.9600
C8—N1	1.274 (4)	C18—H18B	0.9600
C8—C9	1.441 (4)	C18—H18C	0.9600
C8—H7	0.9300	C17A—C18A	1.53 (2)

C9—C10	1.374 (4)	C17A—N2	1.538 (15)
C9—C14	1.414 (5)	C17A—H17C	0.9700
C10—C11	1.370 (4)	C17A—H17D	0.9700
C10—H5	0.9300	C18A—H18D	0.9600
C11—C12	1.411 (5)	C18A—H18E	0.9600
C11—H4	0.9300	C18A—H18F	0.9600
C12—N2	1.365 (4)	O2—H2	0.8200
C12—C13	1.394 (5)	O3—H3	0.8200
C13—C14	1.379 (5)		
O1—C1—O2	122.8 (3)	O3—C14—C9	121.0 (4)
O1—C1—C2	121.2 (3)	C13—C14—C9	121.1 (4)
O2—C1—C2	116.0 (4)	N2—C15—C16	112.3 (4)
C7—C2—C3	118.4 (3)	N2—C15—H2A	109.1
C7—C2—C1	119.7 (4)	C16—C15—H2A	109.1
C3—C2—C1	121.9 (4)	N2—C15—H2B	109.1
C4—C3—C2	120.4 (4)	C16—C15—H2B	109.1
C4—C3—H18	119.8	H2A—C15—H2B	107.9
C2—C3—H18	119.8	C15—C16—H1A	109.5
C5—C4—C3	121.6 (4)	C15—C16—H1B	109.5
C5—C4—H17	119.2	H1A—C16—H1B	109.5
C3—C4—H17	119.2	C15—C16—H1C	109.5
C4—C5—C6	118.3 (3)	H1A—C16—H1C	109.5
C4—C5—N1	116.9 (4)	H1B—C16—H1C	109.5
C6—C5—N1	124.7 (4)	C18—C17—N2	96.6 (12)
C7—C6—C5	119.7 (4)	C18—C17—H17A	112.4
C7—C6—H10	120.1	N2—C17—H17A	112.4
C5—C6—H10	120.1	C18—C17—H17B	112.4
C2—C7—C6	121.6 (4)	N2—C17—H17B	112.4
C2—C7—H11	119.2	H17A—C17—H17B	110.0
C6—C7—H11	119.2	C18A—C17A—N2	109.6 (11)
N1—C8—C9	123.8 (4)	C18A—C17A—H17C	109.8
N1—C8—H7	118.1	N2—C17A—H17C	109.8
C9—C8—H7	118.1	C18A—C17A—H17D	109.8
C10—C9—C14	117.2 (4)	N2—C17A—H17D	109.8
C10—C9—C8	122.0 (4)	H17C—C17A—H17D	108.2
C14—C9—C8	120.8 (4)	C17A—C18A—H18D	109.5
C11—C10—C9	122.7 (4)	C17A—C18A—H18E	109.5
C11—C10—H5	118.7	H18D—C18A—H18E	109.5
C9—C10—H5	118.7	C17A—C18A—H18F	109.5
C10—C11—C12	120.2 (4)	H18D—C18A—H18F	109.5
C10—C11—H4	119.9	H18E—C18A—H18F	109.5
C12—C11—H4	119.9	C8—N1—C5	122.4 (3)
N2—C12—C13	121.8 (4)	C12—N2—C15	122.2 (3)
N2—C12—C11	120.3 (4)	C12—N2—C17A	117.5 (7)
C13—C12—C11	118.0 (4)	C15—N2—C17A	118.8 (7)
C14—C13—C12	120.8 (4)	C12—N2—C17	121.8 (5)
C14—C13—H15	119.6	C15—N2—C17	111.2 (5)

C12—C13—H15	119.6	C1—O2—H2	109.5
O3—C14—C13	117.9 (4)	C14—O3—H3	109.5
O1—C1—C2—C7	10.7 (6)	C12—C13—C14—C9	-2.7 (6)
O2—C1—C2—C7	-170.5 (3)	C10—C9—C14—O3	179.1 (4)
O1—C1—C2—C3	-167.8 (4)	C8—C9—C14—O3	-2.6 (6)
O2—C1—C2—C3	11.1 (6)	C10—C9—C14—C13	1.2 (6)
C7—C2—C3—C4	-1.5 (6)	C8—C9—C14—C13	179.5 (4)
C1—C2—C3—C4	176.9 (4)	C9—C8—N1—C5	-175.9 (3)
C2—C3—C4—C5	1.4 (6)	C4—C5—N1—C8	-169.4 (3)
C3—C4—C5—C6	-0.5 (6)	C6—C5—N1—C8	14.6 (5)
C3—C4—C5—N1	-176.7 (3)	C13—C12—N2—C15	172.2 (4)
C4—C5—C6—C7	-0.1 (6)	C11—C12—N2—C15	-9.0 (5)
N1—C5—C6—C7	175.8 (3)	C13—C12—N2—C17A	-21.8 (10)
C3—C2—C7—C6	0.9 (6)	C11—C12—N2—C17A	157.0 (9)
C1—C2—C7—C6	-177.6 (4)	C13—C12—N2—C17	18.6 (7)
C5—C6—C7—C2	-0.1 (6)	C11—C12—N2—C17	-162.5 (6)
N1—C8—C9—C10	177.6 (4)	C16—C15—N2—C12	88.2 (4)
N1—C8—C9—C14	-0.6 (6)	C16—C15—N2—C17A	-77.6 (10)
C14—C9—C10—C11	0.0 (6)	C16—C15—N2—C17	-115.7 (5)
C8—C9—C10—C11	-178.2 (4)	C18A—C17A—N2—C12	97.2 (10)
C9—C10—C11—C12	0.2 (6)	C18A—C17A—N2—C15	-96.3 (11)
C10—C11—C12—N2	179.6 (4)	C18A—C17A—N2—C17	-9.9 (8)
C10—C11—C12—C13	-1.5 (6)	C18—C17—N2—C12	-98.4 (11)
N2—C12—C13—C14	-178.3 (4)	C18—C17—N2—C15	105.4 (10)
C11—C12—C13—C14	2.8 (6)	C18—C17—N2—C17A	-4.7 (18)
C12—C13—C14—O3	179.4 (4)		

Hydrogen-bond geometry (Å, °)

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
O3—H3...N1	0.82	1.88	2.609 (3)	147
O2—H2...O1 ⁱ	0.82	1.80	2.613 (2)	170

Symmetry code: (i) $-x+2, -y+1, -z+2$.