

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Dimethyl 1-(3-hydroxy-2-iodo-1-phenylpropyl)-1*H*-1,2,3-triazole-4,5-dicarboxylate

 Hoong-Kun Fun,^{a,*‡} Tze Shyang Chia,^a Sivasubramanian Archana,^b Murugan Dinesh^b and Alagusundaram Ponnuswamy^b
^aX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, and ^bDepartment of Organic Chemistry, School of Chemistry, Madurai Kamaraj University, Madurai 625 021, Tamil Nadu, India
Correspondence e-mail: hkfun@usm.my


Received 23 April 2012; accepted 23 April 2012

 Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.001$ Å; R factor = 0.015; wR factor = 0.038; data-to-parameter ratio = 26.6.

In the title compound, $\text{C}_{15}\text{H}_{16}\text{IN}_3\text{O}_5$, the central triazole ring is essentially planar (r.m.s deviation = 0.0034 Å) and makes a dihedral angle of 70.14 (5)° with the pendant benzene ring. The mean planes of the two methoxycarbonyl groups make dihedral angles of 22.52 (7) and 40.93 (4)° with the triazole ring. In the crystal, inversion dimers linked by pairs of $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds generate $R_2^2(18)$ loops. The dimers are linked by $\text{C}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\text{N}$ interactions into sheets lying parallel to the ac plane.

Related literature

For background to the industrial applications of 1,2,3-triazoles, see: Wamhoff (1984). For the stability of the temperature controller used in the data collection, see: Cosier & Glazer (1986).



Experimental

Crystal data

$\text{C}_{15}\text{H}_{16}\text{IN}_3\text{O}_5$
 $M_r = 445.21$
 Triclinic, $P\bar{1}$
 $a = 8.0504$ (3) Å
 $b = 9.6941$ (4) Å
 $c = 11.1893$ (4) Å
 $\alpha = 106.426$ (1)°
 $\beta = 91.798$ (1)°
 $\gamma = 98.606$ (1)°
 $V = 825.65$ (5) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 1.97$ mm⁻¹
 $T = 100$ K
 $0.31 \times 0.24 \times 0.14$ mm

Data collection

Bruker APEX DUO CCD diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2009)
 $T_{\min} = 0.579$, $T_{\max} = 0.763$
 21676 measured reflections
 5934 independent reflections
 5827 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.015$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.015$
 $wR(F^2) = 0.038$
 $S = 1.10$
 5934 reflections
 223 parameters
 H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.49$ e Å⁻³
 $\Delta\rho_{\min} = -0.92$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O5}-\text{H1O5}\cdots\text{O3}^{\text{i}}$	0.74 (2)	2.22 (2)	2.9213 (11)	159 (2)
$\text{C9}-\text{H9A}\cdots\text{O1}^{\text{ii}}$	0.97	2.57	3.3595 (13)	139
$\text{C13}-\text{H13B}\cdots\text{O3}^{\text{iii}}$	0.96	2.48	3.4375 (14)	174
$\text{C13}-\text{H13C}\cdots\text{O5}^{\text{i}}$	0.96	2.58	3.4247 (14)	147
$\text{C15}-\text{H15C}\cdots\text{N2}^{\text{iv}}$	0.96	2.62	3.4903 (14)	151

Symmetry codes: (i) $-x, -y + 2, -z$; (ii) $x - 1, y, z$; (iii) $-x + 1, -y + 2, -z$; (iv) $-x + 1, -y + 2, -z + 1$.

Data collection: APEX2 (Bruker, 2009); cell refinement: SAINT (Bruker, 2009); data reduction: SAINT; program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL and PLATON (Spek, 2009).

HKF and TSC thank Universiti Sains Malaysia (USM) for the Research University Grant (1001/PFIZIK/811160). TSC also thanks the Malaysian Government and USM for the award of a research fellowship.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB6752).

References

- Bruker (2009). SADABS, APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Cosier, J. & Glazer, A. M. (1986). *J. Appl. Cryst.* **19**, 105–107.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
 Wamhoff, H. (1984). *Comprehensive Heterocyclic Chemistry*, Vol. 5, edited by A. R. Katritzky & C. W. Rees, pp. 669–732. Oxford: Pergamon Press.

‡ Thomson Reuters ResearcherID: A-3561-2009.

supporting information

Acta Cryst. (2012). E68, o1559 [doi:10.1107/S1600536812018193]

Dimethyl 1-(3-hydroxy-2-iodo-1-phenylpropyl)-1*H*-1,2,3-triazole-4,5-dicarboxylate

Hoong-Kun Fun, Tze Shyang Chia, Sivasubramanian Archana, Murugan Dinesh and Alagusundaram Ponnuswamy

S1. Comment

1,2,3-Triazoles are important in a wide range of industrial applications such as agrochemicals, corrosion inhibitors, dyes, optical brighteners as well as biologically active agents (Wamhoff *et al.*, 1984). As part of our studies in this area, we now describe the synthesis and structure of the title compound, (I) (Fig. 1).

The central triazole ring (N1–N3/C10/C11) is essentially planar [r.m.s deviation = 0.0034 Å] and makes a dihedral angle of 70.14 (5)° with the terminal benzene ring (C1–C6). The mean planes of the two methyl carboxylate groups (O1/O2/C14/C15 with maximum deviation = 0.0199 (6) Å at atom C14 and O3/O4/C12/C13 with maximum deviation = 0.0060 (6) Å at atom C12) make dihedral angles of 22.52 (7) and 40.93 (4)°, respectively with the triazole ring.


In the crystal (Fig. 2), the molecules are linked by O5—H1O5···O3, C9—H9A···O1, C13—H13B···O3, C13—H13C···O5 and C15—H15C···N2 hydrogen bonds (Table 1) into sheets parallel to *ac* plane.

S2. Experimental


A mixture of 3-azido-2-iodo-3-phenylpropan-1-ol (0.3 g, 0.99 mmol) and dimethyl but-2-ynedioate (0.14 g, 0.99 mmol) was heated to reflux in toluene for 3 h. The solvent was evaporated under reduced pressure to afford the crude reaction mass which was then subjected to column chromatography using silica gel (60–120 mesh) as the stationary phase and petroleum ether:ethyl acetate (80:20) as the mobile phase to give colourless blocks of (I) by slow recrystallization from the eluant. Yield: 0.35 g (81%); *M.p.*: 129–130 °C.

S3. Refinement

Atom H1O5 was located from difference fourier map and refined freely [O—H = 0.74 (2) Å]. The remaining H atoms were positioned geometrically [C—H = 0.93, 0.96, 0.97 and 0.98 Å] and refined using a riding model with $U_{\text{iso}}(\text{H}) = 1.2$ or $1.5 U_{\text{eq}}(\text{C})$. A rotating group model was applied to the methyl groups. Sixteen outliers (5 - 1 4), (-4 1 6), (-4 2 5), (-4 3 4), (-2 3 0), (-4 1 0), (-4 4 4), (-2 - 1 3), (2 1 2), (-2 2 2), (-1 4 0), (-4 - 2 5), (5 0 4), (3 - 1 3), (-3 1 1) and (2 - 4 3) were omitted.

**Figure 1**

The molecular structure of the title compound with 50% probability displacement ellipsoids.

**Figure 2**

The crystal packing of the title compound. The dashed lines represent the hydrogen bonds. For clarity sake, hydrogen atoms not involved in hydrogen bonding have been omitted.

Dimethyl 1-(3-hydroxy-2-iodo-1-phenylpropyl)-1*H*-1,2,3-triazole-4,5-dicarboxylate

Crystal data

C₁₅H₁₆IN₃O₅ $M_r = 445.21$ Triclinic, $P\bar{1}$

Hall symbol: -P 1

 $a = 8.0504 (3) \text{ \AA}$ $b = 9.6941 (4) \text{ \AA}$ $c = 11.1893 (4) \text{ \AA}$ $\alpha = 106.426 (1)^\circ$ $\beta = 91.798 (1)^\circ$ $\gamma = 98.606 (1)^\circ$ $V = 825.65 (5) \text{ \AA}^3$ $Z = 2$ $F(000) = 440$ $D_x = 1.791 \text{ Mg m}^{-3}$ Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 9309 reflections

 $\theta = 2.2\text{--}32.6^\circ$ $\mu = 1.97 \text{ mm}^{-1}$ $T = 100 \text{ K}$

Block, colourless

 $0.31 \times 0.24 \times 0.14 \text{ mm}$

Data collection

Bruker APEX DUO CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 φ and ω scans

Absorption correction: multi-scan

(SADABS; Bruker, 2009)

 $T_{\min} = 0.579$, $T_{\max} = 0.763$

21676 measured reflections

5934 independent reflections

5827 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.015$ $\theta_{\max} = 32.6^\circ$, $\theta_{\min} = 1.9^\circ$ $h = -12 \rightarrow 11$ $k = -14 \rightarrow 14$ $l = -16 \rightarrow 16$

Refinement

Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.015$ $wR(F^2) = 0.038$ $S = 1.10$

5934 reflections

223 parameters

0 restraints

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

 $w = 1/[\sigma^2(F_o^2) + (0.0171P)^2 + 0.3086P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} = 0.002$ $\Delta\rho_{\max} = 0.49 \text{ e \AA}^{-3}$ $\Delta\rho_{\min} = -0.92 \text{ e \AA}^{-3}$

Special details

Experimental. The crystal was placed in the cold stream of an Oxford Cryosystems Cobra open-flow nitrogen cryostat (Cosier & Glazer, 1986) operating at 100.0 (1) K.

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
I1	-0.382668 (7)	0.637377 (7)	0.141512 (5)	0.01603 (2)

O1	0.55901 (10)	1.17479 (8)	0.31744 (7)	0.01765 (13)
O2	0.46348 (10)	1.25049 (8)	0.50897 (7)	0.01713 (13)
O3	0.29674 (10)	1.04776 (7)	0.09582 (6)	0.01498 (12)
O4	0.26687 (9)	0.80305 (7)	0.05704 (6)	0.01300 (12)
O5	-0.09851 (11)	0.94656 (9)	0.12543 (7)	0.01876 (14)
N1	0.13027 (10)	0.84350 (8)	0.29530 (7)	0.01084 (12)
N2	0.13613 (11)	0.87602 (9)	0.42137 (7)	0.01341 (13)
N3	0.25260 (11)	0.99050 (9)	0.46809 (7)	0.01367 (14)
C1	0.11822 (13)	0.50289 (10)	0.15751 (9)	0.01454 (15)
H1A	0.1649	0.5371	0.0942	0.017*
C2	0.15365 (14)	0.37227 (10)	0.17255 (10)	0.01754 (17)
H2A	0.2246	0.3202	0.1202	0.021*
C3	0.08244 (14)	0.31994 (10)	0.26621 (10)	0.01701 (17)
H3A	0.1065	0.2332	0.2771	0.020*
C4	-0.02472 (14)	0.39767 (10)	0.34358 (9)	0.01666 (17)
H4A	-0.0740	0.3616	0.4051	0.020*
C5	-0.05871 (13)	0.52943 (10)	0.32951 (9)	0.01423 (15)
H5A	-0.1295	0.5814	0.3822	0.017*
C6	0.01325 (11)	0.58325 (9)	0.23655 (8)	0.01099 (14)
C7	-0.00868 (11)	0.72960 (9)	0.22038 (8)	0.01059 (13)
H7A	0.0028	0.7236	0.1322	0.013*
C8	-0.17309 (12)	0.78729 (9)	0.25651 (8)	0.01215 (14)
H8A	-0.1878	0.7938	0.3444	0.015*
C9	-0.16977 (13)	0.93812 (10)	0.23842 (9)	0.01566 (16)
H9A	-0.2837	0.9592	0.2369	0.019*
H9B	-0.1044	1.0111	0.3085	0.019*
C10	0.24390 (12)	0.94051 (9)	0.26041 (8)	0.01084 (14)
C11	0.32129 (12)	1.03422 (9)	0.37215 (8)	0.01193 (14)
C12	0.27066 (11)	0.93847 (9)	0.12878 (8)	0.01094 (14)
C13	0.30152 (14)	0.78729 (11)	-0.07305 (8)	0.01628 (16)
H13A	0.2970	0.6859	-0.1166	0.024*
H13B	0.4116	0.8397	-0.0761	0.024*
H13C	0.2187	0.8259	-0.1119	0.024*
C14	0.46100 (12)	1.15881 (10)	0.39366 (8)	0.01275 (14)
C15	0.60447 (14)	1.36969 (11)	0.54233 (10)	0.01921 (18)
H15A	0.5865	1.4391	0.6191	0.029*
H15B	0.6140	1.4164	0.4772	0.029*
H15C	0.7063	1.3326	0.5527	0.029*
H1O5	-0.166 (3)	0.934 (2)	0.0748 (18)	0.032 (5)*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
I1	0.01068 (3)	0.01859 (3)	0.01613 (3)	0.00087 (2)	-0.00104 (2)	0.00174 (2)
O1	0.0165 (3)	0.0192 (3)	0.0152 (3)	-0.0009 (3)	0.0032 (3)	0.0036 (2)
O2	0.0163 (3)	0.0158 (3)	0.0135 (3)	-0.0035 (3)	0.0017 (2)	-0.0018 (2)
O3	0.0186 (3)	0.0131 (3)	0.0134 (3)	0.0007 (2)	0.0005 (2)	0.0052 (2)
O4	0.0170 (3)	0.0114 (3)	0.0099 (3)	0.0024 (2)	0.0029 (2)	0.0018 (2)

O5	0.0203 (4)	0.0221 (3)	0.0174 (3)	0.0057 (3)	0.0010 (3)	0.0101 (3)
N1	0.0121 (3)	0.0110 (3)	0.0088 (3)	0.0010 (2)	0.0003 (2)	0.0024 (2)
N2	0.0157 (4)	0.0143 (3)	0.0089 (3)	0.0004 (3)	0.0004 (3)	0.0024 (2)
N3	0.0148 (4)	0.0140 (3)	0.0108 (3)	0.0000 (3)	0.0007 (3)	0.0024 (2)
C1	0.0162 (4)	0.0122 (3)	0.0152 (4)	0.0034 (3)	0.0042 (3)	0.0033 (3)
C2	0.0192 (4)	0.0130 (4)	0.0205 (4)	0.0052 (3)	0.0041 (3)	0.0033 (3)
C3	0.0191 (4)	0.0117 (3)	0.0202 (4)	0.0019 (3)	-0.0010 (3)	0.0051 (3)
C4	0.0205 (4)	0.0150 (4)	0.0151 (4)	0.0012 (3)	0.0009 (3)	0.0064 (3)
C5	0.0159 (4)	0.0141 (3)	0.0130 (3)	0.0023 (3)	0.0024 (3)	0.0044 (3)
C6	0.0111 (4)	0.0102 (3)	0.0107 (3)	0.0009 (3)	0.0001 (3)	0.0021 (3)
C7	0.0104 (3)	0.0103 (3)	0.0099 (3)	0.0009 (3)	0.0001 (3)	0.0017 (3)
C8	0.0117 (4)	0.0123 (3)	0.0114 (3)	0.0022 (3)	0.0005 (3)	0.0017 (3)
C9	0.0172 (4)	0.0133 (3)	0.0172 (4)	0.0052 (3)	0.0023 (3)	0.0041 (3)
C10	0.0118 (4)	0.0108 (3)	0.0098 (3)	0.0017 (3)	0.0007 (3)	0.0029 (3)
C11	0.0124 (4)	0.0117 (3)	0.0103 (3)	0.0009 (3)	0.0005 (3)	0.0017 (3)
C12	0.0102 (3)	0.0122 (3)	0.0095 (3)	0.0011 (3)	0.0003 (3)	0.0022 (3)
C13	0.0218 (5)	0.0167 (4)	0.0094 (3)	0.0030 (3)	0.0033 (3)	0.0020 (3)
C14	0.0126 (4)	0.0123 (3)	0.0120 (3)	0.0010 (3)	-0.0006 (3)	0.0020 (3)
C15	0.0171 (4)	0.0160 (4)	0.0197 (4)	-0.0040 (3)	-0.0024 (3)	0.0013 (3)

Geometric parameters (Å, °)

I1—C8	2.1649 (9)	C4—C5	1.3953 (13)
O1—C14	1.2059 (11)	C4—H4A	0.9300
O2—C14	1.3414 (11)	C5—C6	1.3942 (12)
O2—C15	1.4491 (12)	C5—H5A	0.9300
O3—C12	1.2091 (11)	C6—C7	1.5148 (12)
O4—C12	1.3264 (10)	C7—C8	1.5347 (13)
O4—C13	1.4602 (11)	C7—H7A	0.9800
O5—C9	1.4227 (12)	C8—C9	1.5283 (13)
O5—H1O5	0.74 (2)	C8—H8A	0.9800
N1—N2	1.3534 (10)	C9—H9A	0.9700
N1—C10	1.3588 (11)	C9—H9B	0.9700
N1—C7	1.4904 (12)	C10—C11	1.3842 (12)
N2—N3	1.3091 (11)	C10—C12	1.4902 (12)
N3—C11	1.3646 (11)	C11—C14	1.4811 (13)
C1—C2	1.3923 (13)	C13—H13A	0.9600
C1—C6	1.3991 (12)	C13—H13B	0.9600
C1—H1A	0.9300	C13—H13C	0.9600
C2—C3	1.3913 (14)	C15—H15A	0.9600
C2—H2A	0.9300	C15—H15B	0.9600
C3—C4	1.3907 (14)	C15—H15C	0.9600
C3—H3A	0.9300		
C14—O2—C15	114.80 (8)	C7—C8—I1	109.17 (6)
C12—O4—C13	115.88 (7)	C9—C8—H8A	109.1
C9—O5—H1O5	110.5 (15)	C7—C8—H8A	109.1
N2—N1—C10	110.40 (7)	I1—C8—H8A	109.1

N2—N1—C7	118.32 (7)	O5—C9—C8	111.51 (7)
C10—N1—C7	130.48 (7)	O5—C9—H9A	109.3
N3—N2—N1	108.01 (7)	C8—C9—H9A	109.3
N2—N3—C11	108.72 (7)	O5—C9—H9B	109.3
C2—C1—C6	120.75 (8)	C8—C9—H9B	109.3
C2—C1—H1A	119.6	H9A—C9—H9B	108.0
C6—C1—H1A	119.6	N1—C10—C11	104.27 (7)
C3—C2—C1	119.75 (9)	N1—C10—C12	124.96 (8)
C3—C2—H2A	120.1	C11—C10—C12	130.77 (8)
C1—C2—H2A	120.1	N3—C11—C10	108.59 (8)
C4—C3—C2	119.87 (9)	N3—C11—C14	122.16 (8)
C4—C3—H3A	120.1	C10—C11—C14	129.19 (8)
C2—C3—H3A	120.1	O3—C12—O4	126.17 (8)
C3—C4—C5	120.40 (9)	O3—C12—C10	123.15 (8)
C3—C4—H4A	119.8	O4—C12—C10	110.64 (7)
C5—C4—H4A	119.8	O4—C13—H13A	109.5
C6—C5—C4	120.08 (9)	O4—C13—H13B	109.5
C6—C5—H5A	120.0	H13A—C13—H13B	109.5
C4—C5—H5A	120.0	O4—C13—H13C	109.5
C5—C6—C1	119.13 (8)	H13A—C13—H13C	109.5
C5—C6—C7	123.26 (8)	H13B—C13—H13C	109.5
C1—C6—C7	117.53 (8)	O1—C14—O2	124.91 (9)
N1—C7—C6	109.14 (7)	O1—C14—C11	124.26 (8)
N1—C7—C8	106.17 (7)	O2—C14—C11	110.83 (8)
C6—C7—C8	118.54 (7)	O2—C15—H15A	109.5
N1—C7—H7A	107.5	O2—C15—H15B	109.5
C6—C7—H7A	107.5	H15A—C15—H15B	109.5
C8—C7—H7A	107.5	O2—C15—H15C	109.5
C9—C8—C7	111.09 (7)	H15A—C15—H15C	109.5
C9—C8—I1	109.34 (6)	H15B—C15—H15C	109.5
C10—N1—N2—N3	-0.86 (10)	I1—C8—C9—O5	-78.79 (9)
C7—N1—N2—N3	-171.70 (8)	N2—N1—C10—C11	0.45 (10)
N1—N2—N3—C11	0.91 (10)	C7—N1—C10—C11	169.83 (9)
C6—C1—C2—C3	0.76 (16)	N2—N1—C10—C12	-179.45 (8)
C1—C2—C3—C4	0.53 (16)	C7—N1—C10—C12	-10.07 (15)
C2—C3—C4—C5	-1.28 (16)	N2—N3—C11—C10	-0.64 (11)
C3—C4—C5—C6	0.75 (15)	N2—N3—C11—C14	-178.05 (8)
C4—C5—C6—C1	0.53 (14)	N1—C10—C11—N3	0.11 (10)
C4—C5—C6—C7	-176.10 (9)	C12—C10—C11—N3	180.00 (9)
C2—C1—C6—C5	-1.29 (15)	N1—C10—C11—C14	177.28 (9)
C2—C1—C6—C7	175.54 (9)	C12—C10—C11—C14	-2.83 (17)
N2—N1—C7—C6	-66.54 (10)	C13—O4—C12—O3	1.45 (14)
C10—N1—C7—C6	124.78 (9)	C13—O4—C12—C10	-176.52 (8)
N2—N1—C7—C8	62.31 (9)	N1—C10—C12—O3	140.62 (10)
C10—N1—C7—C8	-106.37 (10)	C11—C10—C12—O3	-39.26 (16)
C5—C6—C7—N1	90.62 (10)	N1—C10—C12—O4	-41.34 (12)
C1—C6—C7—N1	-86.06 (10)	C11—C10—C12—O4	138.79 (10)

C5—C6—C7—C8	-31.01 (13)	C15—O2—C14—O1	-4.72 (14)
C1—C6—C7—C8	152.31 (8)	C15—O2—C14—C11	175.14 (8)
N1—C7—C8—C9	54.79 (9)	N3—C11—C14—O1	156.03 (10)
C6—C7—C8—C9	177.90 (8)	C10—C11—C14—O1	-20.80 (16)
N1—C7—C8—H1	175.44 (5)	N3—C11—C14—O2	-23.83 (13)
C6—C7—C8—H1	-61.45 (9)	C10—C11—C14—O2	159.34 (9)
C7—C8—C9—O5	41.77 (11)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
O5—H1O5...O3 ⁱ	0.74 (2)	2.22 (2)	2.9213 (11)	159 (2)
C9—H9A...O1 ⁱⁱ	0.97	2.57	3.3595 (13)	139
C13—H13B...O3 ⁱⁱⁱ	0.96	2.48	3.4375 (14)	174
C13—H13C...O5 ⁱ	0.96	2.58	3.4247 (14)	147
C15—H15C...N2 ^{iv}	0.96	2.62	3.4903 (14)	151

Symmetry codes: (i) $-x, -y+2, -z$; (ii) $x-1, y, z$; (iii) $-x+1, -y+2, -z$; (iv) $-x+1, -y+2, -z+1$.