

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

3-(3,4-Dihydroxyphenyl)-1-methoxy-1-oxopropan-2-aminium chloride

Ze-Feng Hou,^a Juan Feng,^b Jing-Tian Liu,^c Xiao-Li Zhen^a and Jian-Rong Han^{a*}

^aCollege of Sciences, Hebei University of Science & Technology, Shijiazhuang 050018, People's Republic of China, ^bCollege of Chemical & Pharmaceutical Engineering, Hebei University of Science & Technology, Shijiazhuang 050018, People's Republic of China, and ^cCollege of Chemistry and Chemical Engineering, Nanjing University of Technology, Nanjing 211816, People's Republic of China
Correspondence e-mail: han_jianrong@126.com


Received 13 April 2012; accepted 18 April 2012

Key indicators: single-crystal X-ray study; $T = 113$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.027; wR factor = 0.061; data-to-parameter ratio = 17.0.

In the title compound, $\text{C}_{10}\text{H}_{14}\text{NO}_4^+\cdot\text{Cl}^-$, the benzene ring makes a dihedral angle of $64.68(4)^\circ$ with the methylamino-propanoate unit, which is bonded to the catechol ring *via* a methylene C atom. A strong intramolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bond occurs. In the crystal, $\text{O}-\text{H}\cdots\text{O}$, $\text{N}-\text{H}\cdots\text{Cl}$ and $\text{O}-\text{H}\cdots\text{Cl}$ hydrogen bonds and weak $\text{C}-\text{H}\cdots\text{O}$ interactions link the molecules into a three-dimensional network.

Related literature

For medicinal applications of the title compound, see: Cooper *et al.* (1984). For a related structure, see: Naicker *et al.* (2012)



Experimental

Crystal data

 $\text{C}_{10}\text{H}_{14}\text{NO}_4^+\cdot\text{Cl}^-$ $M_r = 247.67$ Orthorhombic, $P2_12_12_1$ $a = 4.9969(15)$ Å $b = 14.498(4)$ Å $c = 16.109(5)$ Å $V = 1167.1(6)$ Å³ $Z = 4$ Mo $K\alpha$ radiation $\mu = 0.33$ mm⁻¹ $T = 113$ K $0.20 \times 0.18 \times 0.12$ mm

Data collection

Rigaku Saturn724 CCD diffractometer

Absorption correction: multi-scan (*CrystalClear*; Rigaku/MS, 2005) $T_{\min} = 0.938$, $T_{\max} = 0.962$

12293 measured reflections

2796 independent reflections

2054 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.051$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.027$ $wR(F^2) = 0.061$ $S = 0.93$

2796 reflections

164 parameters

H atoms treated by a mixture of independent and constrained refinement

 $\Delta\rho_{\max} = 0.18$ e Å⁻³ $\Delta\rho_{\min} = -0.23$ e Å⁻³

Absolute structure: Flack (1983),

1117 Friedel pairs

Flack parameter: $-0.03(5)$

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1C}\cdots\text{Cl1}^{\text{i}}$	0.96 (2)	2.29 (2)	3.209 (2)	161 (2)
$\text{N1}-\text{H1B}\cdots\text{Cl1}$	0.92 (2)	2.49 (2)	3.178 (2)	132 (1)
$\text{N1}-\text{H1A}\cdots\text{Cl1}^{\text{ii}}$	1.01 (2)	2.13 (2)	3.112 (2)	163 (2)
$\text{O2}-\text{H2}\cdots\text{Cl1}^{\text{iii}}$	0.84 (2)	2.26 (2)	3.086 (2)	167 (2)
$\text{O1}-\text{H1}\cdots\text{O2}$	0.83 (2)	2.27 (2)	2.698 (2)	113 (2)
$\text{O1}-\text{H1}\cdots\text{O3}^{\text{iv}}$	0.83 (2)	2.26 (2)	3.029 (2)	155 (2)
$\text{C8}-\text{H8}\cdots\text{O3}^{\text{v}}$	1.00	2.44	3.300 (2)	144
$\text{C10}-\text{H10C}\cdots\text{O1}^{\text{v}}$	0.98	2.47	3.137 (3)	125

Symmetry codes: (i) $x - \frac{1}{2}, -y + \frac{3}{2}, -z + 1$; (ii) $x - 1, y, z$; (iii) $-x + \frac{3}{2}, -y + 1, z + \frac{1}{2}$; (iv) $-x + 2, y - \frac{1}{2}, -z + \frac{3}{2}$; (v) $-x + 2, y + \frac{1}{2}, -z + \frac{3}{2}$.

Data collection: *CrystalClear* (Rigaku/MS, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

We are grateful for financial support from the Hebei Natural Science Foundation (No. B2010000836) and the Foundation of the Education Department of Hebei Province (grant No. ZD2010109)

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PV2536).

References

- Cooper, D. R., Marrel, C., Testa, B., van de Waterbeemd, H., Quinn, N., Jenner, P. & Marsden, C. D. (1984). *Clin. Neuropharmacol.* **7**, 89–98.
 Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
 Naicker, T., Govender, T., Kruger, H. G. & Maguire, G. E. M. (2012). *Acta Cryst.* **E68**, o177.
 Rigaku/MS (2005). *CrystalClear*. Rigaku/MS Inc., The Woodlands, Texas, USA.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supporting information

Acta Cryst. (2012). E68, o1516 [doi:10.1107/S160053681201728X]

3-(3,4-Dihydroxyphenyl)-1-methoxy-1-oxopropan-2-aminium chloride

Ze-Feng Hou, Juan Feng, Jing-Tian Liu, Xiao-Li Zhen and Jian-Rong Han

S1. Comment

A systemic or subcutaneous infusion of *L*-Dopa methyl ester to patients suffering from Parkinson's disease who experience response fluctuations may provide a means of maintaining mobility (Cooper *et al.*, 1984). In this article, we report the synthesis and crystal structure of *L*-Dopa methyl ester hydrochloride.

In the title compound (Fig. 1), benzene ring makes a dihedral angle of 64.68 (4)° with the methylaminopropanoate moiety (N1/C8–C10/O3/O4) bonded to the catechol ring *via* a methylene C7 atom. The torsion angles in the chain connected with the aromatic ring (C6/C7—C8/N1) and (C6/C7—C8/C9) are 71.39 (19) and -50.5 (2)°, respectively. The crystal packing is stabilized by strong intermolecular O—H···O, N—H···Cl and O—H···Cl hydrogen bonds and further consolidated by weak C—H···O interactions; all the interactions link the molecules into an infinite network (Table 1 and Fig. 2).


The bond distances and bond angles in the title compound are in agreement with the corresponding bond lengths and bond angles reported in the crystal structure of a closely related compound (Naicker *et al.*, 2012).

S2. Experimental

SOCl₂ (10 ml) was added into MeOH (60 ml) in a reaction flask at 273 K and *L*-3,4-dihydroxyphenylalanine (5.0 g, 25 mmol) was gradually added to this mixture. The temperature was increased to room temperature. After 24 h, the solvent was removed in *Vacuo*, to yield a white solid. The solid product was recrystallized from ethyl acetate by slow evaporation in the form of colorless single crystals of the title compound suitable for X-ray analysis.

S3. Refinement

An absolute structure was determined by the Flack (1983) method using 1117 Friedel pairs of reflections which were not merged. The H atoms bonded to N and O-atoms were located from a difference Fourier map and were allowed to refine freely. The H atoms bonded to C-atoms were positioned geometrically and refined using riding mode, with C—H = 0.95, 0.98, 0.99 and 1.00 Å, for aryl, methyl, methylene and methyne H-atoms, respectively; the $U_{iso}(H)$ were allowed at 1.5 $U_{eq}(C \text{ methyl})$ or 1.2 $U_{eq}(C \text{ non-methyl})$.

**Figure 1**

The molecular structure of the title compound with the atom numbering scheme. Displacement ellipsoids are drawn at the 30% probability level. H atoms are presented as small spheres of arbitrary radius.




Figure 2

A view of the unit cell showing partial packing of the title compound; hydrogen bonds are shown as dotted lines.

3-(3,4-Dihydroxyphenyl)-1-methoxy-1-oxopropan-2-aminium chloride

Crystal data

$C_{10}H_{14}NO_4^+ \cdot Cl^-$

$M_r = 247.67$

Orthorhombic, $P2_12_12_1$

Hall symbol: P 2ac 2ab

$a = 4.9969$ (15) Å

$b = 14.498$ (4) Å

$c = 16.109$ (5) Å

$V = 1167.1$ (6) Å³

$Z = 4$

$F(000) = 520$

$D_x = 1.410$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 3847 reflections

$\theta = 1.9$ – 27.9°

$\mu = 0.33$ mm⁻¹

$T = 113$ K

Prism, colourless

$0.20 \times 0.18 \times 0.12$ mm

Data collection

Rigaku Saturn724 CCD
diffractometer

Radiation source: rotating anode

Multilayer monochromator

Detector resolution: 14.22 pixels mm⁻¹

ω and φ scans

Absorption correction: multi-scan

(*CrystalClear*; Rigaku/MSK, 2005)

$T_{\min} = 0.938$, $T_{\max} = 0.962$

12293 measured reflections

2796 independent reflections

2054 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.051$
 $\theta_{\text{max}} = 27.9^\circ$, $\theta_{\text{min}} = 1.9^\circ$
 $h = -6 \rightarrow 6$

$k = -18 \rightarrow 19$
 $l = -21 \rightarrow 21$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.027$
 $wR(F^2) = 0.061$
 $S = 0.93$
 2796 reflections
 164 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods
 Secondary atom site location: difference Fourier
 map

Hydrogen site location: inferred from
 neighbouring sites
 H atoms treated by a mixture of independent
 and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0218P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.001$
 $\Delta\rho_{\text{max}} = 0.18 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.23 \text{ e } \text{\AA}^{-3}$
 Absolute structure: Flack (1983), 1117 Friedel
 pairs
 Absolute structure parameter: -0.03 (5)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	1.0416 (3)	0.23597 (8)	0.63923 (7)	0.0197 (3)
H1	1.079 (4)	0.2226 (12)	0.6879 (12)	0.029*
O2	0.8286 (3)	0.30070 (8)	0.78169 (8)	0.0217 (3)
H2	0.756 (4)	0.3261 (12)	0.8230 (13)	0.033*
O3	0.6921 (2)	0.64877 (8)	0.71119 (8)	0.0195 (3)
O4	0.3362 (2)	0.59134 (8)	0.77989 (8)	0.0178 (3)
N1	0.4423 (3)	0.63903 (10)	0.56253 (9)	0.0161 (3)
C1	0.5531 (3)	0.39895 (10)	0.55641 (11)	0.0181 (3)
H1D	0.4956	0.4210	0.5038	0.022*
C2	0.7537 (3)	0.33268 (10)	0.56033 (12)	0.0179 (4)
H2A	0.8285	0.3086	0.5106	0.022*
C3	0.8447 (3)	0.30165 (11)	0.63657 (11)	0.0147 (4)
C4	0.7284 (3)	0.33587 (10)	0.70898 (11)	0.0148 (4)
C5	0.5249 (3)	0.39978 (10)	0.70502 (11)	0.0162 (4)
H5	0.4443	0.4213	0.7548	0.019*
C6	0.4353 (3)	0.43341 (10)	0.62813 (10)	0.0146 (4)
C7	0.2087 (3)	0.50298 (11)	0.62467 (11)	0.0171 (4)
H7A	0.0765	0.4871	0.6681	0.020*
H7B	0.1181	0.4972	0.5703	0.020*
C8	0.2934 (3)	0.60417 (11)	0.63659 (11)	0.0140 (4)

H8	0.1278	0.6422	0.6433	0.017*
C9	0.4670 (3)	0.61855 (10)	0.71259 (10)	0.0143 (3)
C10	0.4845 (4)	0.59408 (12)	0.85779 (10)	0.0236 (4)
H10A	0.6548	0.5614	0.8511	0.035*
H10B	0.3790	0.5644	0.9015	0.035*
H10C	0.5195	0.6584	0.8731	0.035*
H1A	0.310 (4)	0.6414 (16)	0.5153 (14)	0.058 (7)*
H1B	0.598 (3)	0.6084 (11)	0.5497 (12)	0.021 (5)*
H1C	0.480 (4)	0.7033 (14)	0.5694 (12)	0.046 (6)*
Cl1	0.95062 (9)	0.63979 (3)	0.44397 (3)	0.01993 (10)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0229 (7)	0.0200 (7)	0.0162 (7)	0.0064 (6)	0.0009 (6)	0.0022 (5)
O2	0.0319 (8)	0.0208 (7)	0.0125 (7)	0.0098 (6)	0.0014 (6)	0.0004 (6)
O3	0.0163 (6)	0.0236 (6)	0.0187 (7)	-0.0039 (5)	-0.0014 (6)	-0.0012 (6)
O4	0.0203 (6)	0.0207 (6)	0.0123 (7)	-0.0020 (5)	0.0007 (5)	0.0007 (5)
N1	0.0176 (7)	0.0152 (7)	0.0157 (7)	-0.0011 (8)	-0.0012 (8)	0.0011 (8)
C1	0.0241 (8)	0.0171 (8)	0.0130 (8)	-0.0048 (8)	-0.0022 (9)	0.0019 (8)
C2	0.0224 (9)	0.0175 (8)	0.0139 (9)	-0.0022 (7)	0.0019 (9)	-0.0013 (8)
C3	0.0158 (9)	0.0114 (8)	0.0169 (10)	-0.0020 (7)	0.0008 (8)	0.0009 (8)
C4	0.0197 (9)	0.0122 (8)	0.0125 (9)	-0.0017 (7)	-0.0013 (8)	0.0014 (7)
C5	0.0221 (10)	0.0114 (8)	0.0152 (9)	-0.0004 (7)	0.0016 (8)	-0.0025 (7)
C6	0.0162 (8)	0.0096 (7)	0.0180 (9)	-0.0041 (7)	-0.0017 (8)	-0.0002 (7)
C7	0.0174 (9)	0.0148 (8)	0.0191 (10)	-0.0023 (7)	-0.0032 (8)	0.0004 (8)
C8	0.0131 (9)	0.0147 (8)	0.0141 (10)	-0.0004 (7)	0.0006 (7)	0.0015 (7)
C9	0.0179 (9)	0.0089 (8)	0.0161 (9)	0.0023 (7)	0.0008 (8)	-0.0006 (7)
C10	0.0315 (12)	0.0255 (10)	0.0138 (10)	-0.0007 (9)	-0.0035 (9)	0.0022 (8)
Cl1	0.0227 (2)	0.0207 (2)	0.0164 (2)	0.00256 (19)	-0.0024 (2)	0.0008 (2)

Geometric parameters (Å, °)

O1—C3	1.370 (2)	C2—H2A	0.9500
O1—H1	0.83 (2)	C3—C4	1.395 (2)
O2—C4	1.372 (2)	C4—C5	1.377 (2)
O2—H2	0.84 (2)	C5—C6	1.404 (2)
O3—C9	1.207 (2)	C5—H5	0.9500
O4—C9	1.326 (2)	C6—C7	1.517 (2)
O4—C10	1.458 (2)	C7—C8	1.539 (2)
N1—C8	1.494 (2)	C7—H7A	0.9900
N1—H1A	1.01 (2)	C7—H7B	0.9900
N1—H1B	0.92 (2)	C8—C9	1.515 (2)
N1—H1C	0.96 (2)	C8—H8	1.0000
C1—C6	1.390 (2)	C10—H10A	0.9800
C1—C2	1.390 (2)	C10—H10B	0.9800
C1—H1D	0.9500	C10—H10C	0.9800
C2—C3	1.385 (2)		

C3—O1—H1	110.8 (13)	C1—C6—C5	118.26 (16)
C4—O2—H2	110.8 (14)	C1—C6—C7	121.64 (15)
C9—O4—C10	116.42 (13)	C5—C6—C7	120.06 (15)
C8—N1—H1A	106.7 (12)	C6—C7—C8	115.09 (13)
C8—N1—H1B	116.0 (11)	C6—C7—H7A	108.5
H1A—N1—H1B	113.7 (16)	C8—C7—H7A	108.5
C8—N1—H1C	109.5 (12)	C6—C7—H7B	108.5
H1A—N1—H1C	100.5 (16)	C8—C7—H7B	108.5
H1B—N1—H1C	109.3 (16)	H7A—C7—H7B	107.5
C6—C1—C2	121.08 (17)	N1—C8—C9	108.29 (14)
C6—C1—H1D	119.5	N1—C8—C7	111.08 (14)
C2—C1—H1D	119.5	C9—C8—C7	112.92 (13)
C3—C2—C1	120.11 (17)	N1—C8—H8	108.1
C3—C2—H2A	119.9	C9—C8—H8	108.1
C1—C2—H2A	119.9	C7—C8—H8	108.1
O1—C3—C2	119.30 (16)	O3—C9—O4	125.64 (16)
O1—C3—C4	121.36 (15)	O3—C9—C8	124.63 (16)
C2—C3—C4	119.31 (16)	O4—C9—C8	109.72 (14)
O2—C4—C5	123.99 (16)	O4—C10—H10A	109.5
O2—C4—C3	115.46 (14)	O4—C10—H10B	109.5
C5—C4—C3	120.55 (16)	H10A—C10—H10B	109.5
C4—C5—C6	120.65 (16)	O4—C10—H10C	109.5
C4—C5—H5	119.7	H10A—C10—H10C	109.5
C6—C5—H5	119.7	H10B—C10—H10C	109.5
C6—C1—C2—C3	-1.7 (2)	C4—C5—C6—C7	179.30 (14)
C1—C2—C3—O1	179.68 (13)	C1—C6—C7—C8	-98.02 (19)
C1—C2—C3—C4	1.6 (2)	C5—C6—C7—C8	84.2 (2)
O1—C3—C4—O2	1.3 (2)	C6—C7—C8—N1	71.39 (19)
C2—C3—C4—O2	179.33 (14)	C6—C7—C8—C9	-50.5 (2)
O1—C3—C4—C5	-177.96 (14)	C10—O4—C9—O3	-3.4 (2)
C2—C3—C4—C5	0.1 (2)	C10—O4—C9—C8	175.40 (12)
O2—C4—C5—C6	179.21 (15)	N1—C8—C9—O3	-3.5 (2)
C3—C4—C5—C6	-1.6 (2)	C7—C8—C9—O3	119.97 (17)
C2—C1—C6—C5	0.2 (2)	N1—C8—C9—O4	177.71 (12)
C2—C1—C6—C7	-177.58 (14)	C7—C8—C9—O4	-58.84 (18)
C4—C5—C6—C1	1.5 (2)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N1—H1C...C11 ⁱ	0.96 (2)	2.29 (2)	3.209 (2)	161 (2)
N1—H1B...C11	0.92 (2)	2.49 (2)	3.178 (2)	132 (1)
N1—H1A...C11 ⁱⁱ	1.01 (2)	2.13 (2)	3.112 (2)	163 (2)
O2—H2...C11 ⁱⁱⁱ	0.84 (2)	2.26 (2)	3.086 (2)	167 (2)
O1—H1...O2	0.83 (2)	2.27 (2)	2.698 (2)	113 (2)
O1—H1...O3 ^{iv}	0.83 (2)	2.26 (2)	3.029 (2)	155 (2)

C8—H8···O3 ⁱⁱ	1.00	2.44	3.300 (2)	144
C10—H10C···O1 ^v	0.98	2.47	3.137 (3)	125

Symmetry codes: (i) $x-1/2, -y+3/2, -z+1$; (ii) $x-1, y, z$; (iii) $-x+3/2, -y+1, z+1/2$; (iv) $-x+2, y-1/2, -z+3/2$; (v) $-x+2, y+1/2, -z+3/2$.