

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

(E)-1-(5-Bromo-2-hydroxyphenyl)-3-[4-(dimethylamino)phenyl]prop-2-en-1-one

Guang-Bing Li, Lu Li and Guo-Xi Wang*

Department of Chemical & Environmental Engineering, Anyang Institute of Technology, Anyang 455000, People's Republic of China
Correspondence e-mail: aywgx@yahoo.com.cn


Received 14 April 2012; accepted 15 April 2012

Key indicators: single-crystal X-ray study; $T = 123$ K; mean $\sigma(\text{C}-\text{C}) = 0.009$ Å; R factor = 0.089; wR factor = 0.287; data-to-parameter ratio = 14.2.

In the title compound, $\text{C}_{17}\text{H}_{16}\text{BrNO}_2$, the two benzene rings make a dihedral angle of 7.4 (3) $^\circ$; the hydroxy group links to the carbonyl group *via* an intramolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bond. In the crystal, weak $\text{C}-\text{H}\cdots\text{O}$ interactions link the molecules into a supramolecular chain running along the c axis.

Related literature

For related compounds and structures, see: Dai & Chen (2011); Xu *et al.* (2011); Fu *et al.* (2011); Zheng *et al.* (2011).



Experimental

Crystal data

$\text{C}_{17}\text{H}_{16}\text{BrNO}_2$
 $M_r = 346.22$

Monoclinic, $P2_1/c$
 $a = 15.1905$ (5) Å

$b = 5.4501$ (2) Å
 $c = 19.7569$ (9) Å
 $\beta = 106.00$ (2) $^\circ$
 $V = 1572.34$ (11) Å 3
 $Z = 4$

Mo $K\alpha$ radiation
 $\mu = 2.62$ mm $^{-1}$
 $T = 123$ K
 $0.30 \times 0.05 \times 0.05$ mm

Data collection

Rigaku Mercury2 diffractometer
Absorption correction: multi-scan
(*CrystalClear*; Rigaku, 2005)
 $T_{\min} = 0.910$, $T_{\max} = 1.000$

8044 measured reflections
2736 independent reflections
1538 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.075$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.089$
 $wR(F^2) = 0.287$
 $S = 1.09$
2736 reflections
193 parameters

1 restraint
H-atom parameters constrained
 $\Delta\rho_{\max} = 1.58$ e Å $^{-3}$
 $\Delta\rho_{\min} = -1.11$ e Å $^{-3}$

Table 1
Hydrogen-bond geometry (Å, $^\circ$).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{O2}-\text{H2}\cdots\text{O1}$	0.82	1.82	2.543 (7)	146
$\text{C17}-\text{H17A}\cdots\text{O2}^i$	0.98	2.55	3.503 (10)	164

Symmetry code: (i) $x, -y + \frac{1}{2}, z - \frac{1}{2}$.

Data collection: *CrystalClear* (Rigaku, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

This work was supported by the start-up fund of Anyang Institute of Technology, China.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU5515).

References

- Dai, J. & Chen, X.-Y. (2011). *Acta Cryst.* **E67**, o287.
Fu, D.-W., Zhao, M.-M. & Ge, J.-Z. (2011). *J. Mol. Struct.* **1006**, 227–233.
Rigaku (2005). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Xu, R.-J., Fu, D.-W., Dai, J., Zhang, Y., Ge, J.-Z. & Ye, H.-Y. (2011). *Inorg. Chem. Commun.* **14**, 1093–1096.
Zheng, W.-N. (2011). *Acta Cryst.* **E67**, m344.

supporting information

Acta Cryst. (2012). E68, o1446 [doi:10.1107/S1600536812016327]

(E)-1-(5-Bromo-2-hydroxyphenyl)-3-[4-(dimethylamino)phenyl]prop-2-en-1-one

Guang-Bing Li, Lu Li and Guo-Xi Wang

S1. Comment

Organic amine derivatives are becoming increasingly important as new molecule-based crystalline materials with the potential optimal physical properties (Dai & Chen, 2011; Xu *et al.*, 2011). In addition, the amino compounds have found a wide range of applications in coordination chemistry as ligands to construct novel crystal structures (Fu *et al.*, 2011; Zheng, 2011). Herein, we report the crystal structure of the title compound, (E)-1-(5-bromo-2-hydroxyphenyl)-3-(4-(dimethylamino)phenyl)prop-2-en-1-one.


The asymmetric unit is composed of one whole amine molecule. The two benzene rings are nearly coplanar and twisted from each other by a dihedral of 7.4 (3)°. The hydroxy H2 atom was involved in intramolecular O—H···O hydrogen bonds interaction (Table 1 and Fig. 1).

S2. Experimental

The title compound (2.0 mmol) was solved in 50 mL methanol. Then the solution was evaporated slowly in the air. Red block crystals suitable for X-ray analysis were obtained after one week.

S3. Refinement

All H atoms attached to C atoms were fixed geometrically and treated as riding with C—H = 0.95 Å (aromatic) and C—H = 0.98 Å (methyl) with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{methyl})$. H atoms bonded to O atom was located in difference Fourier map and restrained with the H—O = 0.82 (2) Å. In the last stage of refinement they were treated as riding on the O atom with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$.

**Figure 1**

Molecular view of the title compound with the atomic numbering scheme. Displacement ellipsoids are drawn at the 30% probability level.

(E)-1-(5-Bromo-2-hydroxyphenyl)-3-[4-(dimethylamino)phenyl]prop-2-en-1-one

Crystal data

$C_{17}H_{16}BrNO_2$

$M_r = 346.22$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 15.1905 (5) \text{ \AA}$

$b = 5.4501 (2) \text{ \AA}$

$c = 19.7569 (9) \text{ \AA}$

$\beta = 106.00 (2)^\circ$

$V = 1572.34 (11) \text{ \AA}^3$

$Z = 4$

$F(000) = 704$

$D_x = 1.463 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 2736 reflections

$\theta = 2.2\text{--}27.5^\circ$

$\mu = 2.62 \text{ mm}^{-1}$

$T = 123 \text{ K}$

Needle, red

$0.30 \times 0.05 \times 0.05 \text{ mm}$

Data collection

Rigaku Mercury2
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

Detector resolution: $13.6612 \text{ pixels mm}^{-1}$

CCD profile fitting scans

Absorption correction: multi-scan

(*CrystalClear*; Rigaku, 2005)

$T_{\min} = 0.910$, $T_{\max} = 1.000$

8044 measured reflections

2736 independent reflections

1538 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.075$

$\theta_{\max} = 25.0^\circ$, $\theta_{\min} = 2.2^\circ$

$h = -18 \rightarrow 17$

$k = -6 \rightarrow 6$

$l = -23 \rightarrow 23$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.089$

$wR(F^2) = 0.287$

$S = 1.09$

2736 reflections

193 parameters

1 restraint

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.141P)^2 + 0.6743P]$

where $P = (F_o^2 + 2F_c^2)/3$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 1.58 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -1.11 \text{ e } \text{\AA}^{-3}$$

Extinction correction: *SHELXTL* (Sheldrick, 2008), $F_c^* = kFc[1 + 0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$
 Extinction coefficient: 0.002

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	1.08032 (7)	-0.2556 (2)	0.57680 (7)	0.1050 (7)
O1	0.6988 (3)	0.0521 (10)	0.6480 (2)	0.0469 (13)
C7	0.7599 (4)	0.0665 (12)	0.6153 (3)	0.0329 (15)
N1	0.5880 (3)	1.1541 (11)	0.3308 (3)	0.0355 (14)
O2	0.7796 (4)	-0.3021 (10)	0.7239 (3)	0.0600 (16)
H2	0.7357	-0.2242	0.7002	0.090*
C10	0.6627 (4)	0.6074 (12)	0.4911 (3)	0.0307 (14)
C9	0.6821 (4)	0.4141 (13)	0.5444 (3)	0.0331 (15)
H9A	0.6394	0.3991	0.5714	0.040*
C14	0.5541 (4)	0.9164 (12)	0.4255 (3)	0.0339 (15)
H14A	0.4969	0.9979	0.4183	0.041*
C13	0.6124 (4)	0.9790 (12)	0.3830 (3)	0.0331 (15)
C5	0.8429 (5)	-0.2911 (12)	0.6878 (4)	0.0386 (17)
C12	0.6988 (4)	0.8490 (12)	0.3966 (3)	0.0362 (16)
H12A	0.7404	0.8867	0.3699	0.043*
C1	0.9105 (4)	-0.1047 (14)	0.6015 (3)	0.0412 (17)
H1A	0.9106	0.0160	0.5669	0.049*
C6	0.8391 (4)	-0.1080 (12)	0.6344 (3)	0.0307 (14)
C2	0.9802 (5)	-0.2758 (13)	0.6193 (4)	0.0446 (19)
C15	0.5789 (5)	0.7398 (12)	0.4769 (4)	0.0342 (16)
H15A	0.5380	0.7044	0.5043	0.041*
C11	0.7207 (4)	0.6699 (12)	0.4483 (3)	0.0364 (16)
H11A	0.7772	0.5849	0.4555	0.044*
C4	0.9150 (5)	-0.4652 (14)	0.7037 (4)	0.0483 (19)
H4A	0.9168	-0.5864	0.7387	0.058*
C8	0.7530 (4)	0.2529 (11)	0.5605 (3)	0.0294 (15)
H8A	0.7987	0.2610	0.5360	0.035*
C16	0.5060 (5)	1.3053 (13)	0.3242 (4)	0.0429 (18)
H16A	0.4515	1.2004	0.3123	0.064*
H16B	0.5012	1.4277	0.2871	0.064*
H16C	0.5105	1.3887	0.3690	0.064*
C3	0.9833 (4)	-0.4609 (14)	0.6688 (4)	0.0438 (18)

H3A	1.0306	-0.5806	0.6784	0.053*
C17	0.6536 (6)	1.2333 (13)	0.2900 (4)	0.0441 (18)
H17A	0.6841	1.0888	0.2775	0.066*
H17B	0.6994	1.3444	0.3190	0.066*
H17C	0.6197	1.3183	0.2470	0.066*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.0628 (9)	0.1585 (14)	0.1133 (11)	0.0652 (8)	0.0570 (8)	0.0639 (8)
O1	0.038 (3)	0.055 (3)	0.057 (3)	0.006 (2)	0.029 (2)	0.010 (2)
C7	0.024 (3)	0.036 (4)	0.041 (4)	-0.005 (3)	0.012 (3)	-0.004 (3)
N1	0.029 (3)	0.041 (3)	0.039 (3)	0.008 (2)	0.014 (3)	-0.003 (2)
O2	0.049 (4)	0.063 (4)	0.069 (4)	0.005 (3)	0.020 (3)	0.029 (3)
C10	0.026 (3)	0.032 (3)	0.034 (3)	0.002 (3)	0.008 (3)	-0.004 (3)
C9	0.024 (3)	0.043 (4)	0.034 (3)	0.000 (3)	0.011 (3)	-0.006 (3)
C14	0.026 (3)	0.033 (3)	0.047 (4)	0.006 (3)	0.017 (3)	-0.009 (3)
C13	0.030 (3)	0.036 (4)	0.033 (3)	-0.002 (3)	0.008 (3)	-0.008 (3)
C5	0.034 (4)	0.040 (4)	0.045 (4)	-0.007 (3)	0.016 (3)	0.001 (3)
C12	0.027 (4)	0.040 (4)	0.045 (4)	0.000 (3)	0.016 (3)	0.002 (3)
C1	0.038 (4)	0.050 (4)	0.039 (4)	0.010 (3)	0.016 (3)	0.012 (3)
C6	0.026 (3)	0.036 (4)	0.027 (3)	-0.003 (3)	0.003 (3)	0.001 (3)
C2	0.032 (4)	0.053 (5)	0.052 (5)	0.011 (3)	0.015 (4)	0.005 (3)
C15	0.028 (4)	0.041 (4)	0.037 (4)	-0.003 (3)	0.015 (3)	-0.006 (3)
C11	0.024 (3)	0.046 (4)	0.043 (4)	0.006 (3)	0.016 (3)	-0.002 (3)
C4	0.044 (4)	0.044 (4)	0.046 (4)	-0.004 (3)	-0.005 (4)	0.004 (3)
C8	0.021 (3)	0.035 (3)	0.033 (4)	-0.001 (3)	0.009 (3)	-0.003 (3)
C16	0.046 (5)	0.044 (4)	0.038 (4)	0.016 (3)	0.011 (3)	0.006 (3)
C3	0.030 (4)	0.043 (4)	0.049 (4)	0.008 (3)	-0.003 (3)	-0.001 (3)
C17	0.054 (5)	0.038 (4)	0.046 (4)	0.000 (3)	0.025 (4)	0.005 (3)

Geometric parameters (Å, °)

Br1—C2	1.933 (8)	C5—C6	1.441 (9)
O1—C7	1.271 (7)	C12—C11	1.386 (9)
C7—C8	1.468 (9)	C12—H12A	0.9500
C7—C6	1.499 (9)	C1—C2	1.382 (9)
N1—C13	1.380 (8)	C1—C6	1.409 (9)
N1—C16	1.469 (8)	C1—H1A	0.9500
N1—C17	1.506 (9)	C2—C3	1.396 (10)
O2—C5	1.347 (9)	C15—H15A	0.9500
O2—H2	0.8200	C11—H11A	0.9500
C10—C11	1.421 (9)	C4—C3	1.397 (9)
C10—C15	1.423 (9)	C4—H4A	0.9500
C10—C9	1.461 (9)	C8—H8A	0.9500
C9—C8	1.358 (9)	C16—H16A	0.9800
C9—H9A	0.9500	C16—H16B	0.9800
C14—C15	1.374 (9)	C16—H16C	0.9800

C14—C13	1.419 (8)	C3—H3A	0.9500
C14—H14A	0.9500	C17—H17A	0.9800
C13—C12	1.450 (9)	C17—H17B	0.9800
C5—C4	1.417 (10)	C17—H17C	0.9800
O1—C7—C8	120.3 (6)	C1—C2—C3	122.8 (7)
O1—C7—C6	118.8 (6)	C1—C2—Br1	119.4 (6)
C8—C7—C6	120.9 (5)	C3—C2—Br1	117.8 (5)
C13—N1—C16	120.0 (5)	C14—C15—C10	123.3 (6)
C13—N1—C17	121.2 (5)	C14—C15—H15A	118.3
C16—N1—C17	117.5 (6)	C10—C15—H15A	118.3
C5—O2—H2	105.1	C12—C11—C10	123.1 (6)
C11—C10—C15	115.4 (6)	C12—C11—H11A	118.5
C11—C10—C9	124.6 (6)	C10—C11—H11A	118.5
C15—C10—C9	120.0 (6)	C3—C4—C5	120.9 (7)
C8—C9—C10	128.6 (6)	C3—C4—H4A	119.6
C8—C9—H9A	115.7	C5—C4—H4A	119.6
C10—C9—H9A	115.7	C9—C8—C7	120.7 (6)
C15—C14—C13	121.1 (6)	C9—C8—H8A	119.7
C15—C14—H14A	119.5	C7—C8—H8A	119.7
C13—C14—H14A	119.5	N1—C16—H16A	109.5
N1—C13—C14	121.7 (6)	N1—C16—H16B	109.5
N1—C13—C12	121.2 (5)	H16A—C16—H16B	109.5
C14—C13—C12	117.0 (6)	N1—C16—H16C	109.5
O2—C5—C4	118.2 (6)	H16A—C16—H16C	109.5
O2—C5—C6	121.8 (6)	H16B—C16—H16C	109.5
C4—C5—C6	119.9 (7)	C2—C3—C4	118.1 (6)
C11—C12—C13	120.1 (6)	C2—C3—H3A	120.9
C11—C12—H12A	119.9	C4—C3—H3A	120.9
C13—C12—H12A	119.9	N1—C17—H17A	109.5
C2—C1—C6	120.5 (6)	N1—C17—H17B	109.5
C2—C1—H1A	119.8	H17A—C17—H17B	109.5
C6—C1—H1A	119.8	N1—C17—H17C	109.5
C1—C6—C5	117.7 (6)	H17A—C17—H17C	109.5
C1—C6—C7	122.8 (6)	H17B—C17—H17C	109.5
C5—C6—C7	119.4 (6)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O2—H2 \cdots O1	0.82	1.82	2.543 (7)	146
C17—H17A \cdots O2 ⁱ	0.98	2.55	3.503 (10)	164

Symmetry code: (i) *x*, $-y+1/2$, $z-1/2$.