

2-Aminoethanaminium iodide

 Alan R. Kennedy^a and Maurice O. Okoth^{b*}

^aDepartment of Pure & Applied Chemistry, WestCHEM, University of Strathclyde, 295 Cathedral Street, Glasgow G1 1XL, Scotland, and ^bDepartment of Chemistry and Biochemistry, Chepkoilel University College, PO Box 1125-30100, Eldoret, Kenya
Correspondence e-mail: okothmdo@mu.ac.ke


Received 24 April 2012; accepted 7 May 2012

Key indicators: single-crystal X-ray study; $T = 123$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.016; wR factor = 0.035; data-to-parameter ratio = 20.2.

The title salt, $[\text{NH}_3\text{CH}_2\text{CH}_2\text{NH}_2]^+\text{I}^-$, has an array structure based on strong intermolecular $\text{N}-\text{H}\cdots\text{N}$ hydrogen bonding formed between the ammonium and amine groups of adjacent cations. This interaction gives a helical chain of cations that runs parallel to the b axis. The four remaining NH group H atoms all form hydrogen bonds to the iodide anion, and these iodide anions lie in channels parallel to the cation-cation chains.

Related literature

For syntheses and structures of salt forms of the related ethylene-1,2-diammonium, see: Chen (2009); Saidi *et al.* (2011). For a structural example of a complex of ethylene-1,2-diammonium, see: Zhang *et al.* (2006). For the synthesis that gave the title compound as a by-product, see: Kennedy *et al.* (2011). For C–N bond length changes in another mono-protonated symmetrical diamine, see: Craig *et al.* (2012).



Experimental

Crystal data

$\text{C}_2\text{H}_9\text{N}_2^+\text{I}^-$	$V = 1178.29$ (6) Å ³
$M_r = 188.01$	$Z = 8$
Orthorhombic, $Pbca$	Mo $K\alpha$ radiation
$a = 8.1380$ (2) Å	$\mu = 5.29$ mm ⁻¹
$b = 8.6259$ (2) Å	$T = 123$ K
$c = 16.7854$ (6) Å	$0.28 \times 0.08 \times 0.04$ mm

Data collection

Oxford Diffraction Gemini S diffractometer	13735 measured reflections
Absorption correction: multi-scan (<i>CrysAlis PRO</i> ; Oxford Diffraction, 2010)	1675 independent reflections
$T_{\min} = 0.405$, $T_{\max} = 0.805$	1405 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.025$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.016$	83 parameters
$wR(F^2) = 0.035$	All H-atom parameters refined
$S = 1.08$	$\Delta\rho_{\text{max}} = 0.58$ e Å ⁻³
1675 reflections	$\Delta\rho_{\text{min}} = -0.39$ e Å ⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1N}\cdots\text{N2}^{\text{i}}$	1.06 (2)	1.75 (2)	2.805 (2)	173.0 (19)
$\text{N1}-\text{H2N}\cdots\text{I1}$	0.882 (19)	3.231 (18)	3.6502 (14)	111.6 (13)
$\text{N1}-\text{H2N}\cdots\text{I1}^{\text{ii}}$	0.882 (19)	2.820 (19)	3.6047 (14)	148.9 (15)
$\text{N1}-\text{H3N}\cdots\text{I1}^{\text{iii}}$	0.84 (2)	2.78 (2)	3.5713 (16)	157.9 (16)
$\text{N2}-\text{H4N}\cdots\text{I1}^{\text{iv}}$	0.84 (2)	2.96 (2)	3.7346 (16)	155.5 (16)
$\text{N2}-\text{H5N}\cdots\text{I1}^{\text{v}}$	0.866 (19)	3.152 (19)	3.9328 (15)	151.2 (14)

Symmetry codes: (i) $-x + \frac{3}{2}, y + \frac{1}{2}, z$; (ii) $-x + \frac{1}{2}, y - \frac{1}{2}, z$; (iii) $x + \frac{1}{2}, y, -z + \frac{3}{2}$; (iv) $-x + 1, -y, -z + 1$; (v) $x + 1, y, z$.

Data collection: *CrysAlis PRO* (Oxford Diffraction, 2010); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SIR92* (Burla *et al.*, 2005); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *X-SEED* (Barbour, 2001); software used to prepare material for publication: *SHELXL97*.

MOO thanks the Commonwealth Scholarship Commission and the British Council for funding and Moi University for sabbatical leave.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BR2201).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
 Burla, M. C., Caliandro, R., Camalli, M., Carrozzini, B., Cascarano, G. L., De Caro, L., Giacovazzo, C., Polidori, G. & Spagna, R. (2005). *J. Appl. Cryst.* **38**, 381–388.
 Chen, L.-Z. (2009). *Acta Cryst.* **E65**, o2625.
 Craig, G. E., Johnson, C. & Kennedy, A. R. (2012). *Acta Cryst.* **E68**, o787.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Kennedy, A. R., Lutta, S. T., Morrison, C. A., Okoth, M. O. & Orang'o, D. M. (2011). *Acta Cryst.* **E67**, o682–o683.
 Oxford Diffraction (2010). *CrysAlis PRO*. Oxford Diffraction Ltd, Yarnton, England.
 Saidi, K., Kamoun, S. & Ayedi, H. F. (2011). *J. Chem. Crystallogr.* **41**, 1258–1261.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Zhang, Z.-J., Guo, G.-C., Xu, G., Fu, M.-L., Zou, J.-P. & Huang, J.-S. (2006). *Inorg. Chem.* **45**, 10028–10030.

supporting information

Acta Cryst. (2012). E68, o1731 [doi:10.1107/S160053681202065X]

2-Aminoethanaminium iodide

Alan R. Kennedy and Maurice O. Okoth

S1. Comment

Despite the common use of ethylene-1,2-diamine as a ligand, there are surprisingly few metal containing crystal structures that feature its cationic form ethylene-1,2-diammonium (for an example see Zhang *et al.*, 2006) and only two structures of simple salt forms of ethylene-1,2-diammonium (Chen, 2009; Saidi *et al.*, 2011). There appears to be no previous reports of structures that contain the singly protonated cation, $\text{NH}_3\text{CH}_2\text{CH}_2\text{NH}_2$.

Crystals of ethylene-2-amine-1-ammonium iodide (I) were recovered whilst trying to replicate the synthesis of the macrocyclic species 5,7,7,12,14,14-hexamethyl-4,8-diaza-1,11-diazoniocyclotetradeca-4,11-diene diiodide, the first step of which is addition of HI to ethylene-1,2-diamine in ethanol (Kennedy *et al.*, 2011). Investigation of the structure clearly showed that the base is protonated at only one site, see Figure 1. This is confirmed by location and independent refinement of the hydrogen atoms and by the slight lengthening of the C1—N1 bond as compared to the C2—N2 bond (compare 1.484 (2) and 1.467 (2) Å). Similar differences are seen in other symmetrical diamines that have been monoprotinated (see for example Craig *et al.*, 2012).


Atom H1N is a hydrogen bond donor that interacts with N2 to form the relatively short cation to cation hydrogen bond that gives the one dimensional helical chain running in the *b* direction, as shown in Figure 2. The four other N—H hydrogen atoms all interact with the iodide anion ($\text{N}\cdots\text{I}$ range 3.5713 (16) to 3.9328 (15) Å), see Table 1 for details. These interactions combine to give the packing motif shown in Figure 3, with channels of anions parallel to the *b* direction and thus also parallel to the cation-cation hydrogen bonded chains.

S2. Experimental


Crystals of (I) were obtained from ethanol solution.

S3. Refinement


All the H-atoms were found through difference synthesis and refined isotropically. The N1 to H1N distance is 1.06 (2) Å. This forms part of the N—H \cdots N hydrogen bond and may reflect some small degree of positional disorder over two sites.

**Figure 1**

The molecular structure of (I). Displacement ellipsoids are drawn at the 50% probability level with H-atoms drawn as spheres of arbitrary size.

**Figure 2**

Hydrogen bonding forms helical, one-dimensional chains of cations that propagate in the crystallographic *b* direction.

**Figure 3**

Packing in (I) viewed along the *b* direction.

2-Aminoethaniminium iodide

Crystal data

$C_2H_9N_2^+I^-$

$M_r = 188.01$

Orthorhombic, *Pbca*

Hall symbol: -P 2ac 2ab

$a = 8.1380$ (2) Å

$b = 8.6259$ (2) Å

$c = 16.7854$ (6) Å

$V = 1178.29$ (6) Å³

$Z = 8$

$F(000) = 704$

$D_x = 2.120$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 7738 reflections

$\theta = 3.4\text{--}30.4^\circ$

$\mu = 5.29$ mm⁻¹

$T = 123$ K

Cut needle, colourless

$0.28 \times 0.08 \times 0.04$ mm

Data collection

Oxford Diffraction Gemini S
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

ω scans

Absorption correction: multi-scan

(*CrysAlis PRO*; Oxford Diffraction, 2010)

$T_{\min} = 0.405$, $T_{\max} = 0.805$

13735 measured reflections

1675 independent reflections

1405 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.025$
 $\theta_{\text{max}} = 30.5^\circ$, $\theta_{\text{min}} = 3.5^\circ$

$h = -11 \rightarrow 11$
 $k = -12 \rightarrow 12$
 $l = -23 \rightarrow 22$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.016$
 $wR(F^2) = 0.035$
 $S = 1.08$
 1675 reflections
 83 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: difference Fourier map
 All H-atom parameters refined
 $w = 1/[\sigma^2(F_o^2) + (0.0188P)^2 + 0.0175P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.003$
 $\Delta\rho_{\text{max}} = 0.58 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.39 \text{ e } \text{\AA}^{-3}$
 Extinction correction: *SHELXL97* (Sheldrick,
 2008), $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
 Extinction coefficient: 0.00618 (17)

Special details

Experimental. Absorption correction: CrysAlis PRO (Oxford Diffraction, 2010). Empirical absorption correction using spherical harmonics, implemented in SCALE3 ABSPACK scaling algorithm.

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
I1	0.153737 (12)	0.243717 (9)	0.639779 (7)	0.01478 (5)
N1	0.57239 (17)	0.09977 (16)	0.66467 (10)	0.0152 (3)
N2	0.91366 (18)	-0.12957 (16)	0.57234 (10)	0.0166 (3)
C1	0.72638 (19)	0.00806 (18)	0.65975 (11)	0.0156 (3)
C2	0.7645 (2)	-0.03399 (18)	0.57391 (10)	0.0164 (3)
H1	0.720 (2)	-0.083 (2)	0.6923 (11)	0.022 (5)*
H2	0.812 (2)	0.073 (2)	0.6768 (12)	0.025 (5)*
H3	0.677 (2)	-0.0965 (19)	0.5534 (11)	0.020 (5)*
H4	0.771 (2)	0.0590 (19)	0.5396 (11)	0.018 (4)*
H1N	0.573 (3)	0.206 (3)	0.6334 (13)	0.030 (5)*
H2N	0.487 (2)	0.042 (2)	0.6523 (12)	0.027 (5)*
H3N	0.561 (2)	0.131 (2)	0.7119 (12)	0.028 (6)*
H4N	0.932 (2)	-0.161 (2)	0.5259 (13)	0.029 (5)*
H5N	0.998 (2)	-0.073 (2)	0.5848 (13)	0.029 (6)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
I1	0.01452 (7)	0.01487 (7)	0.01494 (8)	0.00109 (3)	-0.00020 (4)	-0.00022 (4)

N1	0.0146 (7)	0.0161 (7)	0.0150 (8)	-0.0014 (5)	0.0018 (6)	-0.0016 (5)
N2	0.0151 (7)	0.0194 (7)	0.0152 (8)	0.0000 (5)	0.0019 (6)	-0.0009 (6)
C1	0.0140 (8)	0.0188 (8)	0.0141 (8)	-0.0001 (6)	-0.0006 (6)	0.0007 (6)
C2	0.0164 (8)	0.0191 (7)	0.0136 (8)	0.0010 (6)	0.0007 (7)	0.0010 (6)

Geometric parameters (Å, °)

N1—C1	1.484 (2)	N2—H5N	0.866 (19)
N1—H1N	1.06 (2)	C1—C2	1.518 (2)
N1—H2N	0.882 (19)	C1—H1	0.959 (18)
N1—H3N	0.84 (2)	C1—H2	0.937 (19)
N2—C2	1.467 (2)	C2—H3	0.959 (17)
N2—H4N	0.84 (2)	C2—H4	0.989 (18)
C1—N1—H1N	115.5 (12)	C2—C1—H1	110.8 (11)
C1—N1—H2N	110.4 (12)	N1—C1—H2	107.0 (11)
H1N—N1—H2N	112.4 (17)	C2—C1—H2	106.3 (13)
C1—N1—H3N	108.5 (13)	H1—C1—H2	110.7 (18)
H1N—N1—H3N	100.7 (16)	N2—C2—C1	108.70 (14)
H2N—N1—H3N	108.6 (17)	N2—C2—H3	107.1 (10)
C2—N2—H4N	110.2 (13)	C1—C2—H3	108.8 (11)
C2—N2—H5N	109.6 (12)	N2—C2—H4	113.8 (10)
H4N—N2—H5N	105.2 (18)	C1—C2—H4	111.7 (11)
N1—C1—C2	110.67 (14)	H3—C2—H4	106.6 (15)
N1—C1—H1	111.2 (11)		
N1—C1—C2—N2	-177.75 (12)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N1—H1N...N2 ⁱ	1.06 (2)	1.75 (2)	2.805 (2)	173.0 (19)
N1—H2N...I1	0.882 (19)	3.231 (18)	3.6502 (14)	111.6 (13)
N1—H2N...I1 ⁱⁱ	0.882 (19)	2.820 (19)	3.6047 (14)	148.9 (15)
N1—H3N...I1 ⁱⁱⁱ	0.84 (2)	2.78 (2)	3.5713 (16)	157.9 (16)
N2—H4N...I1 ^{iv}	0.84 (2)	2.96 (2)	3.7346 (16)	155.5 (16)
N2—H5N...I1 ^v	0.866 (19)	3.152 (19)	3.9328 (15)	151.2 (14)

Symmetry codes: (i) $-x+3/2, y+1/2, z$; (ii) $-x+1/2, y-1/2, z$; (iii) $x+1/2, y, -z+3/2$; (iv) $-x+1, -y, -z+1$; (v) $x+1, y, z$.