

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

# (E)-4-Methyl-N'-(3-nitrobenzylidene)-benzohydrazide

Hua-Nan Hu and Shi-Yong Liu\*

College of Chemistry &amp; Pharmacy, Taizhou University, Taizhou Zhejiang 317000, People's Republic of China

Correspondence e-mail: liushiyong2012@yahoo.cn


Received 28 April 2012; accepted 29 April 2012

 Key indicators: single-crystal X-ray study;  $T = 298$  K; mean  $\sigma(\text{C}-\text{C}) = 0.004$  Å;  $R$  factor = 0.049;  $wR$  factor = 0.103; data-to-parameter ratio = 13.0.

In the title compound,  $\text{C}_{15}\text{H}_{13}\text{N}_3\text{O}_3$ , the dihedral angle between the benzene rings is  $1.01(3)^\circ$  and that between the nitro group and its attached ring is  $5.99(15)^\circ$ . In the crystal, molecules are linked by  $\text{N}-\text{H}\cdots\text{O}$  hydrogen bonds generating  $C(4)$  chains along  $[010]$ .

## Related literature

For related structures, see: Liu & You (2010); Liu & Wang (2010); Xu *et al.* (2009); Shafiq *et al.* (2009).



## Experimental

## Crystal data

 $\text{C}_{15}\text{H}_{13}\text{N}_3\text{O}_3$   
 $M_r = 283.28$   
 Orthorhombic,  $Pca2_1$ 
 $a = 32.657(3)$  Å  
 $b = 4.7861(15)$  Å  
 $c = 8.7596(12)$  Å

 $V = 1369.1(5)$  Å<sup>3</sup>  
 $Z = 4$   
 Mo  $K\alpha$  radiation

 $\mu = 0.10$  mm<sup>-1</sup>  
 $T = 298$  K  
 $0.20 \times 0.20 \times 0.17$  mm

## Data collection

 Bruker SMART CCD diffractometer  
 Absorption correction: multi-scan (*SADABS*; Bruker, 2001)  
 $T_{\min} = 0.981$ ,  $T_{\max} = 0.984$ 

 9086 measured reflections  
 2539 independent reflections  
 2036 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.045$ 

## Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.049$   
 $wR(F^2) = 0.103$   
 $S = 1.04$   
 2539 reflections  
 195 parameters  
 2 restraints

 H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\max} = 0.13$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.17$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N3}-\text{H3}\cdots\text{O3}^i$	0.89 (1)	2.13 (2)	2.909 (3)	146 (2)

 Symmetry code: (i)  $x, y - 1, z$ .

Data collection: *SMART* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

The authors acknowledge the Zhejiang Provincial Natural Science Foundation of China (project No. Y12B020017).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB6769).

## References

- Bruker (2001). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Bruker (2007). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Liu, S.-Y. & Wang, X. (2010). *Acta Cryst.* **E66**, o1775.  
 Liu, S.-Y. & You, Z. (2010). *Acta Cryst.* **E66**, o1652.  
 Shafiq, Z., Yaqub, M., Tahir, M. N., Hussain, A. & Iqbal, M. S. (2009). *Acta Cryst.* **E65**, o2898.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
 Xu, L., Huang, S.-S., Zhang, B.-J., Wang, S.-Y. & Zhang, H.-L. (2009). *Acta Cryst.* **E65**, o2412.

## supporting information

*Acta Cryst.* (2012). E68, o1613 [doi:10.1107/S1600536812019198]

**(E)-4-Methyl-N'-(3-nitrobenzylidene)benzohydrazide**

Hua-Nan Hu and Shi-Yong Liu

**S1. Comment**

As a continuation of our work on similar compounds (Liu & You, 2010; Liu & Wang, 2010), we report herein the crystal structure of the title compound a new hydrazone.

The molecular structure of the title compound is shown in Fig. 1. The two benzene ring system are inclined at a dihedral angle of 0.6 (3)°. All the bond lengths are comparable to those observed in related structures (Xu *et al.*, 2009; Shafiq *et al.*, 2009) and those we reported previously.


In the crystal structure, molecules are linked through N–H···O hydrogen bonds, to form one-dimensional chains running along the *b* axis (Fig. 2 and Table 1).

**S2. Experimental**

The title compound was prepared by the condensation reaction of 3-nitrobenzaldehyde (0.05 mol, 7.6 g) and 4-methylbenzohydrazide (0.05 mol, 7.5 g) in anhydrous methanol (100 ml) at ambient temperature. Yellow block-shaped single crystals were obtained by slow evaporation of the solution for several days.


**S3. Refinement**

Anomalous dispersion was negligible and the absolute structure could not be determined in the present experiment. H3 was located from a difference Fourier map and refined isotropically, with the N–H distance restrained to 0.90 (1) Å. The remaining H atoms were positioned geometrically and constrained to ride on their parent atoms, with C–H distances of 0.93–0.96 Å, and with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$  and  $1.5U_{\text{eq}}(\text{C15})$ .



**Figure 1**

The molecular structure of the title compound. Displacement ellipsoids are drawn at the 30% probability level.



**Figure 2**

The molecular packing of the title compound, viewed along the *c* axis. Hydrogen bonds are shown as dashed lines. Hydrogen atoms not involved in hydrogen bonding have been omitted.

**(*E*)-4-Methyl-*N'*-(3-nitrobenzylidene)benzohydrazide**

*Crystal data*

$C_{15}H_{13}N_3O_3$

$M_r = 283.28$

Orthorhombic,  $Pca2_1$

$a = 32.657$  (3) Å

$b = 4.7861$  (15) Å

$c = 8.7596$  (12) Å

$V = 1369.1$  (5) Å<sup>3</sup>

$Z = 4$

$F(000) = 592$

$D_x = 1.374$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 1981 reflections

$\theta = 2.5$ – $24.3^\circ$

$\mu = 0.10$  mm<sup>-1</sup>

$T = 298$  K

Block, yellow

$0.20 \times 0.20 \times 0.17$  mm

*Data collection*

Bruker SMART CCD  
diffractometer  
Radiation source: fine-focus sealed tube  
Graphite monochromator  
 $\omega$  scans  
Absorption correction: multi-scan  
(*SADABS*; Bruker, 2001)  
 $T_{\min} = 0.981$ ,  $T_{\max} = 0.984$

9086 measured reflections  
2539 independent reflections  
2036 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.045$   
 $\theta_{\text{max}} = 25.5^\circ$ ,  $\theta_{\text{min}} = 2.5^\circ$   
 $h = -39 \rightarrow 37$   
 $k = -5 \rightarrow 5$   
 $l = -10 \rightarrow 10$

*Refinement*

Refinement on  $F^2$   
Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.049$   
 $wR(F^2) = 0.103$   
 $S = 1.04$   
2539 reflections  
195 parameters  
2 restraints  
Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map  
Hydrogen site location: inferred from  
neighbouring sites  
H atoms treated by a mixture of independent  
and constrained refinement  
 $w = 1/[\sigma^2(F_o^2) + (0.0473P)^2 + 0.1113P]$   
where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\text{max}} < 0.001$   
 $\Delta\rho_{\text{max}} = 0.13 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -0.17 \text{ e } \text{\AA}^{-3}$

*Special details*

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted R-factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional R-factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\sigma(F^2)$  is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and R-factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.85121 (8)	0.5734 (5)	0.4452 (2)	0.0467 (6)
N2	0.70734 (7)	0.2646 (4)	0.2869 (3)	0.0444 (5)
N3	0.66666 (7)	0.2015 (4)	0.2812 (3)	0.0455 (5)
O1	0.88853 (7)	0.5929 (5)	0.4514 (3)	0.0749 (7)
O2	0.82822 (7)	0.7214 (4)	0.5169 (2)	0.0597 (6)
O3	0.64841 (6)	0.6273 (4)	0.3664 (2)	0.0553 (5)
C1	0.77578 (8)	0.1201 (5)	0.2510 (3)	0.0413 (6)
C2	0.79253 (8)	0.3213 (5)	0.3444 (3)	0.0377 (6)
H2	0.7758	0.4289	0.4072	0.045*
C3	0.83377 (8)	0.3605 (5)	0.3438 (3)	0.0382 (6)
C4	0.85976 (8)	0.2118 (6)	0.2508 (3)	0.0488 (7)
H4	0.8878	0.2448	0.2521	0.059*
C5	0.84325 (9)	0.0139 (6)	0.1563 (3)	0.0527 (7)
H5	0.8601	-0.0881	0.0914	0.063*
C6	0.80177 (8)	-0.0341 (5)	0.1572 (3)	0.0457 (7)

H6	0.7909	-0.1717	0.0942	0.055*
C7	0.73153 (8)	0.0702 (6)	0.2463 (3)	0.0479 (7)
H7	0.7213	-0.1012	0.2140	0.057*
C8	0.63869 (8)	0.3942 (5)	0.3216 (3)	0.0418 (6)
C9	0.59528 (8)	0.3070 (5)	0.3091 (3)	0.0405 (6)
C10	0.56597 (9)	0.4440 (6)	0.3943 (4)	0.0591 (8)
H10	0.5739	0.5841	0.4617	0.071*
C11	0.52522 (10)	0.3763 (7)	0.3809 (4)	0.0677 (9)
H11	0.5062	0.4680	0.4418	0.081*
C12	0.51204 (9)	0.1772 (7)	0.2801 (4)	0.0600 (8)
C13	0.54139 (9)	0.0401 (7)	0.1948 (3)	0.0622 (9)
H13	0.5333	-0.0988	0.1269	0.075*
C14	0.58242 (9)	0.1047 (6)	0.2083 (3)	0.0533 (8)
H14	0.6015	0.0107	0.1488	0.064*
C15	0.46734 (9)	0.0998 (9)	0.2613 (5)	0.0889 (12)
H15A	0.4505	0.2489	0.2990	0.133*
H15B	0.4617	-0.0678	0.3177	0.133*
H15C	0.4615	0.0692	0.1551	0.133*
H3	0.6588 (7)	0.025 (3)	0.266 (3)	0.050 (8)*

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
N1	0.0515 (16)	0.0457 (13)	0.0428 (13)	-0.0090 (12)	-0.0092 (12)	0.0008 (11)
N2	0.0438 (14)	0.0381 (12)	0.0512 (12)	-0.0091 (10)	-0.0044 (11)	-0.0003 (11)
N3	0.0398 (13)	0.0312 (12)	0.0655 (15)	-0.0094 (10)	0.0000 (11)	-0.0075 (12)
O1	0.0505 (14)	0.0829 (16)	0.0915 (17)	-0.0158 (12)	-0.0164 (12)	-0.0160 (13)
O2	0.0715 (14)	0.0567 (13)	0.0507 (12)	-0.0013 (12)	0.0013 (10)	-0.0194 (11)
O3	0.0679 (13)	0.0328 (10)	0.0651 (12)	-0.0097 (9)	0.0031 (11)	-0.0116 (9)
C1	0.0447 (16)	0.0338 (14)	0.0456 (15)	0.0006 (12)	-0.0042 (13)	0.0025 (12)
C2	0.0428 (16)	0.0323 (12)	0.0379 (14)	0.0017 (11)	0.0006 (11)	-0.0043 (11)
C3	0.0446 (15)	0.0355 (13)	0.0343 (14)	-0.0024 (12)	-0.0025 (12)	-0.0003 (11)
C4	0.0428 (16)	0.0509 (17)	0.0526 (17)	0.0012 (13)	-0.0004 (14)	0.0031 (15)
C5	0.0582 (19)	0.0543 (18)	0.0456 (15)	0.0114 (14)	0.0079 (15)	-0.0043 (15)
C6	0.0583 (18)	0.0373 (15)	0.0415 (14)	0.0008 (13)	-0.0084 (14)	-0.0067 (13)
C7	0.0512 (18)	0.0368 (15)	0.0555 (17)	-0.0060 (13)	-0.0034 (14)	-0.0054 (13)
C8	0.0505 (16)	0.0337 (14)	0.0412 (15)	-0.0047 (12)	0.0005 (12)	-0.0008 (12)
C9	0.0473 (16)	0.0310 (13)	0.0433 (15)	0.0013 (11)	0.0024 (12)	0.0023 (11)
C10	0.062 (2)	0.0535 (17)	0.0621 (19)	-0.0044 (15)	0.0135 (16)	-0.0127 (16)
C11	0.059 (2)	0.065 (2)	0.078 (2)	0.0081 (17)	0.0222 (18)	-0.007 (2)
C12	0.0511 (19)	0.068 (2)	0.0614 (19)	0.0021 (15)	0.0041 (16)	0.0096 (18)
C13	0.055 (2)	0.071 (2)	0.0607 (19)	-0.0138 (16)	-0.0062 (15)	-0.0175 (16)
C14	0.0497 (19)	0.0525 (18)	0.0577 (18)	-0.0012 (14)	0.0035 (13)	-0.0163 (14)
C15	0.048 (2)	0.127 (3)	0.091 (3)	-0.002 (2)	0.000 (2)	0.008 (3)

*Geometric parameters (Å, °)*

N1—O2	1.208 (3)	C6—H6	0.9300
N1—O1	1.224 (3)	C7—H7	0.9300
N1—C3	1.467 (3)	C8—C9	1.482 (3)
N2—C7	1.272 (3)	C9—C14	1.376 (3)
N2—N3	1.364 (3)	C9—C10	1.379 (4)
N3—C8	1.345 (3)	C10—C11	1.375 (4)
N3—H3	0.891 (10)	C10—H10	0.9300
O3—C8	1.225 (3)	C11—C12	1.369 (5)
C1—C2	1.377 (3)	C11—H11	0.9300
C1—C6	1.393 (4)	C12—C13	1.381 (4)
C1—C7	1.465 (3)	C12—C15	1.515 (4)
C2—C3	1.360 (3)	C13—C14	1.380 (4)
C2—H2	0.9300	C13—H13	0.9300
C3—C4	1.375 (4)	C14—H14	0.9300
C4—C5	1.369 (4)	C15—H15A	0.9600
C4—H4	0.9300	C15—H15B	0.9600
C5—C6	1.374 (4)	C15—H15C	0.9600
C5—H5	0.9300		
O2—N1—O1	123.4 (2)	O3—C8—N3	122.2 (2)
O2—N1—C3	118.7 (2)	O3—C8—C9	121.9 (2)
O1—N1—C3	117.8 (2)	N3—C8—C9	115.9 (2)
C7—N2—N3	115.6 (2)	C14—C9—C10	118.0 (3)
C8—N3—N2	120.0 (2)	C14—C9—C8	122.5 (2)
C8—N3—H3	119.5 (16)	C10—C9—C8	119.4 (2)
N2—N3—H3	119.7 (16)	C11—C10—C9	120.9 (3)
C2—C1—C6	118.6 (2)	C11—C10—H10	119.5
C2—C1—C7	121.5 (2)	C9—C10—H10	119.5
C6—C1—C7	119.9 (2)	C12—C11—C10	121.6 (3)
C3—C2—C1	119.2 (2)	C12—C11—H11	119.2
C3—C2—H2	120.4	C10—C11—H11	119.2
C1—C2—H2	120.4	C11—C12—C13	117.5 (3)
C2—C3—C4	122.8 (2)	C11—C12—C15	122.9 (3)
C2—C3—N1	118.6 (2)	C13—C12—C15	119.6 (3)
C4—C3—N1	118.6 (2)	C14—C13—C12	121.4 (3)
C5—C4—C3	118.3 (3)	C14—C13—H13	119.3
C5—C4—H4	120.9	C12—C13—H13	119.3
C3—C4—H4	120.9	C9—C14—C13	120.6 (3)
C4—C5—C6	120.0 (3)	C9—C14—H14	119.7
C4—C5—H5	120.0	C13—C14—H14	119.7
C6—C5—H5	120.0	C12—C15—H15A	109.5
C5—C6—C1	121.0 (2)	C12—C15—H15B	109.5
C5—C6—H6	119.5	H15A—C15—H15B	109.5
C1—C6—H6	119.5	C12—C15—H15C	109.5
N2—C7—C1	119.0 (2)	H15A—C15—H15C	109.5
N2—C7—H7	120.5	H15B—C15—H15C	109.5

---

C1—C7—H7                      120.5

---

*Hydrogen-bond geometry (Å, °)*

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
N3—H3...O3 <sup>i</sup>	0.89 (1)	2.13 (2)	2.909 (3)	146 (2)

---

Symmetry code: (i)  $x, y-1, z$ .