

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

(5Z)-5-(2-Hydroxybenzylidene)-3-(4-methylphenyl)-2-sulfanylidene-1,3-thiazolidin-4-one

Durre Shahwar,^a M. Nawaz Tahir,^{b*} Misbah Kashif,^a Afifa Saeed^a and Sana Bukhari^a

^aDepartment of Chemistry, Government College University, Lahore, Pakistan, and^bUniversity of Sargodha, Department of Physics, Sargodha, Pakistan

Correspondence e-mail: dmntahir_uos@yahoo.com


Received 11 May 2012; accepted 13 May 2012

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.047; wR factor = 0.101; data-to-parameter ratio = 13.9.

In the title compound, $\text{C}_{17}\text{H}_{13}\text{NO}_2\text{S}_2$, the dihedral angles between the 2-sulfanylidene-1,3-thiazolidin-4-one group and the pendant toluene and 2-hydroxybenzene rings are 74.62 (6) and 8.73 (12)°, respectively. An intramolecular $\text{C}-\text{H}\cdots\text{S}$ interaction occurs. In the crystal, inversion dimers linked by pairs of $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds generate $R_2^2(16)$ loops. This link is reinforced by a pair of $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds. The dimers are connected by weak $\text{C}-\text{H}\cdots\text{S}$ interactions.

Related literature

For related structures and further synthetic details, see: Shahwar *et al.* (2009a,b). For graph-set notation, see: Bernstein *et al.* (1995).



Experimental

Crystal data

$\text{C}_{17}\text{H}_{13}\text{NO}_2\text{S}_2$
 $M_r = 327.40$
 Monoclinic, $P2_1/c$
 $a = 13.8258$ (6) Å

$b = 5.4278$ (3) Å
 $c = 21.0715$ (9) Å
 $\beta = 101.857$ (3)°
 $V = 1547.54$ (13) Å³

$Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.35$ mm⁻¹

$T = 296$ K
 $0.35 \times 0.15 \times 0.13$ mm

Data collection

Bruker Kappa APEXII CCD diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2005)
 $T_{\min} = 0.945$, $T_{\max} = 0.965$

10652 measured reflections
 2801 independent reflections
 1473 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.069$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.047$
 $wR(F^2) = 0.101$
 $S = 0.93$
 2801 reflections

201 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.19$ e Å⁻³
 $\Delta\rho_{\min} = -0.26$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O2}-\text{H2A}\cdots\text{O1}^i$	0.82	1.92	2.712 (3)	162
$\text{C6}-\text{H6}\cdots\text{S2}^{ii}$	0.93	2.84	3.736 (4)	163
$\text{C11}-\text{H11}\cdots\text{O2}^i$	0.93	2.38	3.294 (4)	167
$\text{C13}-\text{H13}\cdots\text{S1}$	0.93	2.48	3.194 (3)	133

Symmetry codes: (i) $-x + 1, -y + 2, -z$; (ii) $x, y + 1, z$.

Data collection: APEX2 (Bruker, 2009); cell refinement: SAINT (Bruker, 2009); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 for Windows (Farrugia, 1997) and PLATON (Spek, 2009); software used to prepare material for publication: WinGX (Farrugia, 1999) and PLATON.

The authors acknowledge the provision of funds for the purchase of a diffractometer and encouragement by Dr Muhammad Akram Chaudhary, Vice Chancellor, University of Sargodha, Pakistan. The authors also acknowledge the technical support provided by Syed Muhammad Hussain Rizvi of Bana International, Karachi, Pakistan.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB6792).

References

- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2005). SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2009). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Shahwar, D., Tahir, M. N., Raza, M. A. & Iqbal, B. (2009a). *Acta Cryst.* **E65**, o2903.
- Shahwar, D., Tahir, M. N., Raza, M. A. & Iqbal, B. (2009b). *Acta Cryst.* **E65**, o2917.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

supporting information

Acta Cryst. (2012). E68, o1818 [doi:10.1107/S1600536812021630]

(5*Z*)-5-(2-Hydroxybenzylidene)-3-(4-methylphenyl)-2-sulfanylidene-1,3-thiazolidin-4-one

Durre Shahwar, M. Nawaz Tahir, Misbah Kashif, Afifa Saeed and Sana Bukhari

S1. Comment

In the search for new enzyme inhibitors, a series of rohdanine derivatives were prepared and their crystal structures have been reported such as (5*Z*)-5-(2-hydroxybenzylidene)-3-phenyl-2-sulfanylidene-1,3-thiazolidin-4-one (Shahwar *et al.*, 2009*a*) and (5*Z*)-5-(2-methylbenzylidene)-3-phenyl-2-sulfanylidene-1,3-thiazolidin-4-one (Shahwar *et al.*, 2009*b*) which are related to the title compound, (I), (Fig. 1).

In (I), the toluene group A (C1–C7), group B (N1/C8/S1/C10/C9/O1/S2) of 2-sulfanylidene-1,3-thiazolidin-4-one and group C (C11–C17/O2) of 2-methylphenol are planar with r. m. s. deviation of 0.0246 Å, 0.0186 Å and 0.0175 Å, respectively. The dihedral angle between A/B, A/C and B/C is 74.62 (6), 70.16 (7) and 8.73 (12)°, respectively. The molecules are dimerized due to strong O—H···O type of H-bonding with $R_2^2(16)$ (Bernstein *et al.*, 1995) ring motifs (Table 1, Fig. 2). $R_2^2(7)$ ring motifs are formed within the dimers when weak H-bonding of C—H···O type is considered. The dimers are interlinked due to C—H···S type of H-bondings (Table 1, Fig. 2).

S2. Experimental

3-(4-methylphenyl)-2-sulfanylidene-1,3-thiazolidin-4-one (0.419 g, 0.2 mol) (prepared according to the method: Shahwar *et al.* 2009*a*), 2-hydroxybenzaldehyde (0.244 g, 0.2 mol) and K_2CO_3 (0.553 g, 0.4 mol) were dissolved in distilled water (10 ml) and continuously stirred for 24 h. The mixture was neutralized with 5% HCl and the crude product obtained was recrystallized in ethylacetate to get the yellow needles of (I).

S3. Refinement

The H-atoms were positioned geometrically (C—H = 0.93–0.96 Å, O—H = 0.82 Å) and refined as riding with $U_{iso}(H) = xU_{eq}(C, O)$, where $x = 1.5$ for methyl groups and $x = 1.2$ for other H atoms.





Figure 1

View of the title compound with displacement ellipsoids drawn at the 50% probability level.

**Figure 2**

The partial packing, showing that molecules form dimers with $R_2^2(16)$ and other ring motifs.

(5Z)-5-(2-Hydroxybenzylidene)-3-(4-methylphenyl)-2-sulfanylidene-1,3-thiazolidin-4-one

Crystal data

$C_{17}H_{13}NO_2S_2$

$M_r = 327.40$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 13.8258(6) \text{ \AA}$

$b = 5.4278(3) \text{ \AA}$

$c = 21.0715(9) \text{ \AA}$

$\beta = 101.857(3)^\circ$

$V = 1547.54(13) \text{ \AA}^3$

$Z = 4$

$F(000) = 680$

$D_x = 1.405 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 1473 reflections

$\theta = 3.0\text{--}25.3^\circ$

$\mu = 0.35 \text{ mm}^{-1}$

$T = 296 \text{ K}$

Needle, yellow

$0.35 \times 0.15 \times 0.13 \text{ mm}$

Data collection

Bruker Kappa APEXII CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

Detector resolution: $8.20 \text{ pixels mm}^{-1}$

ω scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2005)

$T_{\min} = 0.945$, $T_{\max} = 0.965$

10652 measured reflections

2801 independent reflections

1473 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.069$

$\theta_{\max} = 25.3^\circ$, $\theta_{\min} = 3.0^\circ$

$h = -16 \rightarrow 16$

$k = -6 \rightarrow 6$

$l = -25 \rightarrow 25$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.047$
 $wR(F^2) = 0.101$
 $S = 0.93$
 2801 reflections
 201 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0385P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.19 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.26 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. Bond distances, angles *etc.* have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.32730 (6)	0.24641 (18)	0.08942 (4)	0.0555 (3)
S2	0.14244 (6)	-0.0283 (2)	0.04718 (4)	0.0633 (4)
O1	0.28996 (14)	0.6397 (4)	-0.06441 (10)	0.0510 (8)
O2	0.57469 (16)	1.0037 (5)	0.06997 (10)	0.0773 (10)
N1	0.21224 (16)	0.3302 (5)	-0.02022 (11)	0.0383 (9)
C1	0.1268 (2)	0.3173 (6)	-0.07351 (14)	0.0387 (11)
C2	0.1162 (2)	0.1227 (7)	-0.11500 (14)	0.0486 (13)
C3	0.0316 (2)	0.1030 (7)	-0.16312 (14)	0.0531 (14)
C4	-0.0423 (2)	0.2766 (7)	-0.16993 (15)	0.0486 (13)
C5	-0.0286 (2)	0.4723 (7)	-0.12792 (17)	0.0598 (14)
C6	0.0553 (2)	0.4943 (7)	-0.07940 (15)	0.0534 (12)
C7	-0.1379 (2)	0.2463 (7)	-0.21949 (16)	0.0805 (16)
C8	0.2200 (2)	0.1799 (6)	0.03366 (14)	0.0436 (11)
C9	0.2879 (2)	0.5003 (7)	-0.01944 (15)	0.0404 (11)
C10	0.3612 (2)	0.4829 (6)	0.04263 (13)	0.0408 (10)
C11	0.4379 (2)	0.6400 (6)	0.05800 (13)	0.0418 (11)
C12	0.5128 (2)	0.6614 (6)	0.11718 (13)	0.0391 (11)
C13	0.5208 (2)	0.5059 (7)	0.17093 (14)	0.0530 (14)
C14	0.5919 (2)	0.5422 (7)	0.22614 (15)	0.0592 (14)
C15	0.6561 (2)	0.7355 (7)	0.22960 (15)	0.0564 (14)
C16	0.6513 (2)	0.8925 (7)	0.17828 (15)	0.0520 (13)
C17	0.5808 (2)	0.8546 (6)	0.12202 (14)	0.0446 (11)
H2	0.16539	0.00361	-0.11109	0.0582*
H2A	0.62157	1.09941	0.07623	0.0928*
H3	0.02469	-0.03032	-0.19141	0.0637*
H5	-0.07693	0.59368	-0.13213	0.0718*

H6	0.06290	0.62792	-0.05118	0.0641*
H7A	-0.12882	0.12593	-0.25115	0.1208*
H7B	-0.15615	0.40109	-0.24058	0.1208*
H7C	-0.18916	0.19251	-0.19813	0.1208*
H11	0.44377	0.75226	0.02568	0.0501*
H13	0.47704	0.37487	0.16928	0.0640*
H14	0.59639	0.43528	0.26110	0.0709*
H15	0.70355	0.76061	0.26727	0.0677*
H16	0.69516	1.02395	0.18109	0.0625*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0601 (5)	0.0596 (7)	0.0403 (5)	-0.0257 (5)	-0.0047 (4)	0.0102 (5)
S2	0.0689 (6)	0.0671 (8)	0.0527 (5)	-0.0349 (6)	0.0096 (4)	0.0025 (5)
O1	0.0484 (13)	0.0539 (16)	0.0449 (13)	-0.0174 (12)	-0.0037 (11)	0.0132 (13)
O2	0.0795 (18)	0.087 (2)	0.0509 (14)	-0.0543 (16)	-0.0205 (12)	0.0243 (16)
N1	0.0362 (14)	0.0385 (18)	0.0362 (14)	-0.0112 (14)	-0.0015 (12)	0.0016 (14)
C1	0.0347 (17)	0.041 (2)	0.0385 (18)	-0.0079 (18)	0.0031 (15)	0.0005 (18)
C2	0.0429 (19)	0.053 (3)	0.0473 (19)	0.0042 (19)	0.0032 (17)	-0.006 (2)
C3	0.057 (2)	0.056 (3)	0.0422 (19)	-0.011 (2)	0.0009 (18)	-0.0158 (19)
C4	0.0417 (19)	0.056 (3)	0.0441 (19)	-0.011 (2)	-0.0002 (16)	0.011 (2)
C5	0.049 (2)	0.050 (3)	0.075 (2)	0.006 (2)	0.000 (2)	0.007 (2)
C6	0.053 (2)	0.043 (2)	0.060 (2)	-0.005 (2)	0.0019 (18)	-0.013 (2)
C7	0.055 (2)	0.112 (4)	0.061 (2)	-0.014 (2)	-0.0196 (18)	0.012 (3)
C8	0.0447 (18)	0.044 (2)	0.0401 (18)	-0.0132 (18)	0.0042 (15)	-0.0036 (18)
C9	0.0370 (18)	0.040 (2)	0.0431 (19)	-0.0070 (18)	0.0054 (16)	-0.0045 (19)
C10	0.0372 (17)	0.047 (2)	0.0363 (17)	-0.0089 (18)	0.0032 (14)	0.0034 (18)
C11	0.0417 (18)	0.044 (2)	0.0374 (17)	-0.0101 (18)	0.0026 (15)	0.0061 (17)
C12	0.0345 (17)	0.042 (2)	0.0379 (18)	-0.0092 (18)	0.0007 (15)	0.0006 (18)
C13	0.052 (2)	0.053 (3)	0.047 (2)	-0.014 (2)	-0.0059 (17)	0.008 (2)
C14	0.064 (2)	0.061 (3)	0.044 (2)	-0.012 (2)	-0.0091 (18)	0.014 (2)
C15	0.056 (2)	0.066 (3)	0.0393 (19)	-0.008 (2)	-0.0089 (17)	-0.002 (2)
C16	0.0414 (18)	0.062 (3)	0.046 (2)	-0.0162 (19)	-0.0066 (16)	-0.002 (2)
C17	0.0452 (19)	0.047 (2)	0.0386 (18)	-0.0076 (19)	0.0016 (16)	0.0036 (19)

Geometric parameters (Å, °)

S1—C8	1.731 (3)	C12—C17	1.398 (4)
S1—C10	1.741 (3)	C12—C13	1.399 (4)
S2—C8	1.623 (3)	C13—C14	1.374 (4)
O1—C9	1.218 (4)	C14—C15	1.367 (5)
O2—C17	1.352 (4)	C15—C16	1.368 (5)
O2—H2A	0.8200	C16—C17	1.387 (4)
N1—C1	1.455 (4)	C2—H2	0.9300
N1—C9	1.393 (4)	C3—H3	0.9300
N1—C8	1.384 (4)	C5—H5	0.9300
C1—C2	1.360 (5)	C6—H6	0.9300

C1—C6	1.366 (5)	C7—H7A	0.9600
C2—C3	1.386 (4)	C7—H7B	0.9600
C3—C4	1.376 (5)	C7—H7C	0.9600
C4—C5	1.371 (5)	C11—H11	0.9300
C4—C7	1.516 (4)	C13—H13	0.9300
C5—C6	1.385 (4)	C14—H14	0.9300
C9—C10	1.484 (4)	C15—H15	0.9300
C10—C11	1.347 (4)	C16—H16	0.9300
C11—C12	1.453 (4)		
C8—S1—C10	93.57 (14)	C14—C15—C16	120.7 (3)
C17—O2—H2A	109.00	C15—C16—C17	119.7 (3)
C1—N1—C8	121.2 (2)	O2—C17—C16	121.4 (3)
C1—N1—C9	122.0 (2)	C12—C17—C16	121.2 (3)
C8—N1—C9	116.7 (2)	O2—C17—C12	117.5 (3)
N1—C1—C6	119.6 (3)	C1—C2—H2	120.00
C2—C1—C6	120.5 (3)	C3—C2—H2	120.00
N1—C1—C2	119.8 (3)	C2—C3—H3	119.00
C1—C2—C3	119.6 (3)	C4—C3—H3	119.00
C2—C3—C4	121.4 (3)	C4—C5—H5	119.00
C3—C4—C7	121.4 (3)	C6—C5—H5	119.00
C5—C4—C7	121.0 (3)	C1—C6—H6	120.00
C3—C4—C5	117.5 (3)	C5—C6—H6	120.00
C4—C5—C6	121.8 (3)	C4—C7—H7A	109.00
C1—C6—C5	119.2 (3)	C4—C7—H7B	109.00
S1—C8—S2	122.09 (18)	C4—C7—H7C	109.00
S1—C8—N1	110.2 (2)	H7A—C7—H7B	109.00
S2—C8—N1	127.7 (2)	H7A—C7—H7C	109.00
O1—C9—N1	122.7 (3)	H7B—C7—H7C	110.00
O1—C9—C10	127.2 (3)	C10—C11—H11	115.00
N1—C9—C10	110.1 (3)	C12—C11—H11	115.00
S1—C10—C11	128.4 (2)	C12—C13—H13	119.00
C9—C10—C11	122.2 (3)	C14—C13—H13	119.00
S1—C10—C9	109.4 (2)	C13—C14—H14	120.00
C10—C11—C12	130.1 (3)	C15—C14—H14	120.00
C11—C12—C17	118.2 (3)	C14—C15—H15	120.00
C13—C12—C17	117.0 (3)	C16—C15—H15	120.00
C11—C12—C13	124.8 (3)	C15—C16—H16	120.00
C12—C13—C14	121.5 (3)	C17—C16—H16	120.00
C13—C14—C15	119.9 (3)		
C10—S1—C8—S2	-177.0 (2)	C3—C4—C5—C6	-1.2 (5)
C10—S1—C8—N1	1.8 (2)	C7—C4—C5—C6	175.5 (3)
C8—S1—C10—C9	-2.5 (2)	C4—C5—C6—C1	0.5 (5)
C8—S1—C10—C11	174.7 (3)	O1—C9—C10—S1	-177.6 (3)
C8—N1—C1—C2	-73.7 (4)	O1—C9—C10—C11	5.0 (5)
C8—N1—C1—C6	102.9 (4)	N1—C9—C10—S1	2.6 (3)
C9—N1—C1—C2	109.1 (4)	N1—C9—C10—C11	-174.8 (3)

C9—N1—C1—C6	-74.3 (4)	S1—C10—C11—C12	-1.0 (5)
C1—N1—C8—S1	-177.9 (2)	C9—C10—C11—C12	175.9 (3)
C1—N1—C8—S2	0.9 (4)	C10—C11—C12—C13	2.8 (5)
C9—N1—C8—S1	-0.6 (3)	C10—C11—C12—C17	-175.5 (3)
C9—N1—C8—S2	178.2 (2)	C11—C12—C13—C14	-177.9 (3)
C1—N1—C9—O1	-3.8 (5)	C17—C12—C13—C14	0.4 (5)
C1—N1—C9—C10	176.0 (3)	C11—C12—C17—O2	-2.6 (4)
C8—N1—C9—O1	178.9 (3)	C11—C12—C17—C16	177.0 (3)
C8—N1—C9—C10	-1.4 (4)	C13—C12—C17—O2	179.0 (3)
N1—C1—C2—C3	175.8 (3)	C13—C12—C17—C16	-1.5 (4)
C6—C1—C2—C3	-0.7 (5)	C12—C13—C14—C15	0.7 (5)
N1—C1—C6—C5	-176.1 (3)	C13—C14—C15—C16	-0.8 (5)
C2—C1—C6—C5	0.5 (5)	C14—C15—C16—C17	-0.3 (5)
C1—C2—C3—C4	0.0 (5)	C15—C16—C17—O2	-179.0 (3)
C2—C3—C4—C5	1.0 (5)	C15—C16—C17—C12	1.4 (5)
C2—C3—C4—C7	-175.7 (3)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O2—H2A \cdots O1 ⁱ	0.82	1.92	2.712 (3)	162
C6—H6 \cdots S2 ⁱⁱ	0.93	2.84	3.736 (4)	163
C11—H11 \cdots O2 ⁱ	0.93	2.38	3.294 (4)	167
C13—H13 \cdots S1	0.93	2.48	3.194 (3)	133

Symmetry codes: (i) $-x+1, -y+2, -z$; (ii) $x, y+1, z$.