

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

# *N*'<sup>2</sup>,*N*'<sup>5</sup>-Bis[(*E*)-2-hydroxybenzylidene]-3,4-dimethylthiophene-2,5-dicarbohydrazide

Shao-Lin Zhang, Ling Zhang, Qin-Mei Wen, Rong-Xia Geng\* and Cheng-He Zhou\*

Laboratory of Bioorganic & Medicinal Chemistry, School of Chemistry and Chemical Engineering, Southwest University, Chongqing 400715, People's Republic of China  
Correspondence e-mail: geng0712@swu.edu.cn, zhouch@swu.edu.cn


Received 23 April 2012; accepted 5 May 2012

Key indicators: single-crystal X-ray study;  $T = 296$  K; mean  $\sigma(\text{C}-\text{C}) = 0.004$  Å;  $R$  factor = 0.047;  $wR$  factor = 0.128; data-to-parameter ratio = 13.9.

In the title molecule,  $\text{C}_{22}\text{H}_{20}\text{N}_4\text{O}_4\text{S}$ , both  $\text{C}=\text{N}$  bonds are in an *E* conformation. The benzene rings form dihedral angles of 12.10 (13) and 25.17 (12)° with the thiophene ring. The dihedral angle between the two benzene rings is 17.59 (14)°. There are two intramolecular  $\text{O}-\text{H}\cdots\text{N}$  hydrogen bonds. In the crystal,  $\text{N}-\text{H}\cdots\text{O}$  hydrogen bonds connect molecules into chains along [010].

## Related literature

For the medicinal properties of thiophene derivatives, see: Bondock *et al.* (2010); Geng & Zhou (2008). For a related structure, see: Tang *et al.* (2010).



## Experimental

### Crystal data

$\text{C}_{22}\text{H}_{20}\text{N}_4\text{O}_4\text{S}$   
 $M_r = 436.48$   
Triclinic,  $P\bar{1}$   
 $a = 8.392$  (8) Å

$b = 9.511$  (9) Å  
 $c = 12.937$  (8) Å  
 $\alpha = 99.853$  (17)°  
 $\beta = 90.804$  (18)°

$\gamma = 92.374$  (18)°  
 $V = 1016.2$  (15) Å<sup>3</sup>  
 $Z = 2$   
Mo  $K\alpha$  radiation

$\mu = 0.20$  mm<sup>-1</sup>  
 $T = 296$  K  
 $0.19 \times 0.18 \times 0.16$  mm

### Data collection

Bruker APEXII CCD diffractometer  
Absorption correction: multi-scan (SADABS; Bruker, 2009)  
 $T_{\min} = 0.963$ ,  $T_{\max} = 0.969$

5573 measured reflections  
3925 independent reflections  
2660 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.019$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.047$   
 $wR(F^2) = 0.128$   
 $S = 1.02$   
3925 reflections

283 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.23$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.30$  e Å<sup>-3</sup>

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O1}-\text{H1}\cdots\text{N1}$	0.82	1.91	2.626 (4)	145
$\text{N2}-\text{H2}\cdots\text{O3}^{\text{i}}$	0.86	1.97	2.789 (4)	159
$\text{N4}-\text{H4}\cdots\text{O2}^{\text{ii}}$	0.86	1.97	2.812 (4)	165
$\text{O4}-\text{H4B}\cdots\text{N3}$	0.82	1.85	2.569 (4)	146

Symmetry codes: (i)  $-x + 1, -y + 1, -z + 1$ ; (ii)  $-x + 1, -y, -z + 1$ .

Data collection: APEX2 (Bruker, 2009); cell refinement: SAINT (Bruker, 2009); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

This work was partially supported by the National Natural Science Foundation of China (No. 21172181), the Key Program of the Natural Science Foundation of Chongqing (CSTC2012jjB10026), the Specialized Research Fund for the Doctoral Program of Higher Education of China (SRFDP 20110182110007) and the Research Funds for the Central Universities (XDJK2011D007, XDJK2012B026).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH5463).

## References

- Bondock, S., Metwally, M. A. & Fadaly, W. (2010). *Eur. J. Med. Chem.* **45**, 3692–3701.  
Bruker (2009). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.  
Geng, R.-X. & Zhou, C.-H. (2008). *Chin. J. Org. Chem.* **24**, 163–168.  
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
Tang, Y.-D., Geng, R.-X. & Zhou, C.-H. (2010). *Acta Cryst.* **E66**, o100.

## supporting information

*Acta Cryst.* (2012). E68, o1752 [doi:10.1107/S1600536812020260]

## ***N*'<sup>2</sup>,*N*'<sup>5</sup>-Bis[(*E*)-2-hydroxybenzylidene]-3,4-dimethylthiophene-2,5-dicarbohydrazide**

**Shao-Lin Zhang, Ling Zhang, Qin-Mei Wen, Rong-Xia Geng and Cheng-He Zhou**

### **S1. Comment**

Thiophene is a electron-rich five-membered aromatic heterocycle containing a sulfur atom whose derivatives display various biological activities. Much effort has been devoted to the researches of thiophene-based compounds as medicinal agents (Geng *et al.*, 2008; Bondock *et al.*, 2010). Our interest is to develop novel thiophene compounds with high bioactivities especially broad antimicrobial spectrum. We have already prepared a thiophene compound incorporating Schiff base moieties and determined its crystal structure (Tang *et al.*, 2010). Herein, the crystal structure of title compound (I) is reported.


The molecular structure of (I) is shown in Fig. 1. Both C=N bonds are in an *E* conformation. The benzene rings form dihedral angles of 12.10 (13)° (C17-C22) and 25.17 (12)° (C1-C6) with the thiophene ring (S1/C9/C10/C12/C14). The dihedral angle between the two benzene rings is 17.59 (14)°. There are two intramolecular O—H⋯N hydrogen bonds. In the crystal, N—H⋯O hydrogen bonds connect molecules into chains along [010].

### **S2. Experimental**

A mixture of 3,4-dimethylthiophene-2,5-dicarbohydrazide (0.11 g, 0.5 mmol) and 2-hydroxybenzaldehyde (0.24 g, 2 mmol) in methanol (10.0 ml) was stirred at room temperature. Upon the completion of the reaction (monitored by TLC, eluent, ethyl acetate), the formed precipitate was filtered and then washed with cold methanol to afford a yellow solid of the title compound (0.35 g). A crystal suitable for X-ray analysis was grown from a mixed solution of (I) in chloroform and methanol by slow evaporation at room temperature.

### **S3. Refinement**

H atoms were placed in calculated positions with C—H = 0.93 Å (aromatic), 0.96 Å (methyl), N—H = 0.86 Å and O—H = 0.82 Å. The  $U_{\text{iso}}(\text{H})$  values were set equal to  $1.2U_{\text{eq}}(\text{C}_{\text{aromatic}}, \text{N})$  and  $1.5U_{\text{eq}}(\text{C}_{\text{methyl}}, \text{O})$ .

**Figure 1**

The molecular structure of (I), showing the displacement ellipsoids drawn at the 50% probability level.

### *N'*<sup>2</sup>,*N'*<sup>5</sup>-Bis[(*E*)-2-hydroxybenzylidene]-3,4-dimethylthiophene-2,5-dicarbohydrazide

#### Crystal data

$C_{22}H_{20}N_4O_4S$

$M_r = 436.48$

Triclinic,  $P\bar{1}$

Hall symbol:  $-P\ 1$

$a = 8.392\ (8)\ \text{\AA}$

$b = 9.511\ (9)\ \text{\AA}$

$c = 12.937\ (8)\ \text{\AA}$

$\alpha = 99.853\ (17)^\circ$

$\beta = 90.804\ (18)^\circ$

$\gamma = 92.374\ (18)^\circ$

$V = 1016.2\ (15)\ \text{\AA}^3$

$Z = 2$

$F(000) = 456$

$D_x = 1.426\ \text{Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 1480 reflections

$\theta = 2.9\text{--}24.4^\circ$

$\mu = 0.20\ \text{mm}^{-1}$

$T = 296\ \text{K}$

Block, yellow

$0.19 \times 0.18 \times 0.16\ \text{mm}$

#### Data collection

Bruker APEXII CCD  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

$\varphi$  and  $\omega$  scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2009)

$T_{\min} = 0.963$ ,  $T_{\max} = 0.969$

5573 measured reflections

3925 independent reflections

2660 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.019$

$\theta_{\max} = 26.0^\circ$ ,  $\theta_{\min} = 2.4^\circ$

$h = -9 \rightarrow 10$

$k = -11 \rightarrow 10$

$l = -15 \rightarrow 15$

#### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.047$

$wR(F^2) = 0.128$

$S = 1.02$

3925 reflections

283 parameters

0 restraints

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier

map

Hydrogen site location: inferred from  
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0604P)^2 + 0.0881P]$

where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.23\ \text{e}\ \text{\AA}^{-3}$

$\Delta\rho_{\min} = -0.30\ \text{e}\ \text{\AA}^{-3}$

Extinction correction: *SHELXL97* (Sheldrick,  
2008),  $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$

Extinction coefficient: 0.0062 (17)

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.9005 (2)	0.2862 (2)	0.73465 (15)	0.0457 (5)
N2	0.8176 (2)	0.32288 (19)	0.65213 (15)	0.0463 (6)
H2	0.8112	0.4109	0.6455	0.056*
N3	0.0145 (2)	0.15155 (19)	0.30284 (15)	0.0423 (5)
N4	0.1633 (2)	0.16130 (19)	0.34807 (16)	0.0448 (5)
H4	0.1990	0.0926	0.3759	0.054*
C1	1.0429 (3)	0.2314 (3)	0.92331 (19)	0.0473 (6)
C2	1.1287 (4)	0.2087 (3)	1.0091 (2)	0.0631 (8)
H2A	1.1110	0.1251	1.0363	0.076*
C3	1.2397 (4)	0.3085 (4)	1.0542 (2)	0.0702 (9)
H3A	1.2992	0.2919	1.1118	0.084*
C4	1.2661 (4)	0.4336 (4)	1.0168 (2)	0.0658 (8)
H4A	1.3424	0.5014	1.0488	0.079*
C5	1.1795 (3)	0.4575 (3)	0.9320 (2)	0.0524 (7)
H5A	1.1958	0.5429	0.9071	0.063*
C6	1.0681 (3)	0.3567 (3)	0.88285 (18)	0.0424 (6)
C7	0.9818 (3)	0.3853 (3)	0.79290 (18)	0.0432 (6)
H7A	0.9854	0.4772	0.7771	0.052*
C8	0.7473 (3)	0.2195 (2)	0.5826 (2)	0.0438 (6)
C9	0.6441 (3)	0.2671 (2)	0.50316 (18)	0.0392 (6)
C10	0.6698 (3)	0.3655 (2)	0.44024 (17)	0.0365 (5)
C11	0.8216 (3)	0.4475 (3)	0.4339 (2)	0.0482 (6)
H11A	0.8974	0.4238	0.4837	0.072*
H11B	0.8035	0.5479	0.4493	0.072*
H11C	0.8626	0.4242	0.3644	0.072*
C12	0.5355 (3)	0.3773 (2)	0.37593 (16)	0.0348 (5)
C13	0.5400 (3)	0.4737 (3)	0.29632 (19)	0.0482 (6)
H13A	0.4382	0.4681	0.2607	0.072*
H13B	0.6210	0.4448	0.2464	0.072*
H13C	0.5636	0.5702	0.3306	0.072*
C14	0.4126 (3)	0.2878 (2)	0.39320 (17)	0.0356 (5)
C15	0.2514 (3)	0.2808 (2)	0.34768 (17)	0.0364 (5)
C16	-0.0755 (3)	0.0451 (2)	0.31184 (19)	0.0454 (6)
H16A	-0.0433	-0.0200	0.3532	0.055*
C17	-0.2275 (3)	0.0243 (3)	0.25834 (19)	0.0452 (6)

C18	-0.2791 (3)	0.1158 (3)	0.1937 (2)	0.0490 (7)
C19	-0.4226 (4)	0.0886 (4)	0.1412 (2)	0.0686 (9)
H19A	-0.4569	0.1500	0.0975	0.082*
C20	-0.5150 (4)	-0.0277 (4)	0.1529 (3)	0.0798 (11)
H20A	-0.6133	-0.0446	0.1179	0.096*
C21	-0.4657 (4)	-0.1201 (4)	0.2151 (3)	0.0795 (10)
H21A	-0.5291	-0.2005	0.2218	0.095*
C22	-0.3236 (4)	-0.0936 (3)	0.2669 (2)	0.0639 (8)
H22A	-0.2900	-0.1568	0.3094	0.077*
O1	0.9338 (3)	0.13019 (19)	0.88189 (15)	0.0640 (6)
H1	0.8908	0.1536	0.8305	0.096*
O2	0.7611 (2)	0.09444 (16)	0.58491 (16)	0.0683 (6)
O3	0.1986 (2)	0.38046 (15)	0.31228 (12)	0.0443 (4)
O4	-0.1926 (2)	0.2321 (2)	0.17967 (17)	0.0681 (6)
H4B	-0.1073	0.2355	0.2119	0.102*
S1	0.46125 (8)	0.18434 (6)	0.48365 (5)	0.0450 (2)

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
N1	0.0422 (13)	0.0467 (11)	0.0497 (12)	0.0023 (10)	-0.0194 (10)	0.0140 (9)
N2	0.0483 (13)	0.0348 (10)	0.0563 (13)	-0.0007 (9)	-0.0264 (11)	0.0116 (9)
N3	0.0329 (11)	0.0371 (10)	0.0555 (13)	0.0001 (9)	-0.0143 (10)	0.0054 (9)
N4	0.0351 (12)	0.0350 (10)	0.0654 (14)	-0.0012 (9)	-0.0191 (10)	0.0139 (9)
C1	0.0444 (16)	0.0562 (15)	0.0425 (14)	0.0076 (13)	-0.0018 (12)	0.0105 (12)
C2	0.068 (2)	0.0745 (19)	0.0507 (17)	0.0153 (17)	-0.0077 (15)	0.0200 (15)
C3	0.066 (2)	0.100 (2)	0.0455 (17)	0.022 (2)	-0.0139 (15)	0.0112 (17)
C4	0.0513 (18)	0.090 (2)	0.0512 (17)	0.0001 (17)	-0.0179 (14)	-0.0003 (16)
C5	0.0446 (16)	0.0620 (16)	0.0495 (15)	-0.0029 (13)	-0.0072 (13)	0.0081 (13)
C6	0.0372 (14)	0.0525 (14)	0.0378 (13)	0.0056 (12)	-0.0054 (11)	0.0087 (11)
C7	0.0417 (15)	0.0443 (13)	0.0445 (14)	0.0030 (12)	-0.0087 (12)	0.0103 (11)
C8	0.0358 (14)	0.0387 (12)	0.0574 (15)	-0.0030 (11)	-0.0172 (12)	0.0126 (11)
C9	0.0329 (13)	0.0339 (11)	0.0493 (14)	-0.0015 (10)	-0.0148 (11)	0.0053 (10)
C10	0.0332 (13)	0.0350 (11)	0.0390 (13)	0.0008 (10)	-0.0045 (10)	0.0007 (10)
C11	0.0361 (14)	0.0584 (15)	0.0488 (15)	-0.0058 (12)	-0.0059 (12)	0.0083 (12)
C12	0.0338 (13)	0.0354 (11)	0.0338 (12)	0.0009 (10)	-0.0044 (10)	0.0031 (9)
C13	0.0444 (15)	0.0530 (14)	0.0494 (15)	-0.0021 (12)	-0.0073 (12)	0.0169 (12)
C14	0.0347 (13)	0.0327 (11)	0.0390 (13)	0.0016 (10)	-0.0090 (10)	0.0057 (9)
C15	0.0369 (13)	0.0328 (11)	0.0380 (12)	0.0016 (10)	-0.0082 (10)	0.0028 (10)
C16	0.0406 (15)	0.0412 (13)	0.0547 (15)	-0.0008 (12)	-0.0119 (12)	0.0107 (11)
C17	0.0340 (14)	0.0495 (14)	0.0494 (15)	-0.0022 (12)	-0.0052 (12)	0.0028 (12)
C18	0.0382 (15)	0.0529 (15)	0.0542 (16)	0.0047 (13)	-0.0055 (12)	0.0039 (12)
C19	0.0429 (18)	0.091 (2)	0.069 (2)	0.0100 (18)	-0.0190 (15)	0.0056 (17)
C20	0.0352 (17)	0.114 (3)	0.079 (2)	-0.0092 (19)	-0.0133 (16)	-0.011 (2)
C21	0.050 (2)	0.096 (2)	0.086 (2)	-0.0327 (18)	-0.0060 (18)	0.008 (2)
C22	0.0540 (19)	0.0697 (18)	0.0668 (19)	-0.0183 (15)	-0.0059 (15)	0.0136 (15)
O1	0.0706 (15)	0.0566 (11)	0.0683 (14)	-0.0086 (11)	-0.0167 (11)	0.0253 (10)
O2	0.0728 (15)	0.0334 (9)	0.0996 (15)	-0.0046 (9)	-0.0467 (12)	0.0197 (9)

O3	0.0427 (10)	0.0366 (8)	0.0548 (10)	0.0001 (8)	-0.0179 (8)	0.0135 (7)
O4	0.0557 (13)	0.0601 (11)	0.0935 (16)	0.0018 (10)	-0.0228 (11)	0.0293 (11)
S1	0.0398 (4)	0.0391 (3)	0.0577 (4)	-0.0084 (3)	-0.0212 (3)	0.0167 (3)

*Geometric parameters (Å, °)*

N1—C7	1.270 (3)	C11—H11A	0.9600
N1—N2	1.368 (3)	C11—H11B	0.9600
N2—C8	1.326 (3)	C11—H11C	0.9600
N2—H2	0.8600	C12—C14	1.355 (3)
N3—C16	1.261 (3)	C12—C13	1.491 (3)
N3—N4	1.365 (3)	C13—H13A	0.9600
N4—C15	1.331 (3)	C13—H13B	0.9600
N4—H4	0.8600	C13—H13C	0.9600
C1—O1	1.337 (3)	C14—C15	1.462 (3)
C1—C2	1.368 (3)	C14—S1	1.708 (2)
C1—C6	1.391 (4)	C15—O3	1.217 (3)
C2—C3	1.354 (4)	C16—C17	1.432 (3)
C2—H2A	0.9300	C16—H16A	0.9300
C3—C4	1.370 (4)	C17—C22	1.375 (4)
C3—H3A	0.9300	C17—C18	1.384 (4)
C4—C5	1.364 (4)	C18—O4	1.338 (3)
C4—H4A	0.9300	C18—C19	1.367 (4)
C5—C6	1.377 (3)	C19—C20	1.355 (5)
C5—H5A	0.9300	C19—H19A	0.9300
C6—C7	1.433 (3)	C20—C21	1.361 (5)
C7—H7A	0.9300	C20—H20A	0.9300
C8—O2	1.205 (3)	C21—C22	1.354 (4)
C8—C9	1.476 (3)	C21—H21A	0.9300
C9—C10	1.355 (3)	C22—H22A	0.9300
C9—S1	1.690 (3)	O1—H1	0.8200
C10—C12	1.411 (3)	O4—H4B	0.8200
C10—C11	1.477 (4)		
C7—N1—N2	117.0 (2)	H11A—C11—H11C	109.5
C8—N2—N1	118.42 (19)	H11B—C11—H11C	109.5
C8—N2—H2	120.8	C14—C12—C10	111.9 (2)
N1—N2—H2	120.8	C14—C12—C13	126.7 (2)
C16—N3—N4	118.1 (2)	C10—C12—C13	121.3 (2)
C15—N4—N3	117.69 (19)	C12—C13—H13A	109.5
C15—N4—H4	121.2	C12—C13—H13B	109.5
N3—N4—H4	121.2	H13A—C13—H13B	109.5
O1—C1—C2	117.5 (3)	C12—C13—H13C	109.5
O1—C1—C6	122.1 (2)	H13A—C13—H13C	109.5
C2—C1—C6	120.4 (3)	H13B—C13—H13C	109.5
C3—C2—C1	119.6 (3)	C12—C14—C15	126.6 (2)
C3—C2—H2A	120.2	C12—C14—S1	112.18 (17)
C1—C2—H2A	120.2	C15—C14—S1	121.11 (18)

C2—C3—C4	121.3 (3)	O3—C15—N4	121.2 (2)
C2—C3—H3A	119.4	O3—C15—C14	121.7 (2)
C4—C3—H3A	119.4	N4—C15—C14	117.08 (19)
C5—C4—C3	119.3 (3)	N3—C16—C17	119.9 (2)
C5—C4—H4A	120.3	N3—C16—H16A	120.0
C3—C4—H4A	120.3	C17—C16—H16A	120.0
C4—C5—C6	120.9 (3)	C22—C17—C18	118.1 (3)
C4—C5—H5A	119.6	C22—C17—C16	119.6 (2)
C6—C5—H5A	119.6	C18—C17—C16	122.3 (2)
C5—C6—C1	118.5 (2)	O4—C18—C19	117.7 (3)
C5—C6—C7	119.0 (2)	O4—C18—C17	122.2 (2)
C1—C6—C7	122.6 (2)	C19—C18—C17	120.1 (3)
N1—C7—C6	120.5 (2)	C20—C19—C18	120.0 (3)
N1—C7—H7A	119.8	C20—C19—H19A	120.0
C6—C7—H7A	119.8	C18—C19—H19A	120.0
O2—C8—N2	123.2 (2)	C19—C20—C21	120.9 (3)
O2—C8—C9	121.3 (2)	C19—C20—H20A	119.6
N2—C8—C9	115.5 (2)	C21—C20—H20A	119.6
C10—C9—C8	131.5 (2)	C22—C21—C20	119.2 (3)
C10—C9—S1	112.63 (17)	C22—C21—H21A	120.4
C8—C9—S1	115.89 (19)	C20—C21—H21A	120.4
C9—C10—C12	112.1 (2)	C21—C22—C17	121.6 (3)
C9—C10—C11	125.2 (2)	C21—C22—H22A	119.2
C12—C10—C11	122.6 (2)	C17—C22—H22A	119.2
C10—C11—H11A	109.5	C1—O1—H1	109.5
C10—C11—H11B	109.5	C18—O4—H4B	109.5
H11A—C11—H11B	109.5	C9—S1—C14	91.09 (12)
C10—C11—H11C	109.5		
C7—N1—N2—C8	172.0 (2)	C11—C10—C12—C13	0.5 (3)
C16—N3—N4—C15	-173.2 (2)	C10—C12—C14—C15	174.1 (2)
O1—C1—C2—C3	-179.5 (3)	C13—C12—C14—C15	-9.0 (4)
C6—C1—C2—C3	-0.5 (4)	C10—C12—C14—S1	-2.3 (2)
C1—C2—C3—C4	1.1 (5)	C13—C12—C14—S1	174.64 (18)
C2—C3—C4—C5	-0.4 (5)	N3—N4—C15—O3	3.1 (3)
C3—C4—C5—C6	-1.1 (4)	N3—N4—C15—C14	-178.0 (2)
C4—C5—C6—C1	1.6 (4)	C12—C14—C15—O3	-20.8 (4)
C4—C5—C6—C7	-178.9 (2)	S1—C14—C15—O3	155.27 (18)
O1—C1—C6—C5	178.1 (2)	C12—C14—C15—N4	160.2 (2)
C2—C1—C6—C5	-0.9 (4)	S1—C14—C15—N4	-23.7 (3)
O1—C1—C6—C7	-1.3 (4)	N4—N3—C16—C17	-174.8 (2)
C2—C1—C6—C7	179.7 (2)	N3—C16—C17—C22	178.7 (2)
N2—N1—C7—C6	178.1 (2)	N3—C16—C17—C18	1.8 (4)
C5—C6—C7—N1	167.1 (2)	C22—C17—C18—O4	-179.2 (3)
C1—C6—C7—N1	-13.4 (4)	C16—C17—C18—O4	-2.2 (4)
N1—N2—C8—O2	-4.8 (4)	C22—C17—C18—C19	0.6 (4)
N1—N2—C8—C9	172.8 (2)	C16—C17—C18—C19	177.6 (2)
O2—C8—C9—C10	-133.2 (3)	O4—C18—C19—C20	-179.9 (3)

N2—C8—C9—C10	49.2 (4)	C17—C18—C19—C20	0.3 (4)
O2—C8—C9—S1	45.7 (3)	C18—C19—C20—C21	-1.1 (5)
N2—C8—C9—S1	-132.0 (2)	C19—C20—C21—C22	1.0 (5)
C8—C9—C10—C12	-179.5 (2)	C20—C21—C22—C17	0.0 (5)
S1—C9—C10—C12	1.6 (2)	C18—C17—C22—C21	-0.8 (4)
C8—C9—C10—C11	3.4 (4)	C16—C17—C22—C21	-177.8 (3)
S1—C9—C10—C11	-175.46 (18)	C10—C9—S1—C14	-2.47 (18)
C9—C10—C12—C14	0.4 (3)	C8—C9—S1—C14	178.49 (18)
C11—C10—C12—C14	177.6 (2)	C12—C14—S1—C9	2.72 (17)
C9—C10—C12—C13	-176.7 (2)	C15—C14—S1—C9	-173.90 (18)

*Hydrogen-bond geometry (Å, °)*

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
O1—H1...N1	0.82	1.91	2.626 (4)	145
N2—H2...O3 <sup>i</sup>	0.86	1.97	2.789 (4)	159
N4—H4...O2 <sup>ii</sup>	0.86	1.97	2.812 (4)	165
O4—H4B...N3	0.82	1.85	2.569 (4)	146

Symmetry codes: (i)  $-x+1, -y+1, -z+1$ ; (ii)  $-x+1, -y, -z+1$ .