

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

N,N'-Bis(4-methylphenyl)-*N''*-(2,2,2-trichloroacetyl)phosphoric triamide

 Akbar Raissi Shabari,^{a*} Mehرداد Pourayoubi,^b Hassan Fadaei,^b Marek Nečas^c and Michal Babiak^c
^aFaculty of Chemistry, North Tehran Branch, Islamic Azad University, Tehran, Iran,

^bDepartment of Chemistry, Ferdowsi University of Mashhad, Mashhad, Iran, and

^cDepartment of Chemistry, Faculty of Science, Masaryk University, Kotlarska 2, Brno CZ-61137, Czech Republic

Correspondence e-mail: a.raissi_shabari@yahoo.com


Received 28 April 2012; accepted 11 May 2012

 Key indicators: single-crystal X-ray study; $T = 120$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.028; wR factor = 0.072; data-to-parameter ratio = 13.6.

The P atom in the title compound, $\text{C}_{16}\text{H}_{17}\text{Cl}_3\text{N}_3\text{O}_2\text{P}$, is bonded in a distorted tetrahedral geometry with the phosphoryl and carbonyl groups *anti* with respect to one another. In the crystal, molecules are linked through $(\text{N}-\text{H})_2 \cdots \text{O}(\text{=P})$ and $\text{N}-\text{H} \cdots \text{O}(\text{=C})$ hydrogen bonds into chains along [001]. The phosphoryl O atom acts as a double hydrogen-bond acceptor.

Related literature

For phosphoric triamides having a $\text{C}(\text{=O})\text{NHP}(\text{=O})$ skeleton, see: Pourayoubi *et al.* (2011). For the definition of a double hydrogen-bond acceptor, see: Steiner (2002); Pourayoubi *et al.* (2012).



Experimental

Crystal data

 $\text{C}_{16}\text{H}_{17}\text{Cl}_3\text{N}_3\text{O}_2\text{P}$
 $M_r = 420.65$

 Monoclinic, $P2_1/c$
 $a = 17.5151$ (6) Å
 $b = 10.8638$ (4) Å
 $c = 9.8615$ (3) Å
 $\beta = 97.565$ (3)°
 $V = 1860.12$ (11) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.59$ mm⁻¹
 $T = 120$ K
 $0.60 \times 0.60 \times 0.60$ mm

Data collection

 Oxford Diffraction Xcalibur
 Sapphire2 diffractometer
 Absorption correction: multi-scan
 (*CrysAlis RED*; Oxford
 Diffraction, 2009)
 $T_{\min} = 0.955$, $T_{\max} = 1.000$

 6796 measured reflections
 3265 independent reflections
 2820 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.013$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.028$
 $wR(F^2) = 0.072$
 $S = 1.04$
 3265 reflections
 240 parameters

 H atoms treated by a mixture of
 independent and constrained
 refinement
 $\Delta\rho_{\text{max}} = 0.33$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.26$ e Å⁻³
Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1N}\cdots\text{O1}^{\text{i}}$	0.77 (2)	2.17 (2)	2.8953 (19)	156 (2)
$\text{N2}-\text{H2N}\cdots\text{O1}^{\text{i}}$	0.76 (2)	2.23 (2)	2.948 (2)	159 (2)
$\text{N3}-\text{H3N}\cdots\text{O2}^{\text{ii}}$	0.75 (2)	2.31 (2)	3.008 (2)	157 (2)

 Symmetry codes: (i) $x, -y + \frac{3}{2}, z + \frac{1}{2}$; (ii) $x, -y + \frac{3}{2}, z - \frac{1}{2}$.

Data collection: *CrysAlis CCD* (Oxford Diffraction, 2009); cell refinement: *CrysAlis RED* (Oxford Diffraction, 2009); data reduction: *CrysAlis RED*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *Mercury* (Macrae *et al.*, 2008); software used to prepare material for publication: *enCIFer* (Allen *et al.*, 2004).

Support of this investigation by the Islamic Azad University, North Tehran Branch, is gratefully acknowledged.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH5469).

References

- Allen, F. H., Johnson, O., Shields, G. P., Smith, B. R. & Towler, M. (2004). *J. Appl. Cryst.* **37**, 335–338.
- Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
- Oxford Diffraction (2009). *CrysAlis CCD* and *CrysAlis RED*. Oxford Diffraction Ltd, Yarnton, England.
- Pourayoubi, M., Nečas, M. & Negari, M. (2012). *Acta Cryst.* **C68**, o51–o56.
- Pourayoubi, M., Tarahhomi, A., Saneei, A., Rheingold, A. L. & Golen, J. A. (2011). *Acta Cryst.* **C67**, o265–o272.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Steiner, T. (2002). *Angew. Chem. Int. Ed.* **41**, 48–76.

supporting information

Acta Cryst. (2012). E68, o1813 [doi:10.1107/S160053681202154X]

***N,N'*-Bis(4-methylphenyl)-*N''*-(2,2,2-trichloroacetyl)phosphoric triamide**

Akbar Raissi Shabari, Mehrdad Pourayoubi, Hassan Fadaei, Marek Nečas and Michal Babiak

S1. Comment

The structure determination of the title compound, $\text{P}(\text{O})[\text{NHC}(\text{O})\text{CCl}_3][\text{NHC}_6\text{H}_4(4\text{-CH}_3)]_2$ (Fig. 1), was performed as a part of a project on the synthesis of new phosphoric triamides having a $\text{C}(\text{O})\text{NHP}(\text{O})$ skeleton (Pourayoubi *et al.*, 2011).

The $\text{P}=\text{O}$ (1.4727 (12) Å) and $\text{C}=\text{O}$ (1.211 (2) Å) bond lengths are standard for this category of compounds (Pourayoubi *et al.*, 2011). The P atom has a distorted tetrahedral configuration (Fig. 1). The bond angles at the P atom are in the range 102.25 (8) – 118.28 (8)°. The P—N1 and P—N2 bonds (with lengths of 1.6195 (16) Å and 1.6345 (16) Å) are shorter than the P—N3 bond (1.7071 (16) Å). As might be expected the C15—N3 bond distance (1.349 (2) Å) is shorter than the other C—N bond distances.

In the crystal, each molecule is hydrogen-bonded to two adjacent molecules through $\text{N}_{\text{C}(\text{O})\text{NHP}(\text{O})}-\text{H}\cdots\text{O}(\text{C})$ and $(\text{N}-\text{H})_2\cdots\text{O}(\text{P})$ hydrogen bonds along the *c* axis with the oxygen atom of phosphoryl group as a double-hydrogen bond acceptor (Steiner, 2002; Pourayoubi *et al.*, 2012).

S2. Experimental

$\text{CCl}_3\text{C}(\text{O})\text{NHP}(\text{O})\text{Cl}_2$ was synthesized from a reaction between phosphorus pentachloride (15.5 mmol) and 2,2,2-trichloroacetamide (15.5 mmol) in dry CCl_4 at 353 K (3 h) and then treated with formic acid 85% (15.5 mmol) at ice bath temperature.

To a solution of $\text{CCl}_3\text{C}(\text{O})\text{NHP}(\text{O})\text{Cl}_2$ (1.7 mmol) in dry chloroform (30 ml), a solution of *p*-toluidine (6.8 mmol) in the same solvent (5 ml) was added at ice bath temperature. After 4 h stirring, the solvent was removed and the product was washed with distilled water and recrystallized from methanol at room temperature. IR (KBr, cm^{-1}): 3305, 3248, 3029, 2920, 2858, 1714, 1619, 1514, 1433, 1376, 1277, 1234, 1191, 963, 882, 811, 730, 683.

Single crystals were obtained from a solution of the title compound in CH_3OH after slow evaporation at room temperature.

S3. Refinement

All carbon bound H atoms were placed at calculated positions and were refined as riding with their U_{iso} set to either $1.2U_{\text{eq}}$ or $1.5U_{\text{eq}}$ (methyl) of the respective carrier atoms; in addition, the methyl H atoms were allowed to rotate about the C—C bond. Nitrogen bound H atoms were located in a difference Fourier map and refined isotropically.




Figure 1

The molecular structure of the title compound with ellipsoids shown at the 50% probability level.




Figure 2

Partial packing view showing the formation of a chain through $N_{C(O)NHP(O)}-H \cdots O(C)$ and $(N-H)_2 \cdots O(P)$ hydrogen bonds along the c axis. The dashed lines show the donor...acceptor distances of the hydrogen bonds.

N,N'-Bis(4-methylphenyl)-*N''*-(2,2,2-trichloroacetyl)phosphoric triamide

Crystal data

$C_{16}H_{17}Cl_3N_3O_2P$

$M_r = 420.65$

Monoclinic, $P2_1/c$

Hall symbol: $-P 2_1bc$

$a = 17.5151(6) \text{ \AA}$

$b = 10.8638(4) \text{ \AA}$

$c = 9.8615(3) \text{ \AA}$

$\beta = 97.565(3)^\circ$

$V = 1860.12(11) \text{ \AA}^3$

$Z = 4$

$F(000) = 864$
 $D_x = 1.502 \text{ Mg m}^{-3}$
 Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
 Cell parameters from 5060 reflections
 $\theta = 3.3\text{--}27.7^\circ$

$\mu = 0.59 \text{ mm}^{-1}$
 $T = 120 \text{ K}$
 Prism, colourless
 $0.60 \times 0.60 \times 0.60 \text{ mm}$

Data collection

Oxford Diffraction Xcalibur Sapphire2
 diffractometer
 Radiation source: Enhance (Mo) X-ray Source
 Graphite monochromator
 Detector resolution: $8.4353 \text{ pixels mm}^{-1}$
 ω scan
 Absorption correction: multi-scan
 (CrysAlis RED; Oxford Diffraction, 2009)
 $T_{\min} = 0.955$, $T_{\max} = 1.000$

6796 measured reflections
 3265 independent reflections
 2820 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.013$
 $\theta_{\max} = 25.0^\circ$, $\theta_{\min} = 3.5^\circ$
 $h = -20 \rightarrow 8$
 $k = -12 \rightarrow 12$
 $l = -11 \rightarrow 11$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.028$
 $wR(F^2) = 0.072$
 $S = 1.04$
 3265 reflections
 240 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H atoms treated by a mixture of independent
 and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0371P)^2 + 0.8844P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.33 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.26 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	0.42010 (3)	0.60382 (4)	0.43562 (4)	0.02457 (13)
O1	0.14643 (7)	0.76151 (12)	0.35677 (12)	0.0207 (3)
P1	0.17851 (3)	0.74597 (4)	0.50145 (4)	0.01634 (12)
C1	0.12216 (10)	0.51775 (18)	0.53131 (17)	0.0191 (4)
N1	0.13973 (9)	0.63843 (15)	0.58334 (15)	0.0190 (3)
Cl2	0.46136 (3)	0.60149 (5)	0.72774 (4)	0.02712 (13)
O2	0.30612 (7)	0.68782 (12)	0.72621 (12)	0.0225 (3)
C2	0.07280 (11)	0.5003 (2)	0.41018 (18)	0.0246 (4)
H2	0.0532	0.5692	0.3573	0.030*
N2	0.17663 (9)	0.86377 (15)	0.60406 (16)	0.0187 (3)

Cl3	0.44563 (3)	0.83378 (5)	0.58356 (5)	0.02685 (13)
C3	0.05233 (11)	0.3820 (2)	0.3671 (2)	0.0287 (5)
H3	0.0200	0.3709	0.2829	0.034*
N3	0.27364 (9)	0.71385 (15)	0.49795 (15)	0.0182 (3)
C4	0.07772 (11)	0.2797 (2)	0.4435 (2)	0.0292 (5)
C5	0.12832 (12)	0.2987 (2)	0.5623 (2)	0.0335 (5)
H5	0.1480	0.2298	0.6151	0.040*
C6	0.15074 (11)	0.41629 (19)	0.6055 (2)	0.0276 (5)
H6	0.1859	0.4270	0.6865	0.033*
C7	0.22672 (10)	0.96637 (17)	0.60665 (17)	0.0180 (4)
C8	0.25836 (11)	1.00236 (18)	0.49043 (18)	0.0228 (4)
H8	0.2465	0.9582	0.4072	0.027*
C9	0.30726 (12)	1.10297 (18)	0.4972 (2)	0.0258 (4)
H9	0.3296	1.1253	0.4180	0.031*
C10	0.32484 (11)	1.17239 (18)	0.61502 (19)	0.0250 (4)
C11	0.29305 (12)	1.13365 (19)	0.7299 (2)	0.0280 (5)
H11	0.3047	1.1781	0.8130	0.034*
C12	0.24498 (11)	1.03244 (18)	0.72679 (18)	0.0237 (4)
H12	0.2244	1.0081	0.8072	0.028*
C13	0.05026 (13)	0.1515 (2)	0.4026 (3)	0.0414 (6)
H13A	0.0835	0.0908	0.4550	0.062*
H13B	0.0523	0.1396	0.3047	0.062*
H13C	-0.0029	0.1410	0.4217	0.062*
C14	0.37295 (13)	1.2873 (2)	0.6172 (2)	0.0350 (5)
H14A	0.4040	1.2957	0.7069	0.052*
H14B	0.3392	1.3590	0.5998	0.052*
H14C	0.4071	1.2819	0.5462	0.052*
C15	0.32379 (10)	0.69330 (17)	0.61173 (17)	0.0175 (4)
C16	0.40963 (10)	0.68105 (17)	0.58949 (17)	0.0192 (4)
H1N	0.1460 (12)	0.6449 (19)	0.662 (2)	0.023 (6)*
H2N	0.1638 (12)	0.848 (2)	0.672 (2)	0.026 (6)*
H3N	0.2878 (12)	0.720 (2)	0.430 (2)	0.027 (6)*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl1	0.0236 (2)	0.0341 (3)	0.0165 (2)	0.0062 (2)	0.00442 (17)	-0.00528 (19)
O1	0.0204 (6)	0.0296 (7)	0.0124 (6)	-0.0004 (6)	0.0027 (5)	0.0013 (5)
P1	0.0169 (2)	0.0216 (3)	0.0107 (2)	-0.0005 (2)	0.00235 (17)	0.00029 (18)
C1	0.0155 (9)	0.0273 (10)	0.0154 (8)	-0.0044 (8)	0.0054 (7)	-0.0026 (8)
N1	0.0243 (8)	0.0253 (9)	0.0076 (7)	-0.0037 (7)	0.0025 (6)	-0.0019 (7)
Cl2	0.0263 (3)	0.0359 (3)	0.0180 (2)	0.0092 (2)	-0.00139 (18)	0.00249 (19)
O2	0.0227 (7)	0.0329 (8)	0.0125 (6)	0.0018 (6)	0.0047 (5)	0.0003 (5)
C2	0.0218 (10)	0.0349 (12)	0.0168 (9)	-0.0005 (9)	0.0010 (7)	-0.0019 (8)
N2	0.0224 (8)	0.0230 (9)	0.0118 (7)	-0.0006 (7)	0.0064 (6)	0.0011 (7)
Cl3	0.0271 (3)	0.0285 (3)	0.0252 (2)	-0.0063 (2)	0.00431 (19)	-0.00010 (19)
C3	0.0214 (10)	0.0447 (13)	0.0199 (10)	-0.0068 (10)	0.0015 (8)	-0.0129 (9)
N3	0.0196 (8)	0.0263 (9)	0.0099 (8)	0.0011 (7)	0.0059 (6)	0.0006 (6)

C4	0.0211 (10)	0.0318 (12)	0.0350 (11)	-0.0031 (9)	0.0050 (8)	-0.0110 (9)
C5	0.0296 (11)	0.0267 (11)	0.0411 (12)	-0.0014 (10)	-0.0065 (9)	-0.0007 (10)
C6	0.0271 (11)	0.0277 (11)	0.0254 (10)	-0.0017 (9)	-0.0062 (8)	-0.0013 (8)
C7	0.0173 (9)	0.0186 (9)	0.0177 (9)	0.0027 (8)	0.0011 (7)	0.0023 (7)
C8	0.0298 (10)	0.0225 (10)	0.0164 (9)	0.0016 (9)	0.0039 (7)	0.0014 (8)
C9	0.0287 (10)	0.0237 (11)	0.0262 (10)	0.0006 (9)	0.0080 (8)	0.0065 (8)
C10	0.0216 (10)	0.0229 (10)	0.0292 (10)	0.0010 (8)	-0.0018 (8)	0.0059 (8)
C11	0.0352 (11)	0.0249 (11)	0.0220 (10)	-0.0029 (9)	-0.0033 (8)	-0.0010 (8)
C12	0.0289 (10)	0.0259 (10)	0.0160 (9)	-0.0006 (9)	0.0025 (7)	0.0020 (8)
C13	0.0329 (12)	0.0363 (13)	0.0537 (15)	-0.0078 (11)	0.0003 (11)	-0.0157 (11)
C14	0.0323 (12)	0.0326 (12)	0.0386 (12)	-0.0074 (10)	-0.0010 (9)	0.0042 (10)
C15	0.0205 (9)	0.0174 (9)	0.0146 (9)	0.0002 (8)	0.0026 (7)	-0.0012 (7)
C16	0.0206 (9)	0.0226 (10)	0.0144 (9)	0.0007 (8)	0.0024 (7)	-0.0014 (7)

Geometric parameters (Å, °)

C11—C16	1.7644 (18)	C5—C6	1.387 (3)
O1—P1	1.4727 (12)	C5—H5	0.9500
P1—N1	1.6195 (16)	C6—H6	0.9500
P1—N2	1.6345 (16)	C7—C12	1.386 (3)
P1—N3	1.7071 (16)	C7—C8	1.393 (2)
C1—C6	1.380 (3)	C8—C9	1.385 (3)
C1—C2	1.392 (2)	C8—H8	0.9500
C1—N1	1.427 (2)	C9—C10	1.385 (3)
N1—H1N	0.77 (2)	C9—H9	0.9500
C12—C16	1.7611 (18)	C10—C11	1.392 (3)
O2—C15	1.211 (2)	C10—C14	1.505 (3)
C2—C3	1.386 (3)	C11—C12	1.383 (3)
C2—H2	0.9500	C11—H11	0.9500
N2—C7	1.417 (2)	C12—H12	0.9500
N2—H2N	0.76 (2)	C13—H13A	0.9800
C13—C16	1.7786 (19)	C13—H13B	0.9800
C3—C4	1.383 (3)	C13—H13C	0.9800
C3—H3	0.9500	C14—H14A	0.9800
N3—C15	1.349 (2)	C14—H14B	0.9800
N3—H3N	0.75 (2)	C14—H14C	0.9800
C4—C5	1.388 (3)	C15—C16	1.553 (2)
C4—C13	1.510 (3)		
O1—P1—N1	115.74 (8)	C9—C8—C7	119.49 (18)
O1—P1—N2	118.28 (8)	C9—C8—H8	120.3
N1—P1—N2	102.25 (8)	C7—C8—H8	120.3
O1—P1—N3	104.64 (7)	C8—C9—C10	122.55 (18)
N1—P1—N3	109.71 (8)	C8—C9—H9	118.7
N2—P1—N3	105.77 (8)	C10—C9—H9	118.7
C6—C1—C2	119.16 (18)	C9—C10—C11	116.71 (18)
C6—C1—N1	119.83 (16)	C9—C10—C14	121.72 (18)
C2—C1—N1	120.91 (17)	C11—C10—C14	121.51 (18)

C1—N1—P1	124.64 (12)	C12—C11—C10	122.02 (18)
C1—N1—H1N	116.1 (16)	C12—C11—H11	119.0
P1—N1—H1N	114.9 (16)	C10—C11—H11	119.0
C3—C2—C1	119.68 (19)	C11—C12—C7	120.13 (17)
C3—C2—H2	120.2	C11—C12—H12	119.9
C1—C2—H2	120.2	C7—C12—H12	119.9
C7—N2—P1	124.35 (12)	C4—C13—H13A	109.5
C7—N2—H2N	114.8 (17)	C4—C13—H13B	109.5
P1—N2—H2N	113.8 (17)	H13A—C13—H13B	109.5
C4—C3—C2	121.81 (18)	C4—C13—H13C	109.5
C4—C3—H3	119.1	H13A—C13—H13C	109.5
C2—C3—H3	119.1	H13B—C13—H13C	109.5
C15—N3—P1	123.18 (13)	C10—C14—H14A	109.5
C15—N3—H3N	120.2 (17)	C10—C14—H14B	109.5
P1—N3—H3N	116.2 (17)	H14A—C14—H14B	109.5
C3—C4—C5	117.62 (19)	C10—C14—H14C	109.5
C3—C4—C13	121.82 (19)	H14A—C14—H14C	109.5
C5—C4—C13	120.5 (2)	H14B—C14—H14C	109.5
C6—C5—C4	121.3 (2)	O2—C15—N3	124.41 (16)
C6—C5—H5	119.3	O2—C15—C16	119.89 (15)
C4—C5—H5	119.3	N3—C15—C16	115.66 (14)
C1—C6—C5	120.31 (18)	C15—C16—C12	109.98 (12)
C1—C6—H6	119.8	C15—C16—C11	111.97 (12)
C5—C6—H6	119.8	C12—C16—C11	109.34 (10)
C12—C7—C8	119.06 (17)	C15—C16—C13	106.17 (12)
C12—C7—N2	119.70 (16)	C12—C16—C13	109.57 (10)
C8—C7—N2	121.24 (16)	C11—C16—C13	109.75 (9)
C6—C1—N1—P1	125.17 (17)	P1—N2—C7—C12	-152.11 (15)
C2—C1—N1—P1	-58.6 (2)	P1—N2—C7—C8	27.7 (2)
O1—P1—N1—C1	44.72 (17)	C12—C7—C8—C9	0.0 (3)
N2—P1—N1—C1	174.73 (14)	N2—C7—C8—C9	-179.82 (17)
N3—P1—N1—C1	-73.35 (16)	C7—C8—C9—C10	-1.7 (3)
C6—C1—C2—C3	0.9 (3)	C8—C9—C10—C11	2.3 (3)
N1—C1—C2—C3	-175.35 (16)	C8—C9—C10—C14	-174.99 (19)
O1—P1—N2—C7	-75.36 (16)	C9—C10—C11—C12	-1.3 (3)
N1—P1—N2—C7	156.22 (14)	C14—C10—C11—C12	176.03 (19)
N3—P1—N2—C7	41.39 (16)	C10—C11—C12—C7	-0.3 (3)
C1—C2—C3—C4	2.0 (3)	C8—C7—C12—C11	1.0 (3)
O1—P1—N3—C15	179.94 (15)	N2—C7—C12—C11	-179.19 (17)
N1—P1—N3—C15	-55.29 (17)	P1—N3—C15—O2	4.8 (3)
N2—P1—N3—C15	54.30 (17)	P1—N3—C15—C16	-172.98 (13)
C2—C3—C4—C5	-3.5 (3)	O2—C15—C16—C12	23.4 (2)
C2—C3—C4—C13	174.88 (19)	N3—C15—C16—C12	-158.68 (14)
C3—C4—C5—C6	2.1 (3)	O2—C15—C16—C11	145.21 (15)
C13—C4—C5—C6	-176.3 (2)	N3—C15—C16—C11	-36.9 (2)
C2—C1—C6—C5	-2.3 (3)	O2—C15—C16—C13	-95.03 (18)
N1—C1—C6—C5	174.01 (18)	N3—C15—C16—C13	82.86 (17)

C4—C5—C6—C1 0.8 (3)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N1—H1N \cdots O1 ⁱ	0.77 (2)	2.17 (2)	2.8953 (19)	156 (2)
N2—H2N \cdots O1 ⁱ	0.76 (2)	2.23 (2)	2.948 (2)	159 (2)
N3—H3N \cdots O2 ⁱⁱ	0.75 (2)	2.31 (2)	3.008 (2)	157 (2)

Symmetry codes: (i) $x, -y+3/2, z+1/2$; (ii) $x, -y+3/2, z-1/2$.