

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

4-Cyanopyridinium bromide

Wen-Ni Zheng

College of Chemistry and Chemical Engineering, Southeast University, Nanjing 210096, People's Republic of China

Correspondence e-mail: chenxinyuanseu@yahoo.com.cn


Received 27 April 2012; accepted 16 May 2012

 Key indicators: single-crystal X-ray study; $T = 173$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.032; wR factor = 0.066; data-to-parameter ratio = 19.3.

In the title compound, $\text{C}_6\text{H}_5\text{N}_2^+\cdot\text{Br}^-$, the pyridine N atom is protonated and involved in an intermolecular $\text{N}-\text{H}\cdots\text{Br}$ hydrogen bond which, together with weak $\text{C}-\text{H}\cdots\text{N}$ hydrogen bonds, results in the formation of a chain along the c axis. Weak intermolecular $\text{C}-\text{H}\cdots\text{Br}$ interactions between pyridine H atoms and Br^- anions connect these chains into a network parallel to the bc plane.

Related literature

For the structures and properties of related compounds see: Fu *et al.* (2011a,b); Dai & Chen (2011).



Experimental

Crystal data

 $\text{C}_6\text{H}_5\text{N}_2^+\cdot\text{Br}^-$
 $M_r = 185.03$

 Monoclinic, $P2_1/n$
 $a = 7.3918$ (5) Å

 $b = 12.2587$ (4) Å

 $c = 8.1671$ (3) Å

 $\beta = 111.720$ (1)°

 $V = 687.51$ (6) Å³
 $Z = 4$

 Mo $K\alpha$ radiation

 $\mu = 5.88$ mm⁻¹
 $T = 173$ K

 $0.10 \times 0.05 \times 0.05$ mm

Data collection

Rigaku Mercury2 diffractometer

Absorption correction: multi-scan

 (*CrystalClear*; Rigaku, 2005)

 $T_{\min} = 0.910$, $T_{\max} = 1.000$

4771 measured reflections

1579 independent reflections

 1296 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.047$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.032$
 $wR(F^2) = 0.066$
 $S = 0.95$

1579 reflections

82 parameters

H-atom parameters constrained

 $\Delta\rho_{\max} = 0.57$ e Å⁻³
 $\Delta\rho_{\min} = -0.92$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1}\cdots\text{Br1}^{\text{i}}$	0.90	2.26	3.133 (2)	164
$\text{C1}-\text{H1A}\cdots\text{Br1}$	0.93	2.88	3.615 (2)	137
$\text{C2}-\text{H2A}\cdots\text{Br1}^{\text{ii}}$	0.93	2.77	3.645 (2)	156
$\text{C5}-\text{H5A}\cdots\text{N2}^{\text{iii}}$	0.93	2.66	3.435 (4)	142

 Symmetry codes: (i) $-x + 1, -y + 1, -z$; (ii) $-x + 1, -y + 1, -z + 1$; (iii) $x, y, z - 1$.

Data collection: *CrystalClear* (Rigaku, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

This work was supported by a start-up grant from Southeast University, China.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: MW2067).

References

- Dai, J. & Chen, X.-Y. (2011). *Acta Cryst.* **E67**, o287.
 Fu, D.-W., Zhang, W., Cai, H.-L., Ge, J.-Z., Zhang, Y. & Xiong, R.-G. (2011b). *Adv. Mater.* **23**, 5658–5662.
 Fu, D.-W., Zhang, W., Cai, H.-L., Zhang, Y., Ge, J.-Z., Xiong, R.-G. & Huang, S. P. D. (2011a). *J. Am. Chem. Soc.* **133**, 12780–12786.
 Rigaku (2005). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supporting information

Acta Cryst. (2012). E68, o1825 [doi:10.1107/S160053681202209X]

4-Cyanopyridinium bromide

Wen-Ni Zheng

S1. Comment

Simple organic salts containing strong intermolecular H-bonds have attracted attention as materials which display ferroelectric-paraelectric phase transitions (Fu *et al.*, 2011*a*; Fu *et al.*, 2011*b*). With the purpose of obtaining crystals of organic salts which might undergo such phase transitions, various organic molecules have been studied and a series of new materials have been elaborated (Dai & Chen 2011). Herewith we present the synthesis and crystal structure of the title compound, 4-cyanopyridinium bromide.

In the title compound (Fig. 1), the bond lengths and angles have normal values. The asymmetric unit is composed of one 4-cyanopyridinium cation and one Br⁻ anion. The protonated N atom is involved in a strong N—H \cdots Br hydrogen bond (Table 1) which accompanying the C5—H5A \cdots N2 H-bond generates a linear chain parallel to *c*-axis while weak C1—H1A \cdots Br and C2—H2A \cdots Br1 interactions serve to link the chains into a 3-dimensional layer structure (Fig. 2 and Table 1).

S2. Experimental

Isonicotinonitrile (20 mmol), aqueous HBr (5 mL, 2 mol/L) and ethanol (50 mL) were added to a 100mL flask. The mixture was stirred at 60° C for 2 h, and then the precipitate was filtrated out. Colourless crystals suitable for X-ray diffraction were obtained by slow evaporation of the filtrate.

S3. Refinement

All H atoms attached to C atoms were situated into the idealized positions and treated as riding with C—H = 0.93 Å (aromatic) with $U_{iso}(H)=1.2U_{eq}(C)$. The positional parameters of the H atom (N) were refined freely. And in the last stage of the refinement, they were restrained with the H—N = 0.90 (2)Å, with $U_{iso}(H)=1.2U_{eq}(N)$.




Figure 1

A view of the asymmetric unit with the atomic numbering scheme. The displacement ellipsoids were drawn at the 30% probability level.




Figure 2

The crystal packing of the title compound viewed along the *a* axis showing the N—H···Br, C—H···N and C—H···Br interactions (dotted line) in the title compound.

4-Cyanopyridinium bromide

Crystal data

$C_6H_5N_2^+ \cdot Br^-$

$M_r = 185.03$

Monoclinic, $P2_1/n$

Hall symbol: $-P\ 2_1n$

$a = 7.3918\ (5)\ \text{\AA}$

$b = 12.2587\ (4)\ \text{\AA}$

$c = 8.1671\ (3)\ \text{\AA}$

$\beta = 111.720\ (1)^\circ$

$V = 687.51\ (6)\ \text{\AA}^3$

$Z = 4$

$F(000) = 360$

$D_x = 1.788\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 1579 reflections

$\theta = 3.2\text{--}27.5^\circ$

$\mu = 5.88\ \text{mm}^{-1}$

$T = 173\ \text{K}$

Block, colorless

$0.10 \times 0.05 \times 0.05\ \text{mm}$

Data collection

Rigaku Mercury2
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

Detector resolution: $13.6612\ \text{pixels mm}^{-1}$

ω & φ scans

Absorption correction: multi-scan

(*CrystalClear*; Rigaku, 2005)

$T_{\min} = 0.910$, $T_{\max} = 1.000$

4771 measured reflections

1579 independent reflections

1296 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.047$

$\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 3.2^\circ$

$h = -9 \rightarrow 9$

$k = -15 \rightarrow 15$

$l = -10 \rightarrow 10$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.032$

$wR(F^2) = 0.066$

$S = 0.95$

1579 reflections

82 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0255P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.57 \text{ e } \text{Å}^{-3}$
 $\Delta\rho_{\min} = -0.92 \text{ e } \text{Å}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.61085 (4)	0.41149 (2)	0.32517 (4)	0.01972 (12)
N1	0.4087 (4)	0.74050 (19)	-0.0107 (3)	0.0178 (6)
H1	0.4160	0.7079	-0.1068	0.021*
C1	0.4249 (4)	0.6702 (3)	0.1191 (4)	0.0218 (7)
H1A	0.4410	0.5962	0.1038	0.026*
C2	0.4176 (5)	0.7076 (2)	0.2761 (4)	0.0206 (7)
H2A	0.4295	0.6597	0.3677	0.025*
N2	0.3889 (4)	0.8933 (2)	0.5892 (4)	0.0318 (7)
C3	0.3922 (4)	0.8184 (2)	0.2937 (4)	0.0164 (7)
C4	0.3754 (4)	0.8896 (2)	0.1569 (4)	0.0198 (7)
H4A	0.3575	0.9640	0.1682	0.024*
C5	0.3857 (4)	0.8479 (2)	0.0032 (4)	0.0193 (7)
H5A	0.3767	0.8941	-0.0898	0.023*
C6	0.3875 (5)	0.8612 (3)	0.4580 (4)	0.0202 (7)

Atomic displacement parameters (Å^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.0246 (2)	0.01924 (19)	0.01933 (18)	-0.00107 (15)	0.01284 (14)	-0.00215 (13)
N1	0.0149 (14)	0.0239 (14)	0.0174 (12)	0.0034 (11)	0.0094 (11)	-0.0039 (11)
C1	0.024 (2)	0.0189 (17)	0.0243 (17)	0.0035 (14)	0.0111 (15)	-0.0006 (14)
C2	0.0201 (18)	0.0237 (17)	0.0206 (16)	0.0045 (14)	0.0106 (14)	0.0037 (14)
N2	0.045 (2)	0.0311 (17)	0.0251 (15)	-0.0001 (15)	0.0200 (14)	-0.0045 (13)
C3	0.0082 (16)	0.0260 (17)	0.0163 (15)	0.0014 (13)	0.0060 (13)	0.0006 (13)
C4	0.0194 (18)	0.0212 (17)	0.0221 (16)	0.0005 (14)	0.0118 (14)	-0.0032 (13)
C5	0.0166 (18)	0.0226 (17)	0.0207 (16)	0.0007 (14)	0.0094 (14)	0.0034 (14)
C6	0.0184 (18)	0.0248 (17)	0.0194 (16)	-0.0007 (14)	0.0093 (14)	0.0022 (14)

Geometric parameters (Å , $^\circ$)

N1—C1	1.337 (4)	N2—C6	1.138 (4)
N1—C5	1.337 (4)	C3—C4	1.387 (4)

N1—H1	0.9003	C3—C6	1.453 (4)
C1—C2	1.381 (4)	C4—C5	1.384 (4)
C1—H1A	0.9300	C4—H4A	0.9300
C2—C3	1.385 (4)	C5—H5A	0.9300
C2—H2A	0.9300		
C1—N1—C5	122.9 (2)	C2—C3—C6	120.0 (3)
C1—N1—H1	112.9	C4—C3—C6	119.4 (3)
C5—N1—H1	124.2	C5—C4—C3	118.6 (3)
N1—C1—C2	120.0 (3)	C5—C4—H4A	120.7
N1—C1—H1A	120.0	C3—C4—H4A	120.7
C2—C1—H1A	120.0	N1—C5—C4	119.5 (3)
C1—C2—C3	118.4 (3)	N1—C5—H5A	120.2
C1—C2—H2A	120.8	C4—C5—H5A	120.2
C3—C2—H2A	120.8	N2—C6—C3	178.0 (4)
C2—C3—C4	120.6 (3)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N1—H1 \cdots Br1 ⁱ	0.90	2.26	3.133 (2)	164
C1—H1A \cdots Br1	0.93	2.88	3.615 (2)	137
C2—H2A \cdots Br1 ⁱⁱ	0.93	2.77	3.645 (2)	156
C5—H5A \cdots N2 ⁱⁱⁱ	0.93	2.66	3.435 (4)	142

Symmetry codes: (i) $-x+1, -y+1, -z$; (ii) $-x+1, -y+1, -z+1$; (iii) $x, y, z-1$.