

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

# 1-(3,3-Dichloroallyloxy)-4-methyl-2-nitrobenzene

Dong-mei Ren

 Security and Environment Engineering College, Capital University of Economics and Business, Beijing 10070, People's Republic of China  
 Correspondence e-mail: nanoren@126.com


Received 2 May 2012; accepted 20 May 2012

 Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(\text{C}-\text{C}) = 0.004$  Å;  $R$  factor = 0.048;  $wR$  factor = 0.156; data-to-parameter ratio = 14.5.

In the title compound,  $\text{C}_{10}\text{H}_9\text{Cl}_2\text{NO}_3$ , the dihedral angle between the benzene ring and the plane of the nitro group is  $39.1(1)^\circ$ , while that between the benzene ring and the plane through the three C and two Cl atoms of the dichloroallyloxy unit is  $40.1(1)^\circ$ . In the crystal,  $\text{C}-\text{H}\cdots\text{O}$  hydrogen bonds to the nitro groups form chains along the  $b$  axis. These chains are linked by inversion-related pairs of  $\text{Cl}\cdots\text{O}$  interactions at a distance of  $3.060(3)$  Å, forming sheets approximately parallel to  $[\bar{2}01]$  and generating  $R_2^2(18)$  rings.  $\pi-\pi$  contacts between benzene rings in adjacent sheets, with centroid-centroid distances of  $3.671(2)$  Å, stack molecules along  $c$ .

## Related literature

For background to the applications of the title compound, see: Kolosov *et al.* (2002). For its synthesis, see: Walker *et al.* (2005). For bond-length data, see: Allen *et al.* (1987). For hydrogen-bond motifs, see: Bernstein *et al.* (1995).



## Experimental

### Crystal data

 $\text{C}_{10}\text{H}_9\text{Cl}_2\text{NO}_3$ 
 $M_r = 262.08$ 

 Triclinic,  $P\bar{1}$   
 $a = 7.5430(15)$  Å  
 $b = 7.7630(16)$  Å  
 $c = 10.713(2)$  Å  
 $\alpha = 83.69(3)^\circ$   
 $\beta = 88.78(3)^\circ$   
 $\gamma = 67.23(3)^\circ$ 
 $V = 574.8(2)$  Å<sup>3</sup>  
 $Z = 2$   
 Mo  $K\alpha$  radiation  
 $\mu = 0.56$  mm<sup>-1</sup>  
 $T = 293$  K  
 $0.30 \times 0.20 \times 0.10$  mm

### Data collection

 Enraf-Nonius CAD-4 diffractometer  
 Absorption correction:  $\psi$  scan (North *et al.*, 1968)  
 $T_{\min} = 0.851$ ,  $T_{\max} = 0.947$   
 2277 measured reflections

 2103 independent reflections  
 1638 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.021$   
 3 standard reflections every 200 reflections  
 intensity decay: 1%

### Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.048$   
 $wR(F^2) = 0.156$   
 $S = 1.01$   
 2103 reflections

 145 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.28$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.28$  e Å<sup>-3</sup>
**Table 1**

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C1}-\text{H1A}\cdots\text{O2}^i$	0.93	2.59	3.241 (4)	127

 Symmetry code: (i)  $x, y + 1, z$ .

Data collection: *CAD-4 Software* (Enraf-Nonius, 1985); cell refinement: *CAD-4 Software*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

This study was supported financially by the Capital University of Economics and Business (00891162721716) and by the Scientific Research Level Project of the Beijing Education Commission Foundation. The author thanks the Center of Testing and Analysis, Nanjing University, for the data collection.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SJ5235).

## References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Bernstein, J., Davis, R. E., Shimon, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Enraf-Nonius (1985). *CAD-4 Software*. Enraf-Nonius, Delft, The Netherlands.
- Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.
- Kolosov, S., Adamovich, V., Djurovich, P., Thompson, M. E. & Adachi, C. (2002). *J. Am. Chem. Soc.* **124**, 9945–9954.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Walker, E. R., Leung, S. Y. & Barrett, A. G. M. (2005). *Tetrahedron Lett.* **46**, 6537–6540.

## supporting information

*Acta Cryst.* (2012). E68, o1929 [doi:10.1107/S1600536812023057]

## 1-(3,3-Dichloroallyloxy)-4-methyl-2-nitrobenzene

Dong-mei Ren

### S1. Comment

The title compound is an important intermediate in the synthesis of phenanthrenes, which can be utilized to synthesize organic semiconductors and conjugated polymers (Walker *et al.*, 2005). These materials are of wide current interest with applications in electronic and optoelectronic devices including light-emitting diodes (Kolosov *et al.*, 2002). We report here the crystal structure of the title compound, (I), as we have interests in this field.

The molecular structure of (I) is shown in Fig. 1. The dihedral angle between the (C1...C6) rings and the (C8/C9/C10/C11/C12/H9A) segment of the allyloxy substituent is 40.1 (1)° with the nitro group (N/O1/O2) inclined at 39.1 (1)° to the ring plane. Bond distances in the molecule are normal (Allen *et al.* 1987).


In the crystal structure there is an intermolecular C1—H1A...O2 hydrogen bond that links molecules into chains along the *b* axis (Table 1, Fig. 2). Short C11...O1<sup>i</sup> (<sup>i</sup> = 1-x, -y, 1-z) contacts at a distance of 3.060 (3) Å form inversion dimers and generate R<sup>2</sup><sub>2</sub>(18) rings (Bernstein *et al.*, 1995). These contacts link the hydrogen bonded chains into sheets approximately parallel to [-2 0 1]. Additional  $\pi$ ... $\pi$  contacts with centroid to centroid distances 3.671 (2) Å between benzene rings in adjacent sheets, stack molecules along *c* and generate a three dimensional network structure.

### S2. Experimental

The title compound, (I) was prepared by a method reported in literature (Walker *et al.*, 2005). The crystals were obtained by dissolving (I) (0.1 g) in methanol (30 ml) and evaporating the solvent slowly at room temperature for about 8 d.


### S3. Refinement

All H atoms were positioned geometrically and constrained to ride on their parent atoms, with C—H = 0.93 Å for aromatic H and 0.96 Å for alkyl H, respectively. The  $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C})$ , where  $x = 1.2$  for aromatic H and  $x = 1.5$  for other H.



**Figure 1**

The molecular structure of (I), with the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level.



**Figure 2**

A packing diagram of (I).

### 1-(3,3-Dichloroallyloxy)-4-methyl-2-nitrobenzene

#### Crystal data

$C_{10}H_9Cl_2NO_3$

$M_r = 262.08$

Triclinic,  $P\bar{1}$

Hall symbol:  $-P\ 1$

$a = 7.5430$  (15) Å

$b = 7.7630$  (16) Å

$c = 10.713$  (2) Å

$\alpha = 83.69$  (3)°

$\beta = 88.78$  (3)°

$\gamma = 67.23$  (3)°

$V = 574.8$  (2) Å<sup>3</sup>

$Z = 2$

$F(000) = 268$

$D_x = 1.514$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$   
 Cell parameters from 25 reflections  
 $\theta = 10\text{--}13^\circ$   
 $\mu = 0.56 \text{ mm}^{-1}$

$T = 293 \text{ K}$   
 Block, colourless  
 $0.30 \times 0.20 \times 0.10 \text{ mm}$

*Data collection*

Enraf–Nonius CAD-4  
 diffractometer  
 Radiation source: fine-focus sealed tube  
 Graphite monochromator  
 $\omega/2\theta$  scans  
 Absorption correction:  $\psi$  scan  
 (North *et al.*, 1968)  
 $T_{\min} = 0.851$ ,  $T_{\max} = 0.947$   
 2277 measured reflections

2103 independent reflections  
 1638 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.021$   
 $\theta_{\max} = 25.4^\circ$ ,  $\theta_{\min} = 1.9^\circ$   
 $h = 0 \rightarrow 9$   
 $k = -8 \rightarrow 9$   
 $l = -12 \rightarrow 12$   
 3 standard reflections every 200 reflections  
 intensity decay: 1%

*Refinement*

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.048$   
 $wR(F^2) = 0.156$   
 $S = 1.01$   
 2103 reflections  
 145 parameters  
 0 restraints  
 Primary atom site location: structure-invariant  
 direct methods

Secondary atom site location: difference Fourier  
 map  
 Hydrogen site location: inferred from  
 neighbouring sites  
 H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.1P)^2 + 0.130P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} < 0.001$   
 $\Delta\rho_{\max} = 0.28 \text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.28 \text{ e \AA}^{-3}$

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	0.35049 (14)	0.31042 (13)	0.40617 (7)	0.0723 (3)
Cl2	0.20254 (13)	0.53282 (12)	0.60994 (8)	0.0713 (3)
O1	0.8637 (4)	−0.2999 (3)	0.8304 (2)	0.0767 (8)
O2	0.7766 (4)	−0.4263 (3)	0.9976 (2)	0.0731 (7)
O3	0.6554 (3)	0.0677 (3)	0.82149 (17)	0.0542 (5)
N	0.8110 (4)	−0.2976 (3)	0.9385 (2)	0.0534 (6)
C1	0.6985 (4)	0.1898 (4)	1.0130 (3)	0.0506 (7)
H1A	0.6478	0.3133	0.9755	0.061*
C2	0.7574 (4)	0.1533 (4)	1.1372 (3)	0.0511 (7)
H2A	0.7446	0.2537	1.1816	0.061*
C3	0.8352 (4)	−0.0276 (4)	1.1987 (2)	0.0461 (6)
C4	0.8481 (4)	−0.1727 (4)	1.1298 (2)	0.0457 (6)

H4A	0.8972	-0.2958	1.1681	0.055*
C5	0.7892 (4)	-0.1372 (4)	1.0055 (2)	0.0411 (6)
C6	0.7139 (4)	0.0451 (4)	0.9434 (2)	0.0417 (6)
C7	0.9018 (5)	-0.0623 (5)	1.3344 (3)	0.0661 (9)
H7A	0.8808	0.0554	1.3652	0.099*
H7B	1.0364	-0.1409	1.3409	0.099*
H7C	0.8306	-0.1235	1.3835	0.099*
C8	0.5768 (5)	0.2544 (4)	0.7592 (3)	0.0515 (7)
H8A	0.6681	0.3142	0.7612	0.062*
H8B	0.4597	0.3296	0.7989	0.062*
C9	0.5358 (5)	0.2340 (4)	0.6279 (3)	0.0537 (7)
H9A	0.6257	0.1337	0.5915	0.064*
C10	0.3856 (5)	0.3433 (4)	0.5586 (3)	0.0517 (7)

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C11	0.0946 (7)	0.0785 (6)	0.0430 (4)	-0.0320 (5)	-0.0101 (4)	-0.0062 (4)
C12	0.0730 (6)	0.0587 (5)	0.0630 (5)	-0.0040 (4)	-0.0091 (4)	-0.0066 (4)
O1	0.112 (2)	0.0556 (14)	0.0473 (13)	-0.0128 (13)	-0.0062 (13)	-0.0175 (10)
O2	0.0858 (17)	0.0430 (12)	0.0966 (18)	-0.0307 (12)	0.0023 (14)	-0.0107 (12)
O3	0.0765 (14)	0.0393 (10)	0.0438 (11)	-0.0199 (10)	-0.0116 (10)	0.0006 (8)
N	0.0584 (15)	0.0364 (12)	0.0599 (16)	-0.0107 (11)	-0.0099 (12)	-0.0090 (11)
C1	0.0641 (18)	0.0350 (14)	0.0531 (16)	-0.0199 (13)	-0.0054 (13)	-0.0024 (12)
C2	0.0625 (18)	0.0450 (15)	0.0504 (16)	-0.0239 (13)	0.0007 (13)	-0.0127 (12)
C3	0.0460 (15)	0.0525 (16)	0.0410 (14)	-0.0199 (12)	0.0037 (11)	-0.0078 (12)
C4	0.0489 (15)	0.0438 (14)	0.0412 (14)	-0.0166 (12)	0.0006 (11)	0.0027 (11)
C5	0.0426 (14)	0.0373 (13)	0.0453 (14)	-0.0170 (11)	0.0030 (11)	-0.0065 (11)
C6	0.0453 (14)	0.0415 (14)	0.0376 (13)	-0.0165 (11)	-0.0009 (11)	-0.0020 (11)
C7	0.076 (2)	0.076 (2)	0.0444 (16)	-0.0263 (18)	-0.0025 (15)	-0.0086 (15)
C8	0.0635 (18)	0.0411 (14)	0.0446 (15)	-0.0158 (13)	-0.0059 (13)	0.0023 (12)
C9	0.0617 (18)	0.0477 (16)	0.0449 (15)	-0.0139 (14)	0.0011 (13)	-0.0051 (12)
C10	0.0648 (18)	0.0487 (16)	0.0413 (15)	-0.0218 (14)	-0.0014 (13)	-0.0033 (12)

*Geometric parameters (Å, °)*

C11—C10	1.721 (3)	C3—C7	1.510 (4)
C12—C10	1.717 (3)	C4—C5	1.378 (4)
O1—N	1.216 (3)	C4—H4A	0.9300
O2—N	1.233 (3)	C5—C6	1.398 (4)
O3—C6	1.358 (3)	C7—H7A	0.9600
O3—C8	1.427 (3)	C7—H7B	0.9600
N—C5	1.458 (3)	C7—H7C	0.9600
C1—C2	1.375 (4)	C8—C9	1.484 (4)
C1—C6	1.383 (4)	C8—H8A	0.9700
C1—H1A	0.9300	C8—H8B	0.9700
C2—C3	1.388 (4)	C9—C10	1.307 (4)
C2—H2A	0.9300	C9—H9A	0.9300

C3—C4	1.384 (4)		
C6—O3—C8	117.7 (2)	O3—C6—C5	118.1 (2)
O1—N—O2	123.7 (2)	C1—C6—C5	116.9 (2)
O1—N—C5	119.6 (3)	C3—C7—H7A	109.5
O2—N—C5	116.6 (2)	C3—C7—H7B	109.5
C2—C1—C6	120.8 (3)	H7A—C7—H7B	109.5
C2—C1—H1A	119.6	C3—C7—H7C	109.5
C6—C1—H1A	119.6	H7A—C7—H7C	109.5
C1—C2—C3	122.5 (3)	H7B—C7—H7C	109.5
C1—C2—H2A	118.7	O3—C8—C9	105.5 (2)
C3—C2—H2A	118.7	O3—C8—H8A	110.6
C4—C3—C2	116.9 (2)	C9—C8—H8A	110.6
C4—C3—C7	122.2 (3)	O3—C8—H8B	110.6
C2—C3—C7	120.9 (3)	C9—C8—H8B	110.6
C5—C4—C3	120.9 (3)	H8A—C8—H8B	108.8
C5—C4—H4A	119.5	C10—C9—C8	126.3 (3)
C3—C4—H4A	119.5	C10—C9—H9A	116.9
C4—C5—C6	122.0 (3)	C8—C9—H9A	116.9
C4—C5—N	117.7 (2)	C9—C10—C12	123.7 (2)
C6—C5—N	120.3 (2)	C9—C10—C11	123.2 (2)
O3—C6—C1	124.9 (2)	C12—C10—C11	113.06 (18)
C6—C1—C2—C3	0.3 (5)	C8—O3—C6—C5	179.3 (2)
C1—C2—C3—C4	-1.3 (4)	C2—C1—C6—O3	178.3 (3)
C1—C2—C3—C7	178.8 (3)	C2—C1—C6—C5	0.7 (4)
C2—C3—C4—C5	1.2 (4)	C4—C5—C6—O3	-178.6 (2)
C7—C3—C4—C5	-178.9 (3)	N—C5—C6—O3	2.5 (4)
C3—C4—C5—C6	-0.2 (4)	C4—C5—C6—C1	-0.8 (4)
C3—C4—C5—N	178.7 (3)	N—C5—C6—C1	-179.7 (3)
O1—N—C5—C4	-140.1 (3)	C6—O3—C8—C9	177.3 (2)
O2—N—C5—C4	38.8 (4)	O3—C8—C9—C10	140.5 (3)
O1—N—C5—C6	38.9 (4)	C8—C9—C10—C12	-1.1 (5)
O2—N—C5—C6	-142.3 (3)	C8—C9—C10—C11	179.4 (2)
C8—O3—C6—C1	1.7 (4)		

Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C1—H1A $\cdots$ O2 <sup>i</sup>	0.93	2.59	3.241 (4)	127

Symmetry code: (i)  $x, y+1, z$ .