

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Di- μ -aqua-bis{diaqua[μ -4-({4-[bis(2-hydroxyethyl)amino]-6-chloro-1,3,5-triazin-2-yl]amino)benzenesulfonato]-sodium(I)}

Mei Zhang, Fu-Yu Sun, Gui-Zhe Zhao and Ya-Qing Liu*

Research Center for Engineering Technology of Polymeric Composites of Shanxi Province, College of Materials Science and Engineering, North University of China, Taiyuan 030051, People's Republic of China

Correspondence e-mail: gczx2012@gmail.com


Received 11 May 2012; accepted 24 May 2012

 Key indicators: single-crystal X-ray study; $T = 113$ K; mean $\sigma(\text{C}-\text{C}) = 0.006$ Å; R factor = 0.062; wR factor = 0.131; data-to-parameter ratio = 12.1.

In the dinuclear title compound, $[\text{Na}_2(\text{C}_{13}\text{H}_{15}\text{ClN}_5\text{O}_5\text{S})_2(\text{H}_2\text{O})_6]_n$, two Na^+ cations, disposed about a centre of inversion, are linked by two bridging water molecules. The coordination geometry is based on an O_5 donor set defined by four water molecules and a 4-aminobenzenesulfonate O atom in a distorted trigonal-bipyramidal geometry. In the crystal, significant $\text{O}-\text{H}\cdots\text{O}$, $\text{O}-\text{H}\cdots\text{N}$ and $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds lead to the formation of a three-dimensional architecture.

Related literature

For commercial and synthetic applications of related compounds, see: Candiani & Frigerio (2007); Hollink *et al.* (2005); Konstantion & Petrova (2002).



Experimental

Crystal data

$[\text{Na}_2(\text{C}_{13}\text{H}_{15}\text{ClN}_5\text{O}_5\text{S})_2(\text{H}_2\text{O})_6]$
 $M_r = 931.72$
 Triclinic, $P\bar{1}$
 $a = 7.5628$ (7) Å
 $b = 8.6274$ (8) Å
 $c = 15.532$ (2) Å
 $\alpha = 97.348$ (2)°
 $\beta = 93.363$ (4)°

$\gamma = 102.410$ (7)°
 $V = 977.75$ (18) Å³
 $Z = 1$
 Mo $K\alpha$ radiation
 $\mu = 0.38$ mm⁻¹
 $T = 113$ K
 $0.50 \times 0.04 \times 0.04$ mm

Data collection

Rigaku Saturn724 CCD diffractometer
 Absorption correction: multi-scan (*CrystalClear*; Molecular Structure Corporation & Rigaku, 2005)
 $T_{\min} = 0.834$, $T_{\max} = 0.985$
 8262 measured reflections
 3418 independent reflections
 1459 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.106$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.062$
 $wR(F^2) = 0.131$
 $S = 0.87$
 3418 reflections
 283 parameters
 20 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.52$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.52$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{O1}-\text{H1}\cdots\text{N4}^{\text{i}}$	0.87 (2)	1.96 (3)	2.823 (5)	172 (5)
$\text{O2}-\text{H2}\cdots\text{O3}^{\text{ii}}$	0.86 (3)	1.94 (3)	2.772 (4)	165 (4)
$\text{O6}-\text{H6A}\cdots\text{O5}^{\text{iii}}$	0.99	1.83	2.817 (4)	174
$\text{O6}-\text{H6B}\cdots\text{N3}^{\text{iv}}$	0.99	2.23	3.009 (5)	135
$\text{O7}-\text{H7A}\cdots\text{O8}^{\text{v}}$	0.87 (2)	2.02 (3)	2.861 (5)	164 (4)
$\text{O7}-\text{H7B}\cdots\text{O2}^{\text{ii}}$	0.82 (2)	1.95 (2)	2.767 (4)	169 (5)
$\text{O8}-\text{H8A}\cdots\text{O5}^{\text{vi}}$	0.80 (2)	2.03 (3)	2.797 (4)	162 (4)
$\text{O8}-\text{H8B}\cdots\text{O3}^{\text{iii}}$	0.82 (2)	2.15 (3)	2.907 (5)	154 (4)
$\text{N5}-\text{H5}\cdots\text{O1}^{\text{iv}}$	0.90 (4)	2.01 (4)	2.828 (5)	151 (4)

Symmetry codes: (i) $x, y-1, z$; (ii) $-x+1, -y+1, -z$; (iii) $x+1, y, z$; (iv) $-x+2, -y+2, -z$; (v) $-x+2, -y+1, -z+1$; (vi) $-x+2, -y+2, -z+1$.

Data collection: *CrystalClear* (Molecular Structure Corporation & Rigaku, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *CrystalClear*.

This work was supported financially by the Research Center for Engineering Technology of Polymeric Composites of Shanxi Province, College of Materials Science and Engineering, North University of China,

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK5096).

References

- Candiani, G. & Frigerio, M. (2007). *Chem. Med. Chem.* **2**, 292–296.
- Hollink, E., Simanek, E. E. & Bergbreiter, D. E. (2005). *Tetrahedron Lett.* **46**, 2005–2008.
- Konstantion, T. N. & Petrova, P. (2002). *Dyes Pigm.* **52**, 115–120.
- Molecular Structure Corporation & Rigaku (2005). *CrystalClear*. MSC, The Woodlands, Texas, USA, and Rigaku Corporation, Tokyo, Japan.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supporting information

Acta Cryst. (2012). E68, m847–m848 [doi:10.1107/S1600536812023732]

Di- μ -aqua-bis{diaqua[μ -4-({4-[bis(2-hydroxyethyl)amino]-6-chloro-1,3,5-triazin-2-yl}amino)benzenesulfonato]sodium(I)}

Mei Zhang, Fu-Yu Sun, Gui-Zhe Zhao and Ya-Qing Liu

S1. Comment

Cyanuric chloride derivatives are widely used in commercial chemicals, especially in pesticides, reactive dyes, fluorescent brighteners, liposome and polymer photostabilizers (Hollink *et al.*, 2005; Candiani & Frigerio, 2007). The widespread use of these compounds is due to their higher reactive activity (Konstantion & Petrova, 2002). The title compound belongs to the cyanuric chloride derivatives and its structure is reported herein.


The asymmetric unit is shown in Fig. 1. The dihedral angle between the benzene ring and the triazine ring is $8.6(2)^\circ$. As shown in Fig. 2, the crystal packing displays O—H \cdots O, O—H \cdots N and N—H \cdots O hydrogen bonds, Table 1.

S2. Experimental


Cyanuric chloride (0.1 mol) was dissolved in acetone (120 ml). This solution was poured into distilled water (150 ml) with crushed ice (150 g). The reaction system was stirred maintaining the temperature at 0–5 °C in the ice-bath. An aqueous solution of sodium sulfate (0.1 mol) was slowly dropped into the above reaction vessel within 0.5 h, and then a 20% aqueous solution of sodium carbonate was added drop-wise to the reaction mixture to keep the pH at 7–8. Then the mixture was kept stirring for 5 h at 0–5 °C. After the reaction was completed, the white precipitate was filtered, washed with acetone and water twice, respectively, and dried at room temperature under vacuum to constant weight. This white powder is intermediate I. The intermediate I (0.05 mol) and a mixed solution of water and acetone (160 ml) were added into 250 ml four-neck flask. After stirring for 0.5 h at 25 °C, an aqueous solution of diethanol amine (0.06 mol) was slowly dropped into the reaction vessel within 0.5 h, and then a 20% aqueous solution of sodium carbonate was added drop-wise to the reaction mixture to keep the pH at 8–9. Then the mixture was kept stirring for 6 h at 45 °C. After the reaction was completed, the solution was rotary evaporated, washed with anhydrous alcohol twice, and dried at room temperature under vacuum to constant weight. The target product was obtained. Crystals of the title compound were obtained by slow evaporation of its methanol/*n*-hexane solution held at room temperature.

S3. Refinement

Carbon-bound H-atoms were placed in calculated positions [$C-H = 0.95$ to 0.99 Å, $U_{iso}(H) = 1.2U_{eq}(C)$] and were included in the refinement in the riding model approximation. The positions of the O- and N- bound H-atoms were refined with light distance restraints, and with $U_{iso}(H) = 1.2U_{eq}(\text{water-O,N})$ and $U_{iso}(H) = 1.5U_{eq}(\text{hydroxyl-O})$.

**Figure 1**

The molecular structure of (I). Displacement ellipsoids are drawn at the 30% probability level. H atoms have been omitted.

**Figure 2**

A view of the crystal packing for (I). Hydrogen bonds are shown as dashed lines.

Di- μ -aqua-bis{diaqua[μ -4-({4-[bis(2-hydroxyethyl)amino]-6-chloro-1,3,5-triazin-2-yl}amino)benzenesulfonato]sodium(I)}

Crystal data

[Na₂(C₁₃H₁₅ClN₅O₅S)₂(H₂O)₆]

$M_r = 931.72$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 7.5628$ (7) Å

$b = 8.6274$ (8) Å

$c = 15.532$ (2) Å

$\alpha = 97.348$ (2)°

$\beta = 93.363$ (4)°

$\gamma = 102.410$ (7)°

$V = 977.75$ (18) Å³

$Z = 1$

$F(000) = 484$

$D_x = 1.582$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 3065 reflections

$\theta = 2.4$ – 26.1 °

$\mu = 0.38$ mm⁻¹

$T = 113$ K

Prism, colourless

$0.50 \times 0.04 \times 0.04$ mm

Data collection

Rigaku Saturn724 CCD

diffractometer

Radiation source: rotating anode

Multilayer monochromator

Detector resolution: 14.22 pixels mm⁻¹

ω and φ scans

Absorption correction: multi-scan

(*CrystalClear*; Molecular Structure Corporation & Rigaku, 2005)

$T_{\min} = 0.834$, $T_{\max} = 0.985$

8262 measured reflections

3418 independent reflections

1459 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.106$

$\theta_{\max} = 25.0$ °, $\theta_{\min} = 2.4$ °

$h = -8 \rightarrow 8$

$k = -10 \rightarrow 10$

$l = -18 \rightarrow 18$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.062$

$wR(F^2) = 0.131$

$S = 0.87$

3418 reflections

283 parameters

20 restraints

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0293P)^2]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 0.52$ e Å⁻³

$\Delta\rho_{\min} = -0.52$ e Å⁻³

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Na1	0.9529 (2)	0.8060 (2)	0.46061 (12)	0.0216 (5)

S1	0.55329 (16)	0.91677 (15)	0.34804 (8)	0.0191 (3)
C11	1.02875 (15)	1.27203 (14)	-0.25219 (8)	0.0221 (4)
O1	0.9492 (4)	0.5459 (4)	-0.0875 (2)	0.0213 (9)
H1	0.934 (6)	0.443 (3)	-0.099 (3)	0.032*
O2	0.4874 (4)	0.5163 (4)	-0.3219 (2)	0.0234 (9)
H2	0.504 (6)	0.421 (3)	-0.330 (3)	0.035*
O3	0.4056 (4)	0.7718 (3)	0.32366 (19)	0.0210 (8)
O4	0.7162 (4)	0.8814 (4)	0.3889 (2)	0.0216 (9)
O5	0.4936 (4)	1.0469 (4)	0.40039 (19)	0.0194 (8)
N1	0.7584 (5)	0.7287 (4)	-0.1875 (2)	0.0161 (10)
N2	0.7670 (5)	0.9391 (4)	-0.0807 (2)	0.0162 (10)
N3	0.8817 (5)	0.9841 (4)	-0.2196 (2)	0.0150 (9)
N4	0.8923 (5)	1.2084 (4)	-0.1092 (2)	0.0175 (10)
N5	0.7900 (5)	1.1689 (5)	0.0233 (3)	0.0170 (10)
H5	0.851 (5)	1.272 (5)	0.029 (3)	0.020*
C1	0.7906 (5)	0.5903 (5)	-0.0571 (3)	0.0195 (12)
H1A	0.8270	0.6892	-0.0142	0.023*
H1B	0.7234	0.5042	-0.0271	0.023*
C2	0.6672 (6)	0.6188 (5)	-0.1302 (3)	0.0207 (12)
H2A	0.6108	0.5145	-0.1660	0.025*
H2B	0.5682	0.6632	-0.1048	0.025*
C3	0.8018 (6)	0.6556 (5)	-0.2719 (3)	0.0192 (12)
H3A	0.8331	0.5519	-0.2650	0.023*
H3B	0.9100	0.7264	-0.2901	0.023*
C4	0.6466 (6)	0.6270 (5)	-0.3431 (3)	0.0227 (13)
H4A	0.6151	0.7303	-0.3507	0.027*
H4B	0.6862	0.5828	-0.3989	0.027*
C5	0.8033 (6)	0.8880 (6)	-0.1625 (3)	0.0189 (12)
C6	0.9213 (6)	1.1378 (5)	-0.1863 (3)	0.0169 (12)
C7	0.8149 (6)	1.0990 (6)	-0.0576 (3)	0.0158 (12)
C8	0.7265 (6)	1.0979 (5)	0.0957 (3)	0.0130 (11)
C9	0.7404 (6)	1.2041 (5)	0.1720 (3)	0.0180 (12)
H9	0.7876	1.3154	0.1720	0.022*
C10	0.6857 (6)	1.1488 (5)	0.2486 (3)	0.0180 (12)
H10	0.6997	1.2221	0.3010	0.022*
C11	0.6109 (6)	0.9878 (5)	0.2493 (3)	0.0149 (11)
C12	0.5925 (6)	0.8800 (6)	0.1722 (3)	0.0203 (12)
H12	0.5421	0.7691	0.1720	0.024*
C13	0.6486 (5)	0.9361 (5)	0.0955 (3)	0.0166 (11)
H13	0.6335	0.8634	0.0428	0.020*
O6	1.1240 (4)	1.0471 (3)	0.41494 (19)	0.0191 (8)
H6A	1.2558	1.0514	0.4142	0.023*
H6B	1.0732	1.0718	0.3594	0.023*
O7	0.8355 (4)	0.5359 (4)	0.4260 (2)	0.0349 (11)
H7A	0.825 (6)	0.466 (5)	0.462 (2)	0.042*
H7B	0.747 (5)	0.518 (5)	0.390 (2)	0.042*
O8	1.2365 (4)	0.7394 (4)	0.4857 (2)	0.0226 (9)
H8A	1.297 (5)	0.813 (4)	0.518 (2)	0.027*

H8B 1.267 (5) 0.717 (5) 0.4365 (17) 0.027*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Na1	0.0141 (10)	0.0229 (12)	0.0293 (12)	0.0041 (9)	0.0094 (9)	0.0061 (9)
S1	0.0148 (7)	0.0226 (8)	0.0217 (8)	0.0051 (6)	0.0093 (6)	0.0052 (6)
C11	0.0211 (7)	0.0231 (8)	0.0241 (8)	0.0047 (6)	0.0125 (6)	0.0067 (6)
O1	0.0155 (18)	0.0163 (19)	0.035 (2)	0.0071 (17)	0.0074 (16)	0.0067 (18)
O2	0.0204 (19)	0.019 (2)	0.028 (2)	-0.0016 (18)	0.0032 (16)	0.0022 (18)
O3	0.0175 (18)	0.020 (2)	0.024 (2)	-0.0023 (16)	0.0067 (16)	0.0032 (16)
O4	0.0116 (17)	0.033 (2)	0.026 (2)	0.0147 (16)	0.0025 (15)	0.0102 (16)
O5	0.0145 (18)	0.024 (2)	0.022 (2)	0.0094 (15)	0.0118 (15)	0.0004 (15)
N1	0.018 (2)	0.016 (2)	0.014 (2)	0.0025 (19)	0.0096 (18)	0.0017 (18)
N2	0.013 (2)	0.021 (2)	0.015 (2)	0.0022 (19)	0.0068 (18)	0.0026 (19)
N3	0.012 (2)	0.015 (2)	0.018 (2)	0.0017 (18)	0.0083 (18)	0.0018 (18)
N4	0.015 (2)	0.022 (3)	0.016 (2)	0.002 (2)	0.0080 (19)	0.0045 (19)
N5	0.016 (2)	0.018 (2)	0.016 (2)	-0.0009 (19)	0.0084 (19)	0.004 (2)
C1	0.015 (3)	0.016 (3)	0.030 (3)	0.004 (2)	0.008 (2)	0.010 (2)
C2	0.023 (3)	0.016 (3)	0.026 (3)	0.008 (2)	0.009 (2)	0.004 (2)
C3	0.019 (3)	0.019 (3)	0.019 (3)	0.004 (2)	0.009 (2)	0.000 (2)
C4	0.028 (3)	0.018 (3)	0.023 (3)	0.005 (3)	0.005 (2)	0.006 (2)
C5	0.009 (3)	0.023 (3)	0.025 (3)	0.002 (2)	0.002 (2)	0.005 (2)
C6	0.015 (3)	0.020 (3)	0.020 (3)	0.009 (2)	0.004 (2)	0.007 (2)
C7	0.008 (3)	0.023 (3)	0.018 (3)	0.007 (2)	0.002 (2)	0.004 (2)
C8	0.009 (3)	0.018 (3)	0.014 (3)	0.007 (2)	0.007 (2)	0.003 (2)
C9	0.016 (3)	0.015 (3)	0.024 (3)	0.003 (2)	0.010 (2)	0.001 (2)
C10	0.017 (3)	0.018 (3)	0.020 (3)	0.004 (2)	0.013 (2)	0.002 (2)
C11	0.011 (2)	0.017 (3)	0.020 (3)	0.009 (2)	0.010 (2)	0.002 (2)
C12	0.016 (3)	0.018 (3)	0.028 (3)	0.004 (2)	0.005 (2)	0.005 (2)
C13	0.013 (3)	0.021 (3)	0.017 (3)	0.004 (2)	0.007 (2)	0.003 (2)
O6	0.0091 (17)	0.030 (2)	0.021 (2)	0.0056 (15)	0.0023 (15)	0.0114 (16)
O7	0.021 (2)	0.034 (3)	0.048 (3)	-0.0018 (19)	-0.0035 (19)	0.017 (2)
O8	0.018 (2)	0.026 (2)	0.022 (2)	0.0009 (17)	0.0065 (17)	0.0023 (18)

Geometric parameters (Å, °)

Na1—O7	2.290 (4)	N5—H5	0.90 (4)
Na1—O4	2.306 (3)	C1—C2	1.505 (5)
Na1—O6 ⁱ	2.346 (4)	C1—H1A	0.9900
Na1—O8	2.361 (4)	C1—H1B	0.9900
Na1—O6	2.415 (3)	C2—H2A	0.9900
Na1—C11 ⁱⁱ	3.239 (2)	C2—H2B	0.9900
Na1—Na1 ⁱ	3.322 (4)	C3—C4	1.522 (5)
Na1—H8B	2.68 (4)	C3—H3A	0.9900
S1—O4	1.461 (3)	C3—H3B	0.9900
S1—O5	1.467 (3)	C4—H4A	0.9900
S1—O3	1.478 (3)	C4—H4B	0.9900

S1—C11	1.766 (5)	C8—C9	1.386 (6)
Cl1—C6	1.746 (5)	C8—C13	1.393 (6)
Cl1—Na1 ⁱⁱ	3.238 (2)	C9—C10	1.391 (6)
O1—C1	1.423 (5)	C9—H9	0.9500
O1—H1	0.87 (2)	C10—C11	1.384 (6)
O2—C4	1.451 (5)	C10—H10	0.9500
O2—H2	0.86 (3)	C11—C12	1.400 (6)
N1—C5	1.343 (5)	C12—C13	1.397 (6)
N1—C3	1.467 (5)	C12—H12	0.9500
N1—C2	1.474 (5)	C13—H13	0.9500
N2—C7	1.344 (5)	O6—Na1 ⁱ	2.346 (4)
N2—C5	1.354 (6)	O6—H6A	0.9900
N3—C6	1.323 (5)	O6—H6B	0.9900
N3—C5	1.365 (6)	O7—H7A	0.87 (2)
N4—C6	1.324 (5)	O7—H7B	0.82 (2)
N4—C7	1.374 (5)	O8—H8A	0.80 (2)
N5—C7	1.364 (6)	O8—H8B	0.82 (2)
N5—C8	1.408 (5)		
O7—Na1—O4	95.45 (13)	C1—C2—H2A	108.7
O7—Na1—O6 ⁱ	121.20 (14)	N1—C2—H2B	108.7
O4—Na1—O6 ⁱ	85.08 (12)	C1—C2—H2B	108.7
O7—Na1—O8	86.24 (13)	H2A—C2—H2B	107.6
O4—Na1—O8	159.53 (15)	N1—C3—C4	113.2 (4)
O6 ⁱ —Na1—O8	111.51 (12)	N1—C3—H3A	108.9
O7—Na1—O6	146.92 (15)	C4—C3—H3A	108.9
O4—Na1—O6	81.63 (11)	N1—C3—H3B	108.9
O6 ⁱ —Na1—O6	91.52 (12)	C4—C3—H3B	108.9
O8—Na1—O6	85.87 (12)	H3A—C3—H3B	107.7
O7—Na1—Cl1 ⁱⁱ	74.72 (11)	O2—C4—C3	111.2 (4)
O4—Na1—Cl1 ⁱⁱ	70.57 (9)	O2—C4—H4A	109.4
O6 ⁱ —Na1—Cl1 ⁱⁱ	152.67 (10)	C3—C4—H4A	109.4
O8—Na1—Cl1 ⁱⁱ	90.36 (10)	O2—C4—H4B	109.4
O6—Na1—Cl1 ⁱⁱ	73.26 (9)	C3—C4—H4B	109.4
O7—Na1—Na1 ⁱ	167.15 (14)	H4A—C4—H4B	108.0
O4—Na1—Na1 ⁱ	80.42 (10)	N1—C5—N2	115.8 (5)
O6 ⁱ —Na1—Na1 ⁱ	46.62 (8)	N1—C5—N3	118.7 (5)
O8—Na1—Na1 ⁱ	101.92 (11)	N2—C5—N3	125.5 (4)
O6—Na1—Na1 ⁱ	44.91 (8)	N3—C6—N4	130.0 (5)
Cl1 ⁱⁱ —Na1—Na1 ⁱ	114.70 (8)	N3—C6—Cl1	116.6 (4)
O7—Na1—H8B	82.4 (8)	N4—C6—Cl1	113.4 (3)
O4—Na1—H8B	142.8 (7)	N2—C7—N5	121.3 (5)
O6 ⁱ —Na1—H8B	127.8 (6)	N2—C7—N4	125.9 (5)
O8—Na1—H8B	17.3 (6)	N5—C7—N4	112.8 (4)
O6—Na1—H8B	80.6 (9)	C9—C8—C13	119.3 (4)
Cl1 ⁱⁱ —Na1—H8B	73.1 (6)	C9—C8—N5	114.7 (4)
Na1 ⁱ —Na1—H8B	108.3 (8)	C13—C8—N5	126.0 (4)
O4—S1—O5	112.25 (18)	C8—C9—C10	120.4 (4)

O4—S1—O3	112.36 (19)	C8—C9—H9	119.8
O5—S1—O3	112.56 (18)	C10—C9—H9	119.8
O4—S1—C11	107.4 (2)	C11—C10—C9	120.6 (4)
O5—S1—C11	105.8 (2)	C11—C10—H10	119.7
O3—S1—C11	105.9 (2)	C9—C10—H10	119.7
C6—C11—Na1 ⁱⁱ	125.53 (17)	C10—C11—C12	119.4 (4)
C1—O1—H1	112 (3)	C10—C11—S1	120.3 (4)
C4—O2—H2	110 (3)	C12—C11—S1	120.2 (4)
S1—O4—Na1	173.9 (2)	C13—C12—C11	119.7 (4)
C5—N1—C3	121.7 (4)	C13—C12—H12	120.1
C5—N1—C2	121.4 (4)	C11—C12—H12	120.1
C3—N1—C2	116.9 (4)	C8—C13—C12	120.5 (4)
C7—N2—C5	114.2 (4)	C8—C13—H13	119.8
C6—N3—C5	112.4 (4)	C12—C13—H13	119.8
C6—N4—C7	111.9 (4)	Na1 ⁱ —O6—Na1	88.48 (12)
C7—N5—C8	129.8 (4)	Na1 ⁱ —O6—H6A	113.9
C7—N5—H5	106 (3)	Na1—O6—H6A	113.9
C8—N5—H5	122 (3)	Na1 ⁱ —O6—H6B	113.9
O1—C1—C2	112.1 (4)	Na1—O6—H6B	113.9
O1—C1—H1A	109.2	H6A—O6—H6B	111.1
C2—C1—H1A	109.2	Na1—O7—H7A	126 (3)
O1—C1—H1B	109.2	Na1—O7—H7B	111 (3)
C2—C1—H1B	109.2	H7A—O7—H7B	112 (3)
H1A—C1—H1B	107.9	Na1—O8—H8A	106 (3)
N1—C2—C1	114.4 (4)	Na1—O8—H8B	104 (3)
N1—C2—H2A	108.7	H8A—O8—H8B	120 (3)
O5—S1—O4—Na1	90 (2)	C5—N2—C7—N5	-178.9 (4)
O3—S1—O4—Na1	-38 (2)	C5—N2—C7—N4	0.8 (6)
C11—S1—O4—Na1	-154 (2)	C8—N5—C7—N2	4.3 (7)
O7—Na1—O4—S1	48 (2)	C8—N5—C7—N4	-175.4 (4)
O6 ⁱ —Na1—O4—S1	-73 (2)	C6—N4—C7—N2	-1.9 (6)
O8—Na1—O4—S1	141.4 (19)	C6—N4—C7—N5	177.8 (4)
O6—Na1—O4—S1	-166 (2)	C7—N5—C8—C9	171.8 (4)
C11 ⁱⁱ —Na1—O4—S1	119 (2)	C7—N5—C8—C13	-10.3 (7)
Na1 ⁱ —Na1—O4—S1	-120 (2)	C13—C8—C9—C10	3.2 (7)
C5—N1—C2—C1	-76.6 (5)	N5—C8—C9—C10	-178.8 (4)
C3—N1—C2—C1	101.9 (4)	C8—C9—C10—C11	-2.2 (7)
O1—C1—C2—N1	-50.9 (5)	C9—C10—C11—C12	0.7 (6)
C5—N1—C3—C4	-94.4 (5)	C9—C10—C11—S1	176.5 (3)
C2—N1—C3—C4	87.0 (5)	O4—S1—C11—C10	-86.3 (4)
N1—C3—C4—O2	-61.9 (5)	O5—S1—C11—C10	33.7 (4)
C3—N1—C5—N2	-175.5 (4)	O3—S1—C11—C10	153.4 (3)
C2—N1—C5—N2	2.9 (6)	O4—S1—C11—C12	89.4 (4)
C3—N1—C5—N3	3.9 (6)	O5—S1—C11—C12	-150.5 (4)
C2—N1—C5—N3	-177.7 (3)	O3—S1—C11—C12	-30.8 (4)
C7—N2—C5—N1	179.5 (4)	C10—C11—C12—C13	-0.4 (7)
C7—N2—C5—N3	0.1 (7)	S1—C11—C12—C13	-176.2 (3)

C6—N3—C5—N1	-179.1 (4)	C9—C8—C13—C12	-2.9 (6)
C6—N3—C5—N2	0.3 (7)	N5—C8—C13—C12	179.3 (4)
C5—N3—C6—N4	-1.8 (7)	C11—C12—C13—C8	1.5 (7)
C5—N3—C6—C11	178.1 (3)	O7—Na1—O6—Na1 ⁱ	171.9 (2)
C7—N4—C6—N3	2.6 (7)	O4—Na1—O6—Na1 ⁱ	84.80 (11)
C7—N4—C6—C11	-177.4 (3)	O6 ⁱ —Na1—O6—Na1 ⁱ	0.0
Na1 ⁱⁱ —C11—C6—N3	16.5 (4)	O8—Na1—O6—Na1 ⁱ	-111.45 (12)
Na1 ⁱⁱ —C11—C6—N4	-163.6 (2)	C11 ⁱⁱ —Na1—O6—Na1 ⁱ	156.93 (10)

Symmetry codes: (i) $-x+2, -y+2, -z+1$; (ii) $-x+2, -y+2, -z$.

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
O1—H1...N4 ⁱⁱⁱ	0.87 (2)	1.96 (3)	2.823 (5)	172 (5)
O2—H2...O3 ^{iv}	0.86 (3)	1.94 (3)	2.772 (4)	165 (4)
O6—H6A...O5 ^v	0.99	1.83	2.817 (4)	174
O6—H6B...N3 ⁱⁱ	0.99	2.23	3.009 (5)	135
O7—H7A...O8 ^{vi}	0.87 (2)	2.02 (3)	2.861 (5)	164 (4)
O7—H7B...O2 ^{iv}	0.82 (2)	1.95 (2)	2.767 (4)	169 (5)
O8—H8A...O5 ⁱ	0.80 (2)	2.03 (3)	2.797 (4)	162 (4)
O8—H8B...O3 ^v	0.82 (2)	2.15 (3)	2.907 (5)	154 (4)
N5—H5...O1 ⁱⁱ	0.90 (4)	2.01 (4)	2.828 (5)	151 (4)

Symmetry codes: (i) $-x+2, -y+2, -z+1$; (ii) $-x+2, -y+2, -z$; (iii) $x, y-1, z$; (iv) $-x+1, -y+1, -z$; (v) $x+1, y, z$; (vi) $-x+2, -y+1, -z+1$.